



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:10 AM UTC

PDB ID : 6AWI / pdb_00006awi
Title : Staphylococcus aureus Type II pantothenate kinase in complex with ADP and pantothenate analog Deoxy-N5Pan
Authors : Chen, Y.; Antoshchenko, T.; Strauss, E.; Barnard, L.; Huang, Y.H.
Deposited on : 2017-09-05
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	266	3% 85% 13% .
1	B	266	2% 86% 13% .
1	C	266	5% 84% 14% .
1	D	266	8% 86% 12% .

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8889 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type II pantothenate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	266	Total	C	N	O	S	0	0	0
			2041	1292	350	393	6			
1	B	266	Total	C	N	O	S	0	0	0
			2041	1292	350	393	6			
1	C	265	Total	C	N	O	S	0	0	0
			2032	1287	349	390	6			
1	D	266	Total	C	N	O	S	0	0	0
			2041	1292	350	393	6			

- Molecule 2 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
2	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
2	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 3 is (2R)-2-hydroxy-3,3-dimethyl-N-[3-oxo-3-(pentylamino)propyl]butanamide (CCD ID: C0D) (formula: $C_{14}H_{28}N_2O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			19	14	2	3		
3	B	1	Total	C	N	O	0	0
			19	14	2	3		
3	C	1	Total	C	N	O	0	0
			19	14	2	3		
3	D	1	Total	C	N	O	0	0
			19	14	2	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	148	Total	O	0	0
			148	148		
4	B	179	Total	O	0	0
			179	179		
4	C	117	Total	O	0	0
			117	117		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	106	Total 106	O 106	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type II pantothenate kinase




- Molecule 1: Type II pantothenate kinase




- Molecule 1: Type II pantothenate kinase



- Molecule 1: Type II pantothenate kinase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.39Å 57.04Å 134.08Å 90.00° 99.88° 90.00°	Depositor
Resolution (Å)	50.00 – 1.80 50.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.8 (50.00-1.80) 99.7 (50.00-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.95 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.188 , 0.233 0.202 , 0.245	Depositor DCC
R_{free} test set	4985 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.163	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8889	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 14.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: C0D, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.42	9/2076 (0.4%)	1.19	5/2813 (0.2%)
1	B	1.36	11/2076 (0.5%)	1.17	7/2813 (0.2%)
1	C	1.32	6/2067 (0.3%)	1.19	7/2801 (0.2%)
1	D	1.19	3/2076 (0.1%)	1.14	3/2813 (0.1%)
All	All	1.32	29/8295 (0.3%)	1.17	22/11240 (0.2%)

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	240	TYR	C-O	-7.99	1.14	1.24
1	D	137	TYR	C-O	-7.89	1.14	1.24
1	A	145	GLN	C-O	-7.72	1.14	1.24
1	A	214	GLU	N-CA	-7.10	1.37	1.46
1	C	110	SER	C-O	-6.55	1.16	1.23
1	A	21	GLN	C-O	-6.49	1.16	1.24
1	A	260	ILE	CA-C	-6.37	1.45	1.52
1	A	144	ALA	C-O	-6.25	1.16	1.24
1	B	233	LEU	C-O	-6.02	1.17	1.24
1	B	90	ALA	C-O	-5.88	1.17	1.24
1	C	234	ARG	C-O	-5.88	1.17	1.24
1	A	142	ASP	C-O	-5.83	1.17	1.24
1	A	141	THR	C-O	-5.69	1.17	1.24
1	B	73	ALA	C-O	-5.64	1.17	1.24
1	B	21	GLN	CA-C	5.62	1.59	1.52
1	B	74	ALA	N-CA	5.62	1.53	1.46
1	B	71	PHE	C-O	-5.59	1.17	1.24
1	B	200	VAL	CA-C	5.59	1.59	1.52
1	B	78	LEU	N-CA	5.53	1.53	1.46
1	A	10	THR	CA-C	5.47	1.58	1.52
1	B	232	LEU	C-O	-5.45	1.17	1.24
1	B	72	ASP	C-O	-5.38	1.17	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	112	ARG	C-O	-5.34	1.18	1.23
1	B	234	ARG	C-O	-5.34	1.17	1.24
1	A	123	MET	N-CA	-5.32	1.39	1.46
1	D	170	ASP	CA-C	5.31	1.60	1.52
1	C	200	VAL	N-CA	-5.24	1.40	1.46
1	C	46	LYS	C-O	-5.20	1.18	1.23
1	C	236	VAL	CA-CB	5.13	1.60	1.54

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	184	ASP	N-CA-C	-8.18	101.95	111.03
1	B	70	GLU	N-CA-C	7.08	119.89	111.33
1	B	150	ASN	N-CA-C	6.42	119.10	111.33
1	D	150	ASN	N-CA-C	6.11	117.94	111.28
1	D	128	GLY	N-CA-C	-6.04	105.48	112.73
1	C	150	ASN	N-CA-C	5.69	118.25	111.71
1	B	127	LEU	N-CA-C	-5.63	105.29	111.82
1	C	21	GLN	N-CA-C	-5.56	100.18	109.24
1	A	251	VAL	CB-CA-C	5.52	118.39	110.33
1	B	235	LYS	N-CA-C	5.47	117.25	111.28
1	A	184	ASP	N-CA-C	-5.37	105.60	111.82
1	A	260	ILE	CB-CA-C	-5.20	105.20	112.02
1	B	52	GLY	N-CA-C	5.19	119.17	112.83
1	C	191	ASN	N-CA-C	-5.11	105.79	111.36
1	A	31	ASN	N-CA-C	5.11	118.65	112.93
1	B	69	VAL	N-CA-CB	5.09	116.08	110.53
1	A	75	SER	N-CA-C	-5.09	105.73	111.28
1	C	185	ALA	CA-C-N	5.09	127.06	120.44
1	C	185	ALA	C-N-CA	5.09	127.06	120.44
1	D	41	GLN	N-CA-C	5.07	119.45	113.16
1	C	94	PHE	N-CA-CB	5.01	119.55	110.83
1	B	237	VAL	N-CA-C	5.01	115.24	110.53

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2041	0	2027	20	0
1	B	2041	0	2027	11	0
1	C	2032	0	2021	20	1
1	D	2041	0	2027	18	1
2	A	27	0	12	0	0
2	B	27	0	12	0	0
2	C	27	0	12	0	0
2	D	27	0	12	0	0
3	A	19	0	0	0	0
3	B	19	0	0	0	0
3	C	19	0	0	0	0
3	D	19	0	0	0	0
4	A	148	0	0	6	0
4	B	179	0	0	2	0
4	C	117	0	0	7	0
4	D	106	0	0	1	0
All	All	8889	0	8150	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:433:HOH:O	1:C:234:ARG:HD2	1.58	1.04
1:C:183:LEU:HD22	4:C:410:HOH:O	1.69	0.92
1:D:185:ALA:HB3	4:D:421:HOH:O	1.73	0.88
1:A:27:GLU:OE2	4:A:401:HOH:O	1.92	0.86
1:D:158:HIS:O	1:D:161:LYS:HE2	1.85	0.77
1:A:264:TYR:CE1	4:A:452:HOH:O	2.41	0.72
1:C:20:ASN:ND2	4:C:402:HOH:O	2.22	0.70
1:B:133:GLN:HG3	4:B:466:HOH:O	1.92	0.68
1:C:186:ASP:OD2	1:C:186:ASP:N	2.26	0.67
1:C:252:GLU:HB3	4:C:407:HOH:O	1.97	0.64
1:A:186:ASP:OD2	1:A:186:ASP:N	2.21	0.62
1:D:32:ILE:HD11	1:D:57:ILE:HG13	1.83	0.59
1:A:46:LYS:HG2	1:A:266:GLU:OE1	2.02	0.58
1:C:96:ASN:O	4:C:401:HOH:O	2.17	0.57
1:A:26:THR:HG22	1:A:256:PHE:HZ	1.70	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:114:VAL:HG11	1:C:215:PHE:CE2	2.39	0.56
1:C:84:GLU:HB3	4:C:403:HOH:O	2.05	0.55
1:D:17:GLU:OE1	1:D:264:TYR:OH	2.25	0.54
1:C:55:GLY:HA3	1:D:163:THR:HG21	1.90	0.53
1:D:63:ILE:HG22	1:D:64:PRO:HD2	1.90	0.52
1:C:180:LEU:HA	1:C:183:LEU:HD13	1.90	0.52
1:B:63:ILE:HB	1:B:64:PRO:HD2	1.92	0.52
1:D:19:ASP:C	1:D:21:GLN:H	2.18	0.52
1:A:161:LYS:HD2	1:B:53:ASN:OD1	2.10	0.51
1:A:45:GLU:O	1:A:46:LYS:HE2	2.11	0.51
4:A:526:HOH:O	1:C:21:GLN:HG3	2.10	0.51
1:A:187:PHE:O	1:A:192:LYS:HE2	2.11	0.51
1:A:161:LYS:HD3	1:B:56:VAL:CG2	2.41	0.50
1:A:21:GLN:HA	4:A:521:HOH:O	2.10	0.49
1:A:26:THR:HG22	1:A:256:PHE:CZ	2.47	0.49
1:C:253:ASN:ND2	4:C:407:HOH:O	2.47	0.48
1:D:144:ALA:HB1	1:D:197:ILE:HD12	1.96	0.47
1:C:186:ASP:O	1:C:187:PHE:HB2	2.15	0.47
1:A:182:HIS:O	1:A:185:ALA:HB3	2.15	0.46
1:D:184:ASP:OD1	1:D:184:ASP:C	2.58	0.46
1:D:209:ILE:C	1:D:209:ILE:HD12	2.40	0.46
1:D:25:LYS:HD3	1:D:26:THR:N	2.31	0.46
1:A:14:ILE:HD12	4:A:457:HOH:O	2.16	0.46
1:A:161:LYS:HD3	1:B:56:VAL:HG21	1.99	0.44
1:B:19:ASP:N	1:B:21:GLN:OE1	2.50	0.43
1:C:199:VAL:O	1:C:203:VAL:HG23	2.19	0.43
1:B:32:ILE:HD11	1:B:57:ILE:HG13	2.01	0.43
1:C:73:ALA:O	1:C:261:GLY:HA3	2.19	0.43
1:A:16:GLN:HB2	1:A:23:THR:HB	2.00	0.42
1:A:55:GLY:HA3	1:B:163:THR:HG21	2.00	0.42
1:C:98:GLY:N	4:C:401:HOH:O	2.51	0.42
1:D:19:ASP:O	1:D:20:ASN:HB2	2.17	0.42
1:D:17:GLU:OE1	1:D:264:TYR:CZ	2.73	0.42
1:D:77:GLY:HA3	1:D:257:SER:O	2.19	0.42
1:B:138:LYS:HG2	4:B:529:HOH:O	2.19	0.42
1:D:167:ILE:HB	1:D:168:PRO:HD2	2.02	0.42
1:D:73:ALA:O	1:D:261:GLY:HA3	2.19	0.42
1:C:27:GLU:HG2	1:C:35:VAL:CG2	2.50	0.41
1:B:209:ILE:C	1:B:209:ILE:HD12	2.44	0.41
1:A:182:HIS:O	1:A:185:ALA:N	2.53	0.41
1:B:144:ALA:CB	1:B:197:ILE:HD12	2.51	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:127:LEU:HD11	1:D:127:LEU:HD11	2.03	0.41
1:A:38:TRP:O	1:A:42:GLN:HG2	2.21	0.41
1:A:75:SER:HB2	1:A:111:GLN:HG2	2.02	0.41
1:A:143:MET:HG2	1:A:190:SER:OG	2.21	0.41
1:C:109:GLN:HE21	1:C:109:GLN:HB2	1.56	0.41
1:C:207:MET:SD	1:D:207:MET:SD	3.19	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:60:ASN:OD1	1:D:21:GLN:OE1[2_646]	1.46	0.74

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/266 (99%)	256 (97%)	8 (3%)	0	100	100
1	B	264/266 (99%)	255 (97%)	9 (3%)	0	100	100
1	C	263/266 (99%)	253 (96%)	10 (4%)	0	100	100
1	D	264/266 (99%)	253 (96%)	10 (4%)	1 (0%)	30	19
All	All	1055/1064 (99%)	1017 (96%)	37 (4%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	161	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	216/216 (100%)	208 (96%)	8 (4%)	30	18
1	B	216/216 (100%)	209 (97%)	7 (3%)	34	22
1	C	215/216 (100%)	204 (95%)	11 (5%)	21	9
1	D	216/216 (100%)	205 (95%)	11 (5%)	21	9
All	All	863/864 (100%)	826 (96%)	37 (4%)	26	13

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	THR
1	A	30	LYS
1	A	43	GLN
1	A	112	ARG
1	A	138	LYS
1	A	178	HIS
1	A	251	VAL
1	A	266	GLU
1	B	1	MET
1	B	19	ASP
1	B	21	GLN
1	B	26	THR
1	B	164	GLU
1	B	178	HIS
1	B	266	GLU
1	C	20	ASN
1	C	25	LYS
1	C	41	GLN
1	C	88	ASP
1	C	109	GLN
1	C	161	LYS
1	C	178	HIS
1	C	183	LEU
1	C	186	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	243	LEU
1	C	244	ARG
1	D	20	ASN
1	D	25	LYS
1	D	26	THR
1	D	27	GLU
1	D	43	GLN
1	D	62	ASN
1	D	63	ILE
1	D	112	ARG
1	D	161	LYS
1	D	178	HIS
1	D	184	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	ASN
1	A	66	GLN
1	A	96	ASN
1	A	219	ASN
1	B	16	GLN
1	B	18	GLN
1	B	41	GLN
1	B	66	GLN
1	B	125	GLN
1	B	133	GLN
1	B	181	HIS
1	B	219	ASN
1	C	20	ASN
1	C	42	GLN
1	C	109	GLN
1	C	133	GLN
1	C	253	ASN
1	D	21	GLN
1	D	62	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	C0D	A	302	-	17,18,18	1.59	2 (11%)	22,23,23	0.79	1 (4%)
2	ADP	A	301	-	28,29,29	1.69	9 (32%)	43,45,45	2.39	10 (23%)
3	C0D	D	302	-	17,18,18	0.86	0	22,23,23	1.21	2 (9%)
3	C0D	B	302	-	17,18,18	2.92	2 (11%)	22,23,23	1.73	6 (27%)
2	ADP	C	301	-	28,29,29	1.31	2 (7%)	43,45,45	2.14	14 (32%)
3	C0D	C	302	-	17,18,18	1.02	1 (5%)	22,23,23	1.01	1 (4%)
2	ADP	D	301	-	28,29,29	1.37	5 (17%)	43,45,45	1.96	10 (23%)
2	ADP	B	301	-	28,29,29	1.20	2 (7%)	43,45,45	2.14	8 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	C0D	A	302	-	-	2/22/22/22	-
2	ADP	A	301	-	-	3/16/32/32	0/3/3/3
3	C0D	D	302	-	-	0/22/22/22	-
3	C0D	B	302	-	-	1/22/22/22	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ADP	C	301	-	-	4/16/32/32	0/3/3/3
3	C0D	C	302	-	-	1/22/22/22	-
2	ADP	D	301	-	-	2/16/32/32	0/3/3/3
2	ADP	B	301	-	-	2/16/32/32	0/3/3/3

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	302	C0D	CAW-CAV	11.29	1.62	1.55
3	A	302	C0D	CAW-CAV	5.59	1.58	1.55
2	A	301	ADP	PA-O3A	3.86	1.63	1.59
2	D	301	ADP	C5-C4	3.77	1.45	1.39
2	C	301	ADP	C5-C4	3.73	1.45	1.39
2	A	301	ADP	C5-N7	-3.59	1.32	1.39
2	C	301	ADP	PA-O3A	3.53	1.63	1.59
2	A	301	ADP	C5-C4	3.52	1.45	1.39
2	B	301	ADP	C8-N7	3.04	1.37	1.31
2	D	301	ADP	C4-N9	-2.72	1.32	1.37
2	D	301	ADP	C8-N7	2.51	1.36	1.31
3	C	302	C0D	CAW-CAV	2.49	1.56	1.55
2	D	301	ADP	C2'-C3'	-2.38	1.46	1.53
3	A	302	C0D	CAB-CAW	2.28	1.58	1.53
3	B	302	C0D	OAG-CAV	2.25	1.46	1.42
2	A	301	ADP	C4-N9	-2.21	1.33	1.37
2	D	301	ADP	C5-C6	2.16	1.47	1.41
2	A	301	ADP	C1'-N9	2.16	1.52	1.46
2	A	301	ADP	C8-N9	-2.15	1.33	1.37
2	B	301	ADP	C5-C4	2.13	1.42	1.39
2	A	301	ADP	O3'-C3'	2.05	1.48	1.43
2	A	301	ADP	C5-C6	2.03	1.46	1.41
2	A	301	ADP	O4'-C4'	-2.00	1.40	1.45

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	ADP	N3-C2-N1	-7.13	117.80	128.58
2	A	301	ADP	C5-C4-N3	-6.82	117.33	126.72
2	B	301	ADP	C5-C4-N3	-6.78	117.38	126.72
2	B	301	ADP	N3-C2-N1	-6.70	118.44	128.58
2	D	301	ADP	C5-C4-N3	-6.25	118.12	126.72
2	C	301	ADP	C5-C4-N3	-6.15	118.25	126.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	ADP	C2-N3-C4	5.80	125.99	111.83
2	B	301	ADP	C2-N3-C4	5.42	125.08	111.83
2	A	301	ADP	N3-C4-N9	5.28	136.14	127.17
2	C	301	ADP	N3-C4-N9	5.25	136.09	127.17
2	B	301	ADP	N3-C4-N9	4.90	135.50	127.17
2	C	301	ADP	N3-C2-N1	-4.52	121.74	128.58
3	B	302	C0D	CAN-CAO-CAT	-4.39	105.08	112.39
2	D	301	ADP	N3-C4-N9	4.38	134.61	127.17
2	D	301	ADP	C2-N3-C4	4.20	122.09	111.83
2	C	301	ADP	C2-N3-C4	4.14	121.94	111.83
2	D	301	ADP	N3-C2-N1	-4.07	122.43	128.58
2	D	301	ADP	C4-C5-N7	-3.38	106.72	110.58
2	C	301	ADP	C5-N7-C8	3.34	108.69	103.45
2	C	301	ADP	C4-N9-C8	3.33	109.23	105.74
2	C	301	ADP	N9-C8-N7	-3.17	109.44	113.94
2	A	301	ADP	C2-N1-C6	3.13	123.88	118.73
2	C	301	ADP	C4-C5-N7	-3.12	107.02	110.58
3	B	302	C0D	OAE-CAU-CAV	2.94	129.10	120.89
2	D	301	ADP	O4'-C1'-N9	-2.92	102.49	108.09
3	D	302	C0D	CAN-CAO-CAT	-2.78	107.76	112.39
3	B	302	C0D	OAE-CAU-NAR	-2.72	117.22	122.98
2	D	301	ADP	C4-N9-C8	2.72	108.59	105.74
2	A	301	ADP	C4-C5-N7	-2.69	107.50	110.58
2	C	301	ADP	C1'-N9-C8	-2.68	121.15	127.09
2	A	301	ADP	C6-C5-N7	2.57	137.04	132.09
3	C	302	C0D	CAN-CAO-CAT	-2.55	108.14	112.39
2	C	301	ADP	O2A-PA-O5'	2.52	118.97	107.57
2	A	301	ADP	O3'-C3'-C4'	-2.48	103.97	111.08
2	C	301	ADP	C6-C5-N7	2.46	136.84	132.09
3	A	302	C0D	CAK-CAL-CAM	-2.44	102.30	113.56
3	B	302	C0D	OAG-CAV-CAW	-2.39	105.16	109.38
2	A	301	ADP	C1'-N9-C8	-2.38	121.82	127.09
2	D	301	ADP	C6-C5-N7	2.37	136.66	132.09
3	D	302	C0D	CAM-NAQ-CAT	-2.36	118.42	122.82
2	B	301	ADP	O3A-PA-O1A	2.27	117.54	110.70
3	B	302	C0D	CAK-CAL-CAM	-2.26	103.12	113.56
2	C	301	ADP	C2-N1-C6	2.25	122.43	118.73
2	A	301	ADP	O3A-PB-O1B	-2.24	99.26	111.04
2	B	301	ADP	C2-N1-C6	2.21	122.37	118.73
3	B	302	C0D	CAL-CAM-NAQ	-2.20	106.04	112.20
2	B	301	ADP	O2'-C2'-C1'	-2.17	102.61	110.10
2	B	301	ADP	C5-C6-N6	-2.11	118.05	123.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	301	ADP	O2A-PA-O3A	-2.11	101.57	107.27
2	D	301	ADP	C2'-C1'-N9	2.09	118.51	113.30
2	D	301	ADP	N9-C8-N7	-2.08	110.99	113.94
2	C	301	ADP	O3B-PB-O2B	2.01	115.34	107.80

There are no chirality outliers.

All (15) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	301	ADP	C2'-C1'-N9-C8
2	D	301	ADP	C2'-C1'-N9-C4
2	A	301	ADP	C2'-C1'-N9-C8
2	B	301	ADP	C2'-C1'-N9-C8
2	B	301	ADP	C2'-C1'-N9-C4
3	A	302	C0D	CAK-CAL-CAM-NAQ
2	A	301	ADP	C2'-C1'-N9-C4
2	C	301	ADP	C2'-C1'-N9-C4
2	C	301	ADP	C2'-C1'-N9-C8
3	C	302	C0D	CAJ-CAK-CAL-CAM
3	B	302	C0D	CAK-CAL-CAM-NAQ
3	A	302	C0D	CAJ-CAK-CAL-CAM
2	A	301	ADP	PB-O3A-PA-O1A
2	C	301	ADP	PB-O3A-PA-O1A
2	C	301	ADP	PB-O3A-PA-O2A

There are no ring outliers.


No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand C0D A 302




Ligand ADP A 301









5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/266 (100%)	0.08	8 (3%) 52 52	12, 22, 44, 66	0
1	B	266/266 (100%)	0.00	6 (2%) 61 61	11, 20, 45, 84	0
1	C	265/266 (99%)	0.46	12 (4%) 38 37	17, 28, 52, 69	0
1	D	266/266 (100%)	0.52	21 (7%) 18 16	15, 28, 70, 84	0
All	All	1063/1064 (99%)	0.27	47 (4%) 39 38	11, 25, 56, 84	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	264	TYR	4.4
1	C	146	HIS	3.7
1	C	184	ASP	3.6
1	C	264	TYR	3.5
1	D	63	ILE	3.5
1	D	185	ALA	3.5
1	A	185	ALA	3.3
1	B	186	ASP	3.2
1	D	184	ASP	3.2
1	C	185	ALA	3.2
1	C	108	GLY	3.0
1	D	163	THR	3.0
1	A	184	ASP	2.9
1	D	47	LEU	2.9
1	D	161	LYS	2.7
1	C	20	ASN	2.7
1	D	40	ASN	2.7
1	C	186	ASP	2.6
1	A	59	GLU	2.6
1	D	265	LEU	2.6
1	C	161	LYS	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	58	ALA	2.5
1	C	265	LEU	2.5
1	A	20	ASN	2.5
1	D	1	MET	2.5
1	A	264	TYR	2.4
1	C	181	HIS	2.4
1	D	46	LYS	2.4
1	B	264	TYR	2.4
1	D	54	ALA	2.4
1	D	56	VAL	2.4
1	D	61	ILE	2.4
1	D	64	PRO	2.4
1	D	62	ASN	2.3
1	A	186	ASP	2.2
1	C	80	ILE	2.2
1	B	266	GLU	2.2
1	D	186	ASP	2.2
1	D	44	ILE	2.2
1	A	183	LEU	2.2
1	B	21	GLN	2.2
1	C	38	TRP	2.1
1	B	161	LYS	2.1
1	B	184	ASP	2.1
1	D	164	GLU	2.1
1	D	60	ASN	2.1
1	A	162	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	C0D	D	302	19/19	0.94	0.10	23,29,39,40	0
3	C0D	B	302	19/19	0.95	0.08	16,18,28,29	0
3	C0D	C	302	19/19	0.96	0.06	20,22,25,27	0
3	C0D	A	302	19/19	0.96	0.06	15,17,20,22	0
2	ADP	C	301	27/27	0.97	0.06	19,22,25,28	0
2	ADP	A	301	27/27	0.98	0.05	15,18,22,28	0
2	ADP	D	301	27/27	0.98	0.05	21,25,28,29	0
2	ADP	B	301	27/27	0.98	0.04	13,15,18,19	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around C0D D 302:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C0D B 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around C0D C 302:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around C0D A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ADP C 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around ADP A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ADP D 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.