



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:10 PM UTC

PDB ID : 1AXA / pdb_00001axa
Title : ACTIVE-SITE MOBILITY IN HUMAN IMMUNODEFICIENCY VIRUS
TYPE 1 PROTEASE AS DEMONSTRATED BY CRYSTAL STRUCTURE
OF A28S MUTANT
Authors : Hong, L.; Hartsuck, J.A.; Foundling, S.; Ermolieff, J.; Tang, J.
Deposited on : 1997-10-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	99	 67% 24% 7% .
1	B	99	 58% 27% 11% .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1671 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HIV-1 PROTEASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	99	Total	C	N	O	S	0	1	0
			760	489	130	137	4			
1	B	99	Total	C	N	O	S	0	2	0
			764	491	130	139	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	SER	ALA	engineered mutation	UNP P03366
B	28	SER	ALA	engineered mutation	UNP P03366

- Molecule 2 is N-[[1-[N-ACETAMIDYL]-[1-CYCLOHEXYLMETHYL-2-HYDROXY-4-ISO
PROPYL]-BUT-4-YL]-CARBONYL]-GLUTAMINYL-ARGINYL-AMIDE (CCD ID: U0E)
(formula: C₂₈H₅₂N₈O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	1
			42	28	8	6		
2	A	1	Total	C	N	O	0	1
			42	28	8	6		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	31	Total	O	0	0
			31	31		
3	B	32	Total	O	0	0
			32	32		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: HIV-1 PROTEASE

Chain A:  67% 24% 7% .



- Molecule 1: HIV-1 PROTEASE

Chain B:  58% 27% 11% .



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	63.55Å 63.55Å 83.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.00	Depositor
% Data completeness (in resolution range)	92.6 (20.00-2.00)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
Refinement program	TNT	Depositor
R, R_{free}	0.194 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1671	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: U0E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	4.12	2/778 (0.3%)	2.00	18/1052 (1.7%)
1	B	4.59	2/787 (0.3%)	2.05	24/1064 (2.3%)
All	All	4.37	4/1565 (0.3%)	2.03	42/2116 (2.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	99	PHE	C-OXT	122.33	3.68	1.23
1	A	99	PHE	C-OXT	107.55	3.38	1.23
1	B	66	ILE	CA-C	-5.58	1.45	1.52
1	A	34	GLU	CD-OE1	5.17	1.35	1.25

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	47	ILE	CA-C-N	-10.99	114.60	122.18
1	A	47	ILE	C-N-CA	-10.99	114.60	122.18
1	B	50	ILE	CA-C-N	-9.76	102.56	122.66
1	B	50	ILE	C-N-CA	-9.76	102.56	122.66
1	A	50	ILE	CA-C-N	-9.69	105.39	122.83
1	A	50	ILE	C-N-CA	-9.69	105.39	122.83
1	A	99	PHE	CA-CB-CG	8.57	122.37	113.80
1	A	50	ILE	N-CA-C	8.02	118.80	110.62
1	B	23	LEU	CB-CA-C	-7.35	97.40	109.53
1	B	84	ILE	CA-CB-CG1	6.88	122.09	110.40
1	A	25	ASP	N-CA-C	6.76	119.86	108.02
1	B	36	MET	N-CA-CB	6.75	122.22	111.56
1	A	69	HIS	CA-CB-CG	6.66	120.46	113.80
1	B	87	ARG	N-CA-C	6.49	119.18	111.33
1	B	50	ILE	O-C-N	-6.44	115.50	121.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	36	MET	N-CA-C	6.38	118.46	108.96
1	B	11	VAL	N-CA-CB	-6.35	102.89	112.35
1	B	98	ASN	N-CA-C	6.21	118.57	108.26
1	A	15	ILE	CA-C-O	5.98	126.95	120.43
1	A	79	PRO	N-CA-CB	5.88	106.67	102.79
1	A	50	ILE	O-C-N	-5.77	116.27	121.87
1	B	84	ILE	CA-CB-CG2	5.76	120.29	110.50
1	A	15	ILE	O-C-N	-5.74	116.87	123.07
1	B	40	GLY	CA-C-N	5.71	130.24	122.19
1	B	40	GLY	C-N-CA	5.71	130.24	122.19
1	A	4	THR	N-CA-CB	5.62	119.58	110.41
1	B	38	LEU	CA-C-O	5.58	123.95	119.59
1	B	5	LEU	N-CA-C	5.57	119.77	112.92
1	B	82	VAL	CA-CB-CG1	5.53	119.80	110.40
1	B	89	LEU	N-CA-C	-5.46	106.75	113.41
1	B	44	PRO	N-CA-CB	5.45	109.10	103.38
1	B	54	ILE	CA-CB-CG2	5.43	119.73	110.50
1	B	34	GLU	CB-CG-CD	5.41	121.80	112.60
1	B	61	GLN	N-CA-CB	5.36	119.64	111.70
1	A	32	VAL	N-CA-C	5.35	115.69	107.77
1	A	32	VAL	CB-CA-C	5.33	117.61	110.42
1	B	64	ILE	CB-CA-C	-5.28	102.38	110.50
1	B	57	ARG	CD-NE-CZ	-5.23	117.08	124.40
1	A	59	TYR	CB-CA-C	-5.11	101.69	110.16
1	A	84	ILE	CB-CA-C	-5.10	103.08	110.63
1	A	72	ILE	N-CA-CB	5.09	118.44	111.41
1	B	38	LEU	CB-CA-C	5.03	118.52	110.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	760	0	807	16	0
1	B	764	0	809	16	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	84	0	101	5	0
3	A	31	0	0	0	0
3	B	32	0	0	0	0
All	All	1671	0	1717	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:GLN:H	1:A:7:GLN:HE21	1.27	0.82
1:A:7:GLN:HE21	1:A:7:GLN:N	1.90	0.69
1:B:20:LYS:HZ1	1:B:36:MET:HG2	1.59	0.67
1:A:28[B]:SER:OG	2:A:101[B]:U0E:O4	2.14	0.65
1:B:65:GLU:HG2	1:B:70:LYS:HG2	1.85	0.58
1:A:97:LEU:HB3	1:B:3:ILE:HB	1.87	0.57
1:A:8:ARG:HH12	2:A:100[A]:U0E:HN6	1.54	0.55
1:B:69:HIS:N	1:B:69:HIS:ND1	2.56	0.54
1:B:32:VAL:HG13	1:B:84:ILE:HD12	1.89	0.53
1:A:8:ARG:NH1	2:A:100[A]:U0E:HN6	2.07	0.52
1:B:36:MET:HE1	1:B:38:LEU:HD22	1.96	0.47
1:A:32:VAL:HG13	1:A:84:ILE:HD12	1.96	0.46
1:B:56:VAL:HG12	1:B:78:GLY:HA3	1.98	0.46
1:A:3:ILE:HB	1:B:97:LEU:HB3	1.98	0.45
1:B:57:ARG:HH11	1:B:57:ARG:HD3	1.53	0.45
1:B:66:ILE:O	1:B:67:CYS:C	2.59	0.44
1:B:97:LEU:HD12	1:B:97:LEU:HA	1.84	0.43
1:A:66:ILE:O	1:A:67:CYS:C	2.61	0.43
1:B:64:ILE:HD12	1:B:64:ILE:HG21	1.73	0.43
1:B:89:LEU:HD12	1:B:89:LEU:HA	1.80	0.43
1:A:36:MET:HE3	1:A:36:MET:HB2	1.65	0.42
1:A:10:LEU:HA	1:A:10:LEU:HD23	1.77	0.41
1:B:58:GLN:HE21	1:B:58:GLN:HB2	1.59	0.41
1:A:8:ARG:NH2	2:A:100[A]:U0E:H262	2.35	0.41
1:A:84:ILE:HD13	1:A:84:ILE:HG21	1.66	0.41
1:A:5:LEU:HB3	1:B:87:ARG:HG2	2.03	0.41
1:A:72:ILE:HD13	1:A:72:ILE:HG21	1.73	0.41
1:B:84:ILE:HG21	1:B:84:ILE:HD13	1.50	0.40
1:A:50:ILE:HD12	2:A:101[B]:U0E:H163	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	98/99 (99%)	96 (98%)	1 (1%)	1 (1%)	12	8
1	B	99/99 (100%)	96 (97%)	3 (3%)	0	100	100
All	All	197/198 (100%)	192 (98%)	4 (2%)	1 (0%)	24	21

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	39	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/84 (101%)	69 (81%)	16 (19%)	1	1
1	B	86/84 (102%)	64 (74%)	22 (26%)	0	0
All	All	171/168 (102%)	133 (78%)	38 (22%)	1	0

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	11	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	13	ILE
1	A	24	LEU
1	A	32	VAL
1	A	35	GLU
1	A	37	SER
1	A	38	LEU
1	A	41	ARG
1	A	47	ILE
1	A	50	ILE
1	A	67	CYS
1	A	69	HIS
1	A	75	VAL
1	A	97	LEU
1	A	98	ASN
1	B	3	ILE
1	B	10	LEU
1	B	11	VAL
1	B	15	ILE
1	B	19	LEU
1	B	20	LYS
1	B	21	GLU
1	B	24	LEU
1	B	32	VAL
1	B	36	MET
1	B	38	LEU
1	B	41	ARG
1	B	43	LYS
1	B	46	MET
1	B	50	ILE
1	B	55	LYS
1	B	61	GLN
1	B	64	ILE
1	B	72	ILE
1	B	75	VAL
1	B	82	VAL
1	B	89	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	58	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	61	GLN
1	A	92	GLN
1	B	58	GLN
1	B	92	GLN
1	B	98	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	U0E	A	101[B]	-	42,42,42	0.87	1 (2%)	49,55,55	1.88	15 (30%)
2	U0E	A	100[A]	-	42,42,42	0.79	1 (2%)	49,55,55	2.07	9 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	U0E	A	101[B]	-	-	7/52/60/60	0/1/1/1
2	U0E	A	100[A]	-	-	9/52/60/60	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	101[B]	U0E	C12-C13	-2.27	1.51	1.54
2	A	100[A]	U0E	C19-C18	2.11	1.58	1.53

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	100[A]	U0E	C20-C19-C18	7.78	127.50	113.16
2	A	100[A]	U0E	C19-C18-C22	6.01	124.81	110.11
2	A	101[B]	U0E	C4-C3-C11	-4.52	105.72	112.55
2	A	101[B]	U0E	C19-C18-C22	4.47	121.04	110.11
2	A	100[A]	U0E	C19-C18-N2	4.26	119.34	110.91
2	A	101[B]	U0E	C22-C18-N2	-3.91	100.54	111.11
2	A	100[A]	U0E	C4-C3-C11	-3.64	107.05	112.55
2	A	100[A]	U0E	C24-C23-N4	-3.22	102.43	110.30
2	A	101[B]	U0E	C11-C3-N1	3.12	115.82	109.97
2	A	101[B]	U0E	C13-C17-N2	-3.02	112.35	116.01
2	A	100[A]	U0E	C18-C22-N4	-2.93	110.39	116.63
2	A	101[B]	U0E	C18-C22-N4	-2.84	110.56	116.63
2	A	101[B]	U0E	C16-C14-C13	2.81	118.00	112.01
2	A	100[A]	U0E	O2-C11-C12	2.80	114.75	109.15
2	A	101[B]	U0E	C4-C3-N1	-2.72	106.70	110.20
2	A	101[B]	U0E	O2-C11-C12	2.59	114.33	109.15
2	A	100[A]	U0E	C16-C14-C13	2.58	117.50	112.01
2	A	100[A]	U0E	C13-C17-N2	-2.51	112.97	116.01
2	A	101[B]	U0E	C25-C23-C24	2.42	116.00	110.14
2	A	101[B]	U0E	C20-C19-C18	2.41	117.61	113.16
2	A	101[B]	U0E	N8-C28-N6	2.26	124.41	119.27
2	A	101[B]	U0E	C24-C23-N4	-2.21	104.90	110.30
2	A	101[B]	U0E	O2-C11-C3	-2.18	105.59	109.88
2	A	101[B]	U0E	C23-C24-N5	-2.05	113.27	116.76

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	100[A]	U0E	C3-C4-C5-C10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	100[A]	U0E	N7-C28-N6-C27
2	A	100[A]	U0E	N8-C28-N6-C27
2	A	101[B]	U0E	C3-C4-C5-C10
2	A	101[B]	U0E	N7-C28-N6-C27
2	A	101[B]	U0E	N8-C28-N6-C27
2	A	100[A]	U0E	N2-C18-C19-C20
2	A	100[A]	U0E	C25-C26-C27-N6
2	A	101[B]	U0E	C23-C25-C26-C27
2	A	101[B]	U0E	C24-C23-C25-C26
2	A	101[B]	U0E	C3-C4-C5-C6
2	A	100[A]	U0E	C3-C4-C5-C6
2	A	101[B]	U0E	N4-C23-C25-C26
2	A	100[A]	U0E	C22-C18-C19-C20
2	A	100[A]	U0E	C14-C13-C17-O3
2	A	100[A]	U0E	C18-C19-C20-C21


There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	101[B]	U0E	2	0
2	A	100[A]	U0E	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand U0E A 101 (B)





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.