



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 06:58 AM UTC

PDB ID : 4AX6 / pdb_00004ax6
Title : HYPOCREA JECORINA CEL6A D221A MUTANT SOAKED WITH 6-CHLORO-4-PHENYLUMBELLIFERYL-BETA-CELLOBIOSIDE
Authors : Wu, M.; Nerinckx, W.; Piens, K.; Ishida, T.; Hansson, H.; Stahlberg, J.; Sandgren, M.
Deposited on : 2012-06-10
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	% 93% 7%
1	B	363	% 95% 5%
2	C	2	50% 50%
2	D	2	50% 50%

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 6308 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EXOGLUCANASE 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	363	Total	C	N	O	S	0	2	0
			2750	1746	463	531	10			
1	B	363	Total	C	N	O	S	0	0	0
			2744	1742	463	529	10			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	221	ALA	ASP	engineered mutation	UNP P07987
B	221	ALA	ASP	engineered mutation	UNP P07987

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			22	12	10			
2	D	2	Total	C	O	0	0	0
			22	12	10			

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		
3	B	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is alpha-D-mannopyranose (CCD ID: MAN) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 11 6 5	0	0
4	A	1	Total C O 11 6 5	0	0
4	A	1	Total C O 11 6 5	0	0
4	A	1	Total C O 11 6 5	0	0
4	A	1	Total C O 11 6 5	0	0
4	A	1	Total C O 11 6 5	0	0
4	A	1	Total C O 11 6 5	0	0
4	B	1	Total C O 11 6 5	0	0
4	B	1	Total C O 11 6 5	0	0
4	B	1	Total C O 11 6 5	0	0
4	B	1	Total C O 11 6 5	0	0
4	B	1	Total C O 11 6 5	0	0

- Molecule 5 is 6-chloranyl-7-oxidanyl-4-phenyl-chromen-2-one (CCD ID: UWU) (formula: $C_{15}H_9ClO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total 19	C 15	Cl 1	O 3	0	0
5	B	1	Total 19	C 15	Cl 1	O 3	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	298	Total 298	O 298	0	0
6	B	246	Total 246	O 246	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: EXOGLUCANASE 2



- Molecule 1: EXOGLUCANASE 2



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



- Molecule 2: beta-D-glucopyranose-(1-4)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	49.54Å 52.96Å 66.54Å 77.33° 78.60° 75.66°	Depositor
Resolution (Å)	64.17 – 2.30 64.17 – 2.30	Depositor EDS
% Data completeness (in resolution range)	98.4 (64.17-2.30) 98.4 (64.17-2.30)	Depositor EDS
R_{merge}	0.20	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.43 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.204 , 0.254 0.204 , 0.256	Depositor DCC
R_{free} test set	1379 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	11.2	Xtriage
Anisotropy	0.661	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 31.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	6308	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BGC, UWU, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/2832	0.78	0/3882
1	B	0.57	0/2820	0.78	0/3866
All	All	0.57	0/5652	0.78	0/7748

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2750	0	2617	15	0
1	B	2744	0	2609	11	0
2	C	22	0	19	0	0
2	D	22	0	19	1	0
3	A	28	0	26	0	0
3	B	28	0	26	0	0
4	A	77	0	70	0	0
4	B	55	0	50	0	0
5	A	19	0	8	1	0
5	B	19	0	8	3	0
6	A	298	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	246	0	0	2	0
All	All	6308	0	5452	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:600:UWU:H8	6:B:2074:HOH:O	1.93	0.66
1:B:410:ARG:HH21	1:B:426:GLN:HG3	1.62	0.61
5:B:600:UWU:C16	5:B:600:UWU:H5	2.30	0.61
1:B:264:ALA:HB2	1:B:299:LEU:HG	1.88	0.56
1:A:112:ALA:HA	1:A:432:GLN:HE22	1.73	0.53
1:B:220:PRO:HA	1:B:263:ASP:CG	2.36	0.51
5:B:600:UWU:H5	5:B:600:UWU:H16	1.94	0.49
1:A:277:ASP:OD1	1:A:340:HIS:HE1	1.96	0.48
1:A:87:THR:HG22	1:A:256:PRO:HB2	1.97	0.47
1:B:304:ALA:HB1	2:D:1:BGC:H2	1.96	0.47
1:A:109:SER:HA	1:A:113:ILE:HD12	1.97	0.47
1:A:262:LEU:HG	1:A:296:LEU:HD11	1.98	0.46
1:B:305:ASN:ND2	1:B:306:TYR:H	2.13	0.46
5:A:600:UWU:H5	5:A:600:UWU:C16	2.44	0.46
1:B:109:SER:HA	1:B:113:ILE:HD12	1.96	0.46
1:A:86:ALA:O	1:A:213:ARG:HD3	2.17	0.44
1:A:361:GLN:HG3	1:A:370:VAL:HG11	2.00	0.44
1:A:410:ARG:HB2	1:A:426:GLN:HE22	1.82	0.44
1:B:180:ALA:HB3	6:B:2076:HOH:O	2.17	0.44
1:B:302:ASN:OD1	1:B:305:ASN:HB3	2.18	0.44
1:A:374:GLY:HA2	1:A:438:LEU:HA	2.01	0.42
1:A:220:PRO:HA	1:A:263:ASP:CG	2.44	0.42
1:B:374:GLY:HA2	1:B:438:LEU:HA	2.02	0.42
1:B:137:ASP:HA	1:B:169:TYR:O	2.20	0.41
1:A:305:ASN:ND2	1:A:306:TYR:H	2.17	0.41
1:A:255:LEU:HB3	1:A:256:PRO:HD2	2.03	0.41
1:A:277:ASP:HB3	1:A:278:PRO:HD3	2.03	0.41
1:A:287:TYR:HD1	1:A:296:LEU:HD23	1.85	0.41
1:B:443:ASN:HA	1:B:444:PRO:C	2.46	0.41
1:A:133:PHE:HB2	1:A:394:VAL:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	363/363 (100%)	351 (97%)	12 (3%)	0	100	100
1	B	361/363 (99%)	347 (96%)	14 (4%)	0	100	100
All	All	724/726 (100%)	698 (96%)	26 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	285/283 (101%)	285 (100%)	0	100	100
1	B	283/283 (100%)	282 (100%)	1 (0%)	84	92
All	All	568/566 (100%)	567 (100%)	1 (0%)	87	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	299	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	158	ASN
1	A	182	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	237	ASN
1	A	285	ASN
1	A	305	ASN
1	A	361	GLN
1	A	362	GLN
1	A	426	GLN
1	A	432	GLN
1	A	441	ASN
1	B	161	ASN
1	B	182	ASN
1	B	237	ASN
1	B	285	ASN
1	B	305	ASN
1	B	319	GLN
1	B	351	GLN
1	B	437	GLN
1	B	441	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BGC	C	1	2,5	11,11,12	1.20	2 (18%)	15,15,17	1.93	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	C	2	2	11,11,12	0.78	0	15,15,17	0.69	0
2	BGC	D	1	2,5	11,11,12	1.43	2 (18%)	15,15,17	2.17	5 (33%)
2	BGC	D	2	2	11,11,12	0.81	0	15,15,17	0.70	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2,5	-	1/2/19/22	0/1/1/1
2	BGC	C	2	2	-	0/2/19/22	0/1/1/1
2	BGC	D	1	2,5	-	2/2/19/22	0/1/1/1
2	BGC	D	2	2	-	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	BGC	O5-C1	2.89	1.48	1.43
2	D	1	BGC	O5-C5	2.63	1.48	1.43
2	C	1	BGC	O5-C5	2.49	1.48	1.43
2	C	1	BGC	O5-C1	2.35	1.47	1.43

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1	BGC	C1-O5-C5	5.84	120.01	112.19
2	C	1	BGC	C1-O5-C5	5.57	119.65	112.19
2	D	1	BGC	O5-C5-C6	3.25	113.99	107.66
2	C	1	BGC	O5-C5-C6	2.48	112.50	107.66
2	D	1	BGC	C6-C5-C4	-2.40	107.12	113.02
2	D	1	BGC	C1-C2-C3	2.09	112.68	109.64
2	D	1	BGC	O5-C5-C4	2.08	115.89	110.83
2	C	1	BGC	O4-C4-C3	-2.00	105.66	110.38

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1	BGC	O5-C5-C6-O6
2	D	1	BGC	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	1	BGC	O5-C5-C6-O6


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	BGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	MAN	B	506	1	11,11,12	0.59	0	15,15,17	0.77	0
4	MAN	B	507	1	11,11,12	0.56	0	15,15,17	0.74	1 (6%)
4	MAN	B	508	1	11,11,12	0.56	0	15,15,17	0.93	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	UWU	B	600	2	21,21,21	1.80	4 (19%)	30,30,30	1.51	6 (20%)
4	MAN	B	504	1	11,11,12	0.58	0	15,15,17	0.85	1 (6%)
4	MAN	A	503	1	11,11,12	0.43	0	15,15,17	1.19	1 (6%)
4	MAN	A	506	1	11,11,12	0.59	0	15,15,17	0.92	1 (6%)
3	NAG	A	501	1	14,14,15	0.51	0	17,19,21	0.69	0
3	NAG	B	501	1	14,14,15	0.53	0	17,19,21	0.86	0
4	MAN	A	509	1	11,11,12	0.58	0	15,15,17	0.74	0
3	NAG	B	502	1	14,14,15	0.50	0	17,19,21	0.67	0
4	MAN	A	505	1	11,11,12	0.48	0	15,15,17	1.24	1 (6%)
3	NAG	A	502	1	14,14,15	0.51	0	17,19,21	0.87	0
4	MAN	A	507	1	11,11,12	0.55	0	15,15,17	0.85	1 (6%)
4	MAN	A	508	1	11,11,12	0.56	0	15,15,17	0.90	1 (6%)
4	MAN	B	509	1	11,11,12	0.61	0	15,15,17	0.80	0
5	UWU	A	600	2	21,21,21	1.80	4 (19%)	30,30,30	1.54	6 (20%)
4	MAN	A	504	1	11,11,12	0.54	0	15,15,17	1.02	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	B	506	1	-	2/2/19/22	0/1/1/1
4	MAN	B	507	1	-	2/2/19/22	0/1/1/1
4	MAN	B	508	1	-	0/2/19/22	0/1/1/1
5	UWU	B	600	2	-	0/4/4/4	0/3/3/3
4	MAN	B	504	1	-	0/2/19/22	0/1/1/1
4	MAN	A	503	1	-	0/2/19/22	0/1/1/1
4	MAN	A	506	1	-	2/2/19/22	0/1/1/1
3	NAG	A	501	1	-	0/6/23/26	0/1/1/1
3	NAG	B	501	1	-	0/6/23/26	0/1/1/1
4	MAN	A	509	1	-	2/2/19/22	0/1/1/1
3	NAG	B	502	1	-	0/6/23/26	0/1/1/1
4	MAN	A	505	1	-	0/2/19/22	0/1/1/1
3	NAG	A	502	1	-	0/6/23/26	0/1/1/1
4	MAN	A	507	1	-	0/2/19/22	0/1/1/1
4	MAN	A	508	1	-	0/2/19/22	0/1/1/1
4	MAN	B	509	1	-	0/2/19/22	0/1/1/1
5	UWU	A	600	2	-	0/4/4/4	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	A	504	1	-	2/2/19/22	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	600	UWU	C11-C4	-5.75	1.40	1.49
5	B	600	UWU	C11-C4	-5.68	1.41	1.49
5	B	600	UWU	C3-C4	3.67	1.39	1.35
5	A	600	UWU	C3-C4	3.44	1.39	1.35
5	B	600	UWU	C4A-C4	-3.20	1.40	1.47
5	A	600	UWU	C4A-C4	-3.16	1.40	1.47
5	A	600	UWU	C3-C2	-2.41	1.39	1.44
5	B	600	UWU	C3-C2	-2.19	1.39	1.44

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	505	MAN	C1-O5-C5	4.44	118.13	112.19
5	A	600	UWU	O8-C2-C3	4.24	122.33	117.16
5	B	600	UWU	O8-C2-C3	3.97	121.99	117.16
4	A	503	MAN	C1-O5-C5	3.88	117.38	112.19
5	A	600	UWU	O2-C2-C3	-3.69	118.50	125.92
5	B	600	UWU	O2-C2-C3	-3.59	118.70	125.92
5	A	600	UWU	C8A-O8-C2	-3.40	117.43	121.67
5	B	600	UWU	C8A-O8-C2	-3.30	117.56	121.67
4	A	504	MAN	C1-O5-C5	2.89	116.06	112.19
5	A	600	UWU	C4-C3-C2	-2.71	120.59	122.89
4	A	508	MAN	C1-O5-C5	2.65	115.74	112.19
4	A	507	MAN	C1-O5-C5	2.56	115.62	112.19
4	B	508	MAN	C1-O5-C5	2.52	115.57	112.19
4	B	504	MAN	C1-O5-C5	2.48	115.51	112.19
5	B	600	UWU	C4-C3-C2	-2.48	120.79	122.89
4	A	506	MAN	C1-O5-C5	2.35	115.34	112.19
5	B	600	UWU	C4A-C4-C11	2.30	124.40	120.41
5	A	600	UWU	C4A-C4-C11	2.29	124.40	120.41
5	B	600	UWU	O8-C2-O2	2.24	119.31	116.40
5	A	600	UWU	O8-C2-O2	2.13	119.17	116.40
4	B	507	MAN	C1-O5-C5	2.05	114.93	112.19

There are no chirality outliers.

All (10) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	507	MAN	C4-C5-C6-O6
4	A	506	MAN	C4-C5-C6-O6
4	B	507	MAN	O5-C5-C6-O6
4	A	506	MAN	O5-C5-C6-O6
4	B	506	MAN	C4-C5-C6-O6
4	A	509	MAN	C4-C5-C6-O6
4	B	506	MAN	O5-C5-C6-O6
4	A	504	MAN	C4-C5-C6-O6
4	A	509	MAN	O5-C5-C6-O6
4	A	504	MAN	O5-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	600	UWU	3	0
5	A	600	UWU	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	363/363 (100%)	0.08	2 (0%) 85 86	8, 11, 17, 21	3 (0%)
1	B	363/363 (100%)	0.24	3 (0%) 82 83	8, 14, 22, 28	3 (0%)
All	All	726/726 (100%)	0.16	5 (0%) 84 85	8, 12, 19, 28	6 (0%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	158	ASN	3.1
1	B	119	ALA	2.6
1	B	319	GLN	2.3
1	A	159	GLY	2.2
1	B	237	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	D	1	11/12	0.76	0.14	15,15,15,15	0
2	BGC	D	2	11/12	0.86	0.10	14,15,15,15	0
2	BGC	C	1	11/12	0.90	0.09	11,11,11,11	0
2	BGC	C	2	11/12	0.91	0.08	10,10,11,11	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around Chain D:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MAN	A	503	11/12	0.58	0.20	19,20,21,21	0
4	MAN	B	507	11/12	0.66	0.16	29,30,30,31	0
4	MAN	A	509	11/12	0.72	0.16	19,20,20,20	0
5	UWU	B	600	19/19	0.77	0.14	15,15,16,16	0
4	MAN	B	504	11/12	0.78	0.12	19,19,19,19	0
3	NAG	B	501	14/15	0.79	0.14	18,19,19,19	0
4	MAN	B	509	11/12	0.80	0.12	28,28,28,28	0
4	MAN	A	508	11/12	0.80	0.12	22,23,23,23	0
3	NAG	A	501	14/15	0.82	0.13	16,16,16,17	0
4	MAN	B	506	11/12	0.83	0.10	25,25,26,26	0
5	UWU	A	600	19/19	0.84	0.11	12,12,12,13	0
4	MAN	A	504	11/12	0.84	0.10	16,16,16,16	0
4	MAN	A	507	11/12	0.86	0.11	18,18,18,18	0
3	NAG	B	502	14/15	0.87	0.10	14,14,15,15	0
4	MAN	A	506	11/12	0.88	0.10	16,16,16,16	0
4	MAN	A	505	11/12	0.88	0.09	16,16,16,16	0
4	MAN	B	508	11/12	0.89	0.08	29,30,30,30	0
3	NAG	A	502	14/15	0.93	0.07	10,10,10,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around UWU B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around UWU A 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.