



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 04:37 PM UTC

PDB ID : 4AXG / pdb_00004axg
Title : Structure of eIF4E-Cup complex
Authors : Kinkelin, K.; Veith, K.; Gruenwald, M.; Bono, F.
Deposited on : 2012-06-12
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	248	
1	B	248	
2	C	130	
2	D	130	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3047 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EUKARYOTIC TRANSLATION INITIATION FACTOR 4E.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	181	Total	C	N	O	S	0	0	0
			1439	912	245	276	6			
1	B	167	Total	C	N	O	S	0	0	0
			1205	769	204	230	2			

- Molecule 2 is a protein called PROTEIN CUP.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	37	Total	C	N	O	S	0	0	0
			288	176	59	50	3			
2	D	15	Total	C	N	O	S	0	0	0
			101	63	18	19	1			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	13	Total	O	0	0
			13	13		
3	B	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: EUKARYOTIC TRANSLATION INITIATION FACTOR 4E



• Molecule 1: EUKARYOTIC TRANSLATION INITIATION FACTOR 4E





• Molecule 2: PROTEIN CUP





● Molecule 2: PROTEIN CUP



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	142.58Å 142.58Å 108.05Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	81.31 – 2.80 81.31 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (81.31-2.80) 100.0 (81.31-2.80)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.82Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.228 , 0.244 (Not available) , 0.272	Depositor DCC
R_{free} test set	825 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	67.1	Xtriage
Anisotropy	0.409	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 71.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3047	wwPDB-VP
Average B, all atoms (Å ²)	77.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.38	0/1472	0.71	0/2002
1	B	0.36	0/1230	0.75	0/1686
2	C	0.27	0/290	0.72	0/386
2	D	0.26	0/101	0.63	0/136
All	All	0.36	0/3093	0.72	0/4210

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1439	0	1347	50	0
1	B	1205	0	1042	25	0
2	C	288	0	280	19	0
2	D	101	0	80	1	0
3	A	13	0	0	0	0
3	B	1	0	0	0	0
All	All	3047	0	2749	75	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

All (75) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:73:MET:CE	2:C:324:VAL:HA	1.36	1.54
1:A:73:MET:HE1	2:C:324:VAL:HA	1.22	1.20
1:A:73:MET:CE	2:C:324:VAL:CA	2.28	1.11
1:A:73:MET:HE3	2:C:325:LYS:H	1.01	1.10
1:A:73:MET:HE2	2:C:324:VAL:HA	1.15	1.07
1:A:73:MET:HE3	2:C:325:LYS:N	1.79	0.96
1:A:222:ARG:HD2	1:A:222:ARG:H	1.35	0.90
1:A:222:ARG:HD2	1:A:222:ARG:N	1.90	0.86
1:A:73:MET:CE	2:C:325:LYS:H	1.87	0.84
1:B:97:THR:CG2	1:B:108:LEU:HD22	2.09	0.83
1:A:73:MET:HE2	2:C:324:VAL:CA	2.01	0.81
1:A:73:MET:HE1	2:C:324:VAL:CA	2.00	0.80
1:A:222:ARG:HH11	1:A:222:ARG:HA	1.47	0.79
1:A:132:ARG:NH1	1:A:134:MET:SD	2.56	0.78
1:A:81:LEU:HB3	1:A:123:ASP:HB2	1.65	0.77
1:A:236:VAL:CG1	1:A:238:GLN:O	2.36	0.73
1:A:118:ILE:HD11	1:A:187:ILE:HD13	1.72	0.72
1:A:99:PHE:CZ	1:A:126:LEU:HD11	2.26	0.70
1:A:234:THR:HG22	1:A:234:THR:O	1.95	0.66
1:B:94:ASN:N	1:B:94:ASN:OD1	2.30	0.64
1:B:97:THR:HG21	1:B:108:LEU:HD22	1.80	0.64
1:B:208:LEU:HD21	1:B:246:TYR:CD2	2.33	0.63
1:A:222:ARG:HG3	1:A:222:ARG:NH1	2.12	0.63
1:A:99:PHE:HZ	1:A:126:LEU:HD11	1.62	0.62
1:A:234:THR:O	1:A:234:THR:CG2	2.48	0.62
1:A:134:MET:O	1:A:140:ASN:ND2	2.30	0.62
1:A:222:ARG:HH11	1:A:222:ARG:CG	2.14	0.61
1:A:73:MET:HE2	2:C:323:PRO:O	2.01	0.60
1:A:222:ARG:HH11	1:A:222:ARG:CA	2.14	0.60
1:A:196:ILE:HG13	1:A:214:LEU:HD11	1.86	0.58
1:A:222:ARG:HH11	1:A:222:ARG:HG3	1.67	0.58
1:B:81:LEU:HB3	1:B:123:ASP:HB2	1.86	0.57
1:A:125:SER:OG	1:A:184:VAL:HG23	2.04	0.57
1:A:76:TRP:HB2	1:A:99:PHE:CZ	2.41	0.56
1:B:80:TYR:O	1:B:94:ASN:OD1	2.24	0.56
1:A:236:VAL:HG11	1:A:238:GLN:O	2.05	0.55
1:A:106:TRP:CE2	2:C:336:ARG:HB2	2.42	0.55
1:A:167:LEU:HD22	2:C:332:LEU:HD13	1.89	0.55
1:B:97:THR:HG22	1:B:108:LEU:HD22	1.86	0.55
1:A:76:TRP:HB2	1:A:99:PHE:CE2	2.42	0.54
1:A:222:ARG:HA	1:A:222:ARG:NH1	2.19	0.54
1:A:73:MET:CE	2:C:325:LYS:N	2.53	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:TRP:HB2	1:A:125:SER:HB2	1.91	0.53
1:B:179:GLN:NE2	1:B:203:ASN:HD22	2.07	0.53
1:A:222:ARG:NH1	1:A:222:ARG:CG	2.73	0.51
1:B:110:ASN:OD1	2:D:336:ARG:NH1	2.42	0.51
1:B:79:TRP:HB2	1:B:125:SER:HB2	1.93	0.51
1:B:79:TRP:CZ3	1:B:127:PHE:HB2	2.46	0.51
1:A:73:MET:HE1	2:C:324:VAL:CB	2.41	0.51
1:B:161:LEU:HD11	1:B:220:LEU:HD21	1.92	0.50
1:B:74:ASN:HD22	1:B:128:LYS:HD3	1.77	0.50
1:A:73:MET:CE	2:C:324:VAL:C	2.85	0.50
1:A:215:ARG:HH21	1:A:222:ARG:NH2	2.10	0.49
1:A:86:SER:HB2	1:A:90:GLU:HB2	1.95	0.49
1:B:179:GLN:HE22	1:B:203:ASN:HD22	1.61	0.48
1:B:127:PHE:CE2	1:B:133:PRO:HA	2.47	0.48
1:A:96:ILE:HG12	2:C:371:MET:HE1	1.96	0.47
1:B:71:PRO:HA	1:B:101:THR:HG22	1.95	0.47
1:A:80:TYR:HB2	1:A:96:ILE:HD11	1.96	0.47
1:B:180:ILE:HG22	1:B:198:THR:HG22	1.96	0.47
1:A:137:ASP:OD1	1:A:138:ALA:N	2.48	0.46
1:A:222:ARG:N	1:A:222:ARG:CD	2.73	0.46
1:B:128:LYS:HE2	1:B:180:ILE:O	2.15	0.46
1:B:128:LYS:HB2	1:B:131:ILE:HD12	1.97	0.45
1:B:126:LEU:HD22	1:B:166:LEU:HD22	1.98	0.44
1:B:187:ILE:O	1:B:188:ARG:HD3	2.17	0.44
1:A:100:ASP:OD1	1:A:101:THR:HG23	2.18	0.43
1:A:81:LEU:HD21	1:A:84:ASP:HA	2.01	0.42
1:A:80:TYR:CD2	2:C:373:ILE:HD11	2.55	0.41
1:B:208:LEU:HD21	1:B:246:TYR:HD2	1.82	0.41
1:B:183:ALA:HB2	1:B:196:ILE:HD13	2.02	0.41
1:B:180:ILE:HA	1:B:198:THR:HA	2.02	0.41
1:A:73:MET:HE1	2:C:324:VAL:HG13	2.03	0.41
1:A:113:LYS:HG3	2:C:373:ILE:O	2.21	0.41
1:B:165:VAL:HG13	1:B:196:ILE:HD11	2.01	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/248 (71%)	173 (98%)	4 (2%)	0	100	100
1	B	161/248 (65%)	160 (99%)	1 (1%)	0	100	100
2	C	33/130 (25%)	33 (100%)	0	0	100	100
2	D	13/130 (10%)	13 (100%)	0	0	100	100
All	All	384/756 (51%)	379 (99%)	5 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	152/215 (71%)	148 (97%)	4 (3%)	40	75
1	B	108/215 (50%)	104 (96%)	4 (4%)	30	65
2	C	29/118 (25%)	28 (97%)	1 (3%)	32	68
2	D	7/118 (6%)	7 (100%)	0	100	100
All	All	296/666 (44%)	287 (97%)	9 (3%)	36	72

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	95	GLU
1	A	97	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	222	ARG
1	A	225	SER
1	B	77	THR
1	B	94	ASN
1	B	161	LEU
1	B	177	SER
2	C	375	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	231	HIS
1	B	142	GLN
1	B	203	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	181/248 (72%)	0.42	14 (7%) 19 14	31, 55, 106, 155	0
1	B	167/248 (67%)	1.32	48 (28%) 1 1	44, 95, 140, 154	0
2	C	37/130 (28%)	1.26	9 (24%) 2 1	36, 82, 116, 127	0
2	D	15/130 (11%)	1.21	4 (26%) 1 1	86, 111, 133, 133	0
All	All	400/756 (52%)	0.90	75 (18%) 3 2	31, 75, 134, 155	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	238	GLN	5.7
1	B	206	ALA	5.5
2	C	372	ASN	5.1
1	A	241	ASN	5.1
1	B	83	ASN	4.4
1	B	208	LEU	4.1
1	B	134	MET	4.0
1	B	201	GLY	3.8
1	B	92	MET	3.8
1	B	178	ASP	3.7
1	B	177	SER	3.7
1	A	66	HIS	3.6
1	A	223	ASN	3.6
1	B	241	ASN	3.6
1	B	220	LEU	3.5
2	C	374	TRP	3.5
1	B	180	ILE	3.5
1	B	120	LEU	3.5
1	A	132	ARG	3.4
1	B	203	ASN	3.4
1	B	72	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	200	ASP	3.3
1	B	205	GLU	3.2
2	D	339	MET	3.2
1	B	130	ASN	3.2
2	C	373	ILE	3.0
1	B	216	ASP	3.0
1	B	235	MET	3.0
1	B	210	ILE	3.0
1	B	174	PHE	3.0
2	D	327	TYR	2.9
1	B	151	ASN	2.9
1	A	134	MET	2.9
1	B	73	MET	2.9
1	B	246	TYR	2.8
2	C	364	LEU	2.8
1	B	199	ALA	2.7
1	B	191	SER	2.7
1	B	99	PHE	2.7
2	D	325	LYS	2.6
2	C	319	VAL	2.6
1	A	236	VAL	2.5
1	B	219	ARG	2.5
2	C	362	ILE	2.5
1	A	227	GLN	2.5
1	B	234	THR	2.5
1	B	82	GLU	2.5
1	B	138	ALA	2.5
1	B	233	ASP	2.4
2	C	363	GLU	2.4
1	B	223	ASN	2.4
1	B	222	ARG	2.4
2	D	326	SER	2.4
1	B	142	GLN	2.4
2	C	376	THR	2.3
1	B	243	LYS	2.3
2	C	321	SER	2.3
1	B	168	CYS	2.3
1	B	247	THR	2.3
1	A	191	SER	2.2
1	B	248	LEU	2.2
1	A	222	ARG	2.2
1	B	131	ILE	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	189	GLY	2.1
1	B	176	HIS	2.1
1	B	212	HIS	2.1
1	B	202	ASN	2.1
1	B	175	ASP	2.1
1	B	135	TRP	2.1
1	A	235	MET	2.0
1	B	221	GLY	2.0
1	A	91	ASP	2.0
1	B	133	PRO	2.0
1	B	110	ASN	2.0
1	A	100	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.