



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:09 AM UTC

PDB ID : 4AYX / pdb_00004ayx
Title : STRUCTURE OF THE HUMAN MITOCHONDRIAL ABC TRANSPORTER, ABCB10 (ROD FORM B)
Authors : Pike, A.C.W.; Shintre, C.A.; Li, Q.; Kim, J.; von Delft, F.; Barr, A.J.; Das, S.; Chaikuad, A.; Xia, X.; Quigley, A.; Dong, Y.; Dong, L.; Krojer, T.; Vollmar, M.; Muniz, J.R.C.; Bray, J.E.; Berridge, G.; Chalk, R.; Gileadi, O.; Burgess-Brown, N.; Shrestha, L.; Goubin, S.; Yang, J.; Mahajan, P.; Mukhopadhyay, S.; Bullock, A.N.; Arrowsmith, C.H.; Weigelt, J.; Bountra, C.; Edwards, A.M.; Carpenter, E.P.
Deposited on : 2012-06-22
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	: 4-5-2 with Phenix2.0
Mogul	: 2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	: 2.0
EDS	: 3.0
Buster-report	: wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	: 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	: 9.0.010 (Gargrove)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	595	12% 88% 8% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

Density-Fitness	: 1.0.12
Ideal geometry (proteins)	: Engh & Huber (2001)
Ideal geometry (DNA, RNA)	: Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	: 2.49

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	CDL	A	1721	X	-	-	-

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4428 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ATP-BINDING CASSETTE SUB-FAMILY B MEMBER 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	571	4200	2687	726	771	16	0	1	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	151	MET	-	expression tag	UNP Q9NRK6
A	739	ALA	-	expression tag	UNP Q9NRK6
A	740	GLU	-	expression tag	UNP Q9NRK6
A	741	ASN	-	expression tag	UNP Q9NRK6
A	742	LEU	-	expression tag	UNP Q9NRK6
A	743	TYR	-	expression tag	UNP Q9NRK6
A	744	PHE	-	expression tag	UNP Q9NRK6
A	745	GLN	-	expression tag	UNP Q9NRK6

- Molecule 2 is GLYCINE (CCD ID: GLY) (formula: C₂H₅NO₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			5	2	1	2		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		

- Molecule 4 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (CCD ID: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 5 is DODECYL-BETA-D-MALTOSIDE (CCD ID: LMT) (formula: $C_{24}H_{46}O_{11}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			35	24	11		
5	A	1	Total	C	O	0	0
			35	24	11		

- Molecule 6 is CARDIOLIPIN (CCD ID: CDL) (formula: C₈₁H₁₅₆O₁₇P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	O	P	0	0
			74	55	17	2		
6	A	1	Total	C	O		0	0
			14	12	2			


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	33	Total 33	O 33	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP-BINDING CASSETTE SUB-FAMILY B MEMBER 10



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	180.14Å 180.14Å 50.56Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	39.00 – 2.90 39.00 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (39.00-2.90) 99.3 (39.00-2.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.91 (at 2.90Å)	Xtriage
Refinement program	BUSTER 2.11.2	Depositor
R, R_{free}	0.223 , 0.267 0.232 , 0.261	Depositor DCC
R_{free} test set	1088 reflections (5.17%)	wwPDB-VP
Wilson B-factor (Å ²)	79.9	Xtriage
Anisotropy	0.501	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 59.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.046 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4428	wwPDB-VP
Average B, all atoms (Å ²)	90.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CDL, LMT, ACP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	1/4276 (0.0%)	1.39	2/5811 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	287	THR	CA-C	5.72	1.55	1.52

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	264	ASP	CA-CB-CG	5.17	117.77	112.60
1	A	557	ASP	CA-CB-CG	5.09	117.69	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4200	0	4192	19	0
2	A	5	0	2	0	0
3	A	1	0	0	0	0
4	A	31	0	14	0	0
5	A	70	0	92	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	88	0	116	1	0
7	A	33	0	0	0	0
All	All	4428	0	4416	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:LEU:HD23	1:A:436:MET:HE1	1.72	0.70
1:A:537:LEU:HD22	1:A:656:LEU:HB3	1.76	0.68
1:A:171:LEU:HD21	1:A:287:THR:HG22	1.84	0.59
1:A:182:SER:O	1:A:186:MET:HG2	2.05	0.57
1:A:178:LEU:HB2	1:A:232:ARG:HD2	1.88	0.55
1:A:258:GLN:HA	1:A:473:PRO:HB3	1.89	0.55
1:A:498:HIS:HB2	1:A:548:SER:HB3	1.90	0.54
1:A:571:GLY:HA3	1:A:652:PRO:HG3	1.92	0.50
1:A:420:LEU:HB2	1:A:426:MET:HE2	1.95	0.47
1:A:702:ALA:HB1	1:A:709:ILE:HD12	1.96	0.47
1:A:497:VAL:HG13	1:A:546:PRO:HB3	1.96	0.47
1:A:522:VAL:HG22	1:A:685:VAL:HB	1.99	0.45
1:A:533:LYS:HB3	1:A:688:ILE:HG12	1.98	0.44
1:A:409:ILE:HG13	5:A:1724:LMT:H122	2.00	0.43
1:A:627:VAL:HG11	1:A:634:LEU:HD11	2.00	0.42
5:A:1724:LMT:H5B	5:A:1724:LMT:H6D	2.02	0.41
1:A:184:ILE:HB	6:A:1726:CDL:H381	2.03	0.41
1:A:285:SER:HA	1:A:289:ASN:HB3	2.03	0.40
1:A:290:LEU:O	1:A:294:LEU:HB2	2.21	0.40
1:A:294:LEU:HD12	1:A:294:LEU:HA	1.98	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	570/595 (96%)	561 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	427/489 (87%)	415 (97%)	12 (3%)	38	72

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	175	VAL
1	A	281	LEU
1	A	290	LEU
1	A	291	SER
1	A	365	ARG
1	A	440	TRP
1	A	475	LEU
1	A	493	GLU
1	A	507	VAL
1	A	526	VAL
1	A	630	LYS
1	A	648	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	299	GLN
1	A	386	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 7 ligands modelled in this entry, 1 is monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	LMT	A	1720	-	36,36,36	0.19	0	47,47,47	0.41	0
4	ACP	A	900	3	31,33,33	0.42	0	47,52,52	0.93	4 (8%)
6	CDL	A	1726	-	13,13,99	0.91	1 (7%)	13,13,111	0.71	0
2	GLY	A	1717	-	4,4,4	1.02	0	3,4,4	0.89	0
6	CDL	A	1721	-	73,73,99	0.73	2 (2%)	79,85,111	0.63	4 (5%)
5	LMT	A	1724	-	36,36,36	0.22	0	47,47,47	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	LMT	A	1720	-	-	4/21/61/61	0/2/2/2
4	ACP	A	900	3	-	3/19/38/38	0/3/3/3
6	CDL	A	1726	-	-	2/12/12/110	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLY	A	1717	-	-	2/2/2/2	-
6	CDL	A	1721	-	1/1/9/9	30/84/84/110	-
5	LMT	A	1724	-	-	4/21/61/61	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1721	CDL	C79-C78	-3.88	1.32	1.51
6	A	1721	CDL	C39-C38	-3.85	1.32	1.51
6	A	1726	CDL	C39-C38	-3.10	1.32	1.51

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	900	ACP	O2A-PA-O5'	-3.14	93.32	107.57
4	A	900	ACP	O1B-PB-C3B	3.14	117.31	109.05
6	A	1721	CDL	C39-C38-C37	2.47	126.86	114.37
4	A	900	ACP	O3A-PA-O1A	2.47	118.12	110.70
6	A	1721	CDL	C79-C78-C77	2.35	126.27	114.37
6	A	1721	CDL	C80-C79-C78	2.33	126.13	114.37
4	A	900	ACP	O2B-PB-O1B	2.32	117.50	109.95
6	A	1721	CDL	C40-C39-C38	2.20	125.50	114.37

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
6	A	1721	CDL	CA4

All (45) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1717	GLY	OXT-C-CA-N
4	A	900	ACP	C5'-O5'-PA-O2A
6	A	1721	CDL	CA2-C1-CB2-OB2
6	A	1721	CDL	CA3-OA5-PA1-OA2
6	A	1721	CDL	CA3-OA5-PA1-OA3
6	A	1721	CDL	CA3-OA5-PA1-OA4
6	A	1721	CDL	CB2-OB2-PB2-OB3
6	A	1721	CDL	CB2-OB2-PB2-OB4
6	A	1721	CDL	CB2-OB2-PB2-OB5
6	A	1721	CDL	CA4-CA6-OA8-CA7

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	A	1721	CDL	O1-C1-CB2-OB2
2	A	1717	GLY	O-C-CA-N
5	A	1720	LMT	O1'-C1-C2-C3
5	A	1720	LMT	C1-C2-C3-C4
5	A	1724	LMT	C7-C8-C9-C10
6	A	1726	CDL	C31-C32-C33-C34
5	A	1724	LMT	C11-C10-C9-C8
6	A	1721	CDL	OA5-CA3-CA4-CA6
6	A	1721	CDL	C39-C40-C41-C42
6	A	1721	CDL	OB5-CB3-CB4-CB6
6	A	1721	CDL	CA7-C31-C32-C33
6	A	1721	CDL	C79-C80-C81-C82
5	A	1720	LMT	C3-C4-C5-C6
6	A	1721	CDL	OA5-CA3-CA4-OA6
6	A	1721	CDL	OB5-CB3-CB4-OB6
4	A	900	ACP	C5'-O5'-PA-O1A
4	A	900	ACP	C5'-O5'-PA-O3A
6	A	1721	CDL	CB3-OB5-PB2-OB2
5	A	1724	LMT	O1'-C1-C2-C3
6	A	1721	CDL	CB6-CB4-OB6-CB5
6	A	1721	CDL	OB6-CB4-CB6-OB8
6	A	1721	CDL	CA5-C11-C12-C13
6	A	1721	CDL	CA3-CA4-OA6-CA5
5	A	1724	LMT	C4-C5-C6-C7
6	A	1721	CDL	CB4-CB3-OB5-PB2
6	A	1721	CDL	C42-C43-C44-C45
6	A	1721	CDL	C11-CA5-OA6-CA4
5	A	1720	LMT	C2-C3-C4-C5
6	A	1721	CDL	C12-C11-CA5-OA6
6	A	1721	CDL	C80-C81-C82-C83
6	A	1721	CDL	CA6-CA4-OA6-CA5
6	A	1721	CDL	CB3-CB4-CB6-OB8
6	A	1721	CDL	C12-C11-CA5-OA7
6	A	1721	CDL	C82-C83-C84-C85
6	A	1726	CDL	C32-C31-CA7-OA8


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1726	CDL	1	0
5	A	1724	LMT	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	571/595 (95%)	0.79	70 (12%) 8 7	51, 89, 123, 189	1 (0%)

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	689	ALA	8.5
1	A	693	SER	6.9
1	A	526	VAL	6.8
1	A	309	PHE	6.4
1	A	703	VAL	5.5
1	A	724	GLY	5.2
1	A	524	ALA	5.0
1	A	690	HIS	4.9
1	A	692	LEU	4.7
1	A	450	SER	4.3
1	A	525	LEU	4.3
1	A	200[A]	TYR	4.0
1	A	695	ILE	4.0
1	A	715	HIS	4.0
1	A	289	ASN	4.0
1	A	290	LEU	4.0
1	A	453	SER	4.0
1	A	669	GLU	3.9
1	A	687	VAL	3.8
1	A	454	GLU	3.7
1	A	662	SER	3.7
1	A	292	ASP	3.7
1	A	720	SER	3.6
1	A	163	LEU	3.6
1	A	379	SER	3.5
1	A	468	LEU	3.5
1	A	503	ALA	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	285	SER	3.3
1	A	666	ALA	3.3
1	A	310	VAL	3.2
1	A	664	LEU	3.2
1	A	212	THR	3.2
1	A	710	THR	3.1
1	A	691	ARG	3.0
1	A	181	SER	2.9
1	A	668	ASN	2.9
1	A	702	ALA	2.8
1	A	723	ASN	2.8
1	A	470	GLU	2.8
1	A	598	VAL	2.7
1	A	722	PRO	2.7
1	A	670	TYR	2.7
1	A	253	SER	2.7
1	A	154	LEU	2.7
1	A	335	TYR	2.6
1	A	694	THR	2.6
1	A	660	ALA	2.5
1	A	665	ASP	2.5
1	A	704	LEU	2.5
1	A	632	VAL	2.5
1	A	451	PHE	2.4
1	A	487	SER	2.4
1	A	257	ARG	2.3
1	A	580	PHE	2.3
1	A	625	THR	2.3
1	A	600	ALA	2.3
1	A	667	GLU	2.3
1	A	399	PHE	2.3
1	A	455	LEU	2.3
1	A	185	SER	2.2
1	A	394	ALA	2.2
1	A	397	GLY	2.2
1	A	488	PHE	2.2
1	A	459	LEU	2.2
1	A	209	ASP	2.1
1	A	646	ARG	2.1
1	A	190	PHE	2.1
1	A	716	GLU	2.1
1	A	337	ARG	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	697	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	MG	A	801	1/1	0.41	0.22	130,130,130,130	0
2	GLY	A	1717	5/5	0.69	0.18	96,101,104,119	0
5	LMT	A	1720	35/35	0.81	0.18	98,116,124,124	0
6	CDL	A	1726	14/100	0.81	0.23	114,121,128,129	0
6	CDL	A	1721	74/100	0.86	0.21	89,101,108,110	74
4	ACP	A	900	31/31	0.86	0.12	132,134,150,151	0
5	LMT	A	1724	35/35	0.88	0.15	99,107,122,122	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around LMT A 1720:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around CDL A 1726:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CDL A 1721:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ACP A 900:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around LMT A 1724:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.