



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 07:57 PM UTC

PDB ID : 5AYD / pdb\_00005ayd  
Title : Crystal structure of Ruminococcus albus beta-(1,4)-mannooligosaccharide phosphorylase (RaMP2) in complexes with phosphate  
Authors : Ye, Y.; Saburi, W.; Kato, K.; Yao, M.  
Deposited on : 2015-08-13  
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	85% 13% .
1	B	335	87% 12% .
1	C	335	84% 14% .
1	D	335	85% 14% .
1	E	335	88% 11% ..

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	335	 89% 11%

## 2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 17442 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-1,4-mannooligosaccharide phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	335	2700	1737	446	497	20	0	0	0
1	B	335	2700	1737	446	497	20	0	0	0
1	C	335	2700	1737	446	497	20	0	0	0
1	D	335	2700	1737	446	497	20	0	0	0
1	E	335	2700	1737	446	497	20	0	0	0
1	F	335	2700	1737	446	497	20	0	0	0

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	192	Total	O	0	0
			192	192		
3	B	212	Total	O	0	0
			212	212		
3	C	221	Total	O	0	0
			221	221		
3	D	185	Total	O	0	0
			185	185		
3	E	172	Total	O	0	0
			172	172		
3	F	230	Total	O	0	0
			230	230		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-1,4-mannooligosaccharide phosphorylase




- Molecule 1: Beta-1,4-mannooligosaccharide phosphorylase



- Molecule 1: Beta-1,4-mannooligosaccharide phosphorylase




- Molecule 1: Beta-1,4-mannooligosaccharide phosphorylase





- Molecule 1: Beta-1,4-mannooligosaccharide phosphorylase

Chain E: 88% 11% ..



- Molecule 1: Beta-1,4-mannooligosaccharide phosphorylase

Chain F: 89% 11%



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.68Å 166.18Å 92.97Å 90.00° 118.91° 90.00°	Depositor
Resolution (Å)	44.76 – 2.30 44.76 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.76-2.30) 98.8 (44.76-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.13 (at 2.29Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692), Coot	Depositor
R, $R_{free}$	0.168 , 0.212 0.171 , 0.213	Depositor DCC
$R_{free}$ test set	5437 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtrriage
Anisotropy	0.435	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 25.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.034 for -h-l,k,h 0.034 for l,k,-h-l 0.045 for h,-k,-h-l 0.038 for -h-l,-k,l 0.196 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	17442	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/2787	0.76	2/3791 (0.1%)
1	B	0.51	0/2787	0.80	1/3791 (0.0%)
1	C	0.50	0/2787	0.82	0/3791
1	D	0.48	0/2787	0.81	2/3791 (0.1%)
1	E	0.46	0/2787	0.80	3/3791 (0.1%)
1	F	0.49	0/2787	0.80	1/3791 (0.0%)
All	All	0.49	0/16722	0.80	9/22746 (0.0%)

There are no bond length outliers.

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	261	GLU	CA-C-N	7.14	126.85	119.56
1	E	261	GLU	C-N-CA	7.14	126.85	119.56
1	B	60	GLY	N-CA-C	6.12	118.40	110.45
1	D	50	SER	N-CA-C	6.07	119.43	109.72
1	A	107	VAL	N-CA-C	5.45	115.65	110.42

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2700	0	2564	32	0
1	B	2700	0	2564	28	0
1	C	2700	0	2564	43	0
1	D	2700	0	2564	29	0
1	E	2700	0	2564	29	0
1	F	2700	0	2564	23	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	192	0	0	7	0
3	B	212	0	0	7	0
3	C	221	0	0	14	0
3	D	185	0	0	6	0
3	E	172	0	0	6	0
3	F	230	0	0	5	0
All	All	17442	0	15384	179	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 179 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:327:ASP:OD1	3:E:501:HOH:O	1.85	0.94
1:F:5:ILE:HD11	1:F:322:VAL:HG21	1.50	0.91
1:B:5:ILE:HD11	1:B:322:VAL:HG21	1.53	0.91
1:C:230:THR:HG22	1:C:232:GLU:H	1.41	0.84
1:A:5:ILE:HD11	1:A:322:VAL:HG21	1.62	0.81

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	333/335 (99%)	315 (95%)	16 (5%)	2 (1%)	21	27
1	B	333/335 (99%)	316 (95%)	15 (4%)	2 (1%)	21	27
1	C	333/335 (99%)	318 (96%)	14 (4%)	1 (0%)	36	46
1	D	333/335 (99%)	316 (95%)	15 (4%)	2 (1%)	21	27
1	E	333/335 (99%)	315 (95%)	16 (5%)	2 (1%)	21	27
1	F	333/335 (99%)	317 (95%)	14 (4%)	2 (1%)	21	27
All	All	1998/2010 (99%)	1897 (95%)	90 (4%)	11 (1%)	21	27

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	GLU
1	B	122	GLU
1	C	122	GLU
1	D	122	GLU
1	D	334	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	294/294 (100%)	289 (98%)	5 (2%)	53	72
1	B	294/294 (100%)	289 (98%)	5 (2%)	53	72
1	C	294/294 (100%)	288 (98%)	6 (2%)	48	67
1	D	294/294 (100%)	288 (98%)	6 (2%)	48	67
1	E	294/294 (100%)	286 (97%)	8 (3%)	39	58
1	F	294/294 (100%)	292 (99%)	2 (1%)	76	87
All	All	1764/1764 (100%)	1732 (98%)	32 (2%)	51	70

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	265	VAL
1	E	334	SER
1	C	122	GLU
1	C	102	LYS
1	F	1	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	277	GLN
1	F	99	GLN
1	F	288	ASN
1	F	152	GLN
1	F	196	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PO4	B	401	-	4,4,4	0.95	0	6,6,6	0.64	0
2	PO4	C	401	-	4,4,4	0.85	0	6,6,6	0.47	0
2	PO4	F	401	-	4,4,4	1.04	0	6,6,6	0.57	0
2	PO4	E	401	-	4,4,4	1.04	0	6,6,6	0.77	0
2	PO4	A	401	-	4,4,4	0.81	0	6,6,6	0.75	0
2	PO4	D	401	-	4,4,4	0.78	0	6,6,6	0.99	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	335/335 (100%)	-1.61	0 100 100	14, 26, 40, 53	0
1	B	335/335 (100%)	-1.64	0 100 100	13, 23, 40, 58	0
1	C	335/335 (100%)	-1.65	0 100 100	14, 23, 39, 58	0
1	D	335/335 (100%)	-1.58	1 (0%) 90 90	15, 26, 42, 60	0
1	E	335/335 (100%)	-1.60	0 100 100	16, 27, 44, 55	0
1	F	335/335 (100%)	-1.66	0 100 100	14, 23, 39, 59	0
All	All	2010/2010 (100%)	-1.62	1 (0%) 100 100	13, 24, 42, 60	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	335	VAL	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	PO4	A	401	5/5	1.00	0.02	18,19,22,22	0
2	PO4	B	401	5/5	1.00	0.02	15,17,19,20	0
2	PO4	C	401	5/5	1.00	0.02	15,15,19,20	0
2	PO4	D	401	5/5	1.00	0.01	14,19,22,24	0
2	PO4	E	401	5/5	1.00	0.01	17,19,20,20	0
2	PO4	F	401	5/5	1.00	0.01	15,16,21,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.