



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:44 PM UTC

PDB ID : 6AYT / pdb_00006ayt
Title : Crystal structure of Campylobacter jejuni 5'-methylthioadenosine/S-adenosyl homocysteine nucleosidase (MTAN) complexed with pyrazinylthio-DADMe-I mmucillin-A
Authors : Harijan, R.K.; Ducati, R.G.; Bonanno, J.B.; Almo, S.C.; Schramm, V.L.
Deposited on : 2017-09-08
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	10% 94% . .
1	B	238	16% 92% . .
1	C	238	22% 93% . .
1	D	238	16% 94% . .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7683 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5'-methylthioadenosine/S-adenosylhomocysteine nucleosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	229	Total	C	N	O	S	0	0	0
			1764	1138	278	337	11			
1	B	229	Total	C	N	O	S	0	1	0
			1767	1140	278	338	11			
1	C	229	Total	C	N	O	S	0	1	0
			1767	1140	278	338	11			
1	D	229	Total	C	N	O	S	0	0	0
			1764	1138	278	337	11			

There are 44 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-9	MET	-	initiating methionine	UNP A0A1E7P7U4
A	-8	GLY	-	expression tag	UNP A0A1E7P7U4
A	-7	HIS	-	expression tag	UNP A0A1E7P7U4
A	-6	HIS	-	expression tag	UNP A0A1E7P7U4
A	-5	HIS	-	expression tag	UNP A0A1E7P7U4
A	-4	HIS	-	expression tag	UNP A0A1E7P7U4
A	-3	HIS	-	expression tag	UNP A0A1E7P7U4
A	-2	HIS	-	expression tag	UNP A0A1E7P7U4
A	-1	SER	-	expression tag	UNP A0A1E7P7U4
A	0	GLY	-	expression tag	UNP A0A1E7P7U4
A	40	ASP	ASN	conflict	UNP A0A1E7P7U4
B	-9	MET	-	initiating methionine	UNP A0A1E7P7U4
B	-8	GLY	-	expression tag	UNP A0A1E7P7U4
B	-7	HIS	-	expression tag	UNP A0A1E7P7U4
B	-6	HIS	-	expression tag	UNP A0A1E7P7U4
B	-5	HIS	-	expression tag	UNP A0A1E7P7U4
B	-4	HIS	-	expression tag	UNP A0A1E7P7U4
B	-3	HIS	-	expression tag	UNP A0A1E7P7U4
B	-2	HIS	-	expression tag	UNP A0A1E7P7U4
B	-1	SER	-	expression tag	UNP A0A1E7P7U4
B	0	GLY	-	expression tag	UNP A0A1E7P7U4

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	40	ASP	ASN	conflict	UNP A0A1E7P7U4
C	-9	MET	-	initiating methionine	UNP A0A1E7P7U4
C	-8	GLY	-	expression tag	UNP A0A1E7P7U4
C	-7	HIS	-	expression tag	UNP A0A1E7P7U4
C	-6	HIS	-	expression tag	UNP A0A1E7P7U4
C	-5	HIS	-	expression tag	UNP A0A1E7P7U4
C	-4	HIS	-	expression tag	UNP A0A1E7P7U4
C	-3	HIS	-	expression tag	UNP A0A1E7P7U4
C	-2	HIS	-	expression tag	UNP A0A1E7P7U4
C	-1	SER	-	expression tag	UNP A0A1E7P7U4
C	0	GLY	-	expression tag	UNP A0A1E7P7U4
C	40	ASP	ASN	conflict	UNP A0A1E7P7U4
D	-9	MET	-	initiating methionine	UNP A0A1E7P7U4
D	-8	GLY	-	expression tag	UNP A0A1E7P7U4
D	-7	HIS	-	expression tag	UNP A0A1E7P7U4
D	-6	HIS	-	expression tag	UNP A0A1E7P7U4
D	-5	HIS	-	expression tag	UNP A0A1E7P7U4
D	-4	HIS	-	expression tag	UNP A0A1E7P7U4
D	-3	HIS	-	expression tag	UNP A0A1E7P7U4
D	-2	HIS	-	expression tag	UNP A0A1E7P7U4
D	-1	SER	-	expression tag	UNP A0A1E7P7U4
D	0	GLY	-	expression tag	UNP A0A1E7P7U4
D	40	ASP	ASN	conflict	UNP A0A1E7P7U4

- Molecule 2 is (3R,4S)-1-[(4-amino-5H-pyrrolo[3,2-d]pyrimidin-7-yl)methyl]-4-[(pyrazin-2-ylsulfanylmethyl)pyrrolidin-3-ol (CCD ID: 4EH) (formula: C₁₆H₁₉N₇OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			25	16	7	1	1		
2	B	1	Total	C	N	O	S	0	0
			25	16	7	1	1		
2	C	1	Total	C	N	O	S	0	0
			25	16	7	1	1		
2	D	1	Total	C	N	O	S	0	0
			25	16	7	1	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	140	Total	O	0	0
			140	140		
4	B	120	Total	O	0	0
			120	120		
4	C	117	Total	O	0	0
			117	117		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	132	Total 132	O 132	0	0



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-methylthioadenosine/S-adenosylhomocysteine nucleosidase



- Molecule 1: 5'-methylthioadenosine/S-adenosylhomocysteine nucleosidase



- Molecule 1: 5'-methylthioadenosine/S-adenosylhomocysteine nucleosidase



- Molecule 1: 5'-methylthioadenosine/S-adenosylhomocysteine nucleosidase



Y221
L222
K223
N224
C225
E226
K227
L228

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	71.08Å 90.98Å 72.45Å 90.00° 111.44° 90.00°	Depositor
Resolution (Å)	67.44 – 1.85 67.44 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.7 (67.44-1.85) 99.7 (67.44-1.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.219 , 0.245 0.231 , 0.251	Depositor DCC
R_{free} test set	3703 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	19.4	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 41.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for l,-k,h	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7683	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 67.06 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.4684e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, 4EH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/1792	0.76	0/2414
1	B	0.54	0/1798	0.77	1/2422 (0.0%)
1	C	0.53	0/1798	0.74	0/2422
1	D	0.52	0/1792	0.76	0/2414
All	All	0.53	0/7180	0.76	1/9672 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	124	LYS	CA-CB-CG	5.99	126.08	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1764	0	1789	5	0
1	B	1767	0	1794	5	0
1	C	1767	0	1794	3	0
1	D	1764	0	1789	2	0
2	A	25	0	19	2	0
2	B	25	0	19	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	25	0	19	1	0
2	D	25	0	19	0	0
3	A	4	0	6	0	0
3	C	4	0	6	0	0
3	D	4	0	6	0	0
4	A	140	0	0	0	0
4	B	120	0	0	1	0
4	C	117	0	0	0	0
4	D	132	0	0	0	0
All	All	7683	0	7260	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:MET:HG2	1:A:50:ILE:HD11	1.81	0.62
1:A:50:ILE:HD12	2:A:301:4EH:O3'	1.99	0.61
1:B:50:ILE:HD11	1:C:112:VAL:HG12	1.90	0.54
1:B:26:ILE:HG22	1:B:35:PHE:HE1	1.75	0.52
1:B:122:ASP:OD2	1:B:124:LYS:HG3	2.09	0.52
1:B:116:GLU:OE1	4:B:401:HOH:O	2.20	0.46
1:D:92:ALA:HB2	1:D:145:ILE:HD12	1.97	0.46
1:C:92:ALA:HB2	1:C:145:ILE:HD12	1.98	0.46
1:A:9:MET:CG	1:A:50:ILE:HD11	2.45	0.46
1:B:92:ALA:HB2	1:B:145:ILE:HD12	1.98	0.45
1:A:92:ALA:HB2	1:A:145:ILE:HD12	2.00	0.43
1:D:2:LYS:HD3	1:D:66:PHE:O	2.18	0.43
1:A:50:ILE:HG23	2:A:301:4EH:H2	2.01	0.42
1:C:50:ILE:HG23	2:C:301:4EH:H2	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	227/238 (95%)	219 (96%)	8 (4%)	0	100	100
1	B	228/238 (96%)	220 (96%)	8 (4%)	0	100	100
1	C	228/238 (96%)	220 (96%)	8 (4%)	0	100	100
1	D	227/238 (95%)	219 (96%)	8 (4%)	0	100	100
All	All	910/952 (96%)	878 (96%)	32 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	186/195 (95%)	185 (100%)	1 (0%)	81	77
1	B	187/195 (96%)	184 (98%)	3 (2%)	55	44
1	C	187/195 (96%)	184 (98%)	3 (2%)	55	44
1	D	186/195 (95%)	184 (99%)	2 (1%)	65	57
All	All	746/780 (96%)	737 (99%)	9 (1%)	63	55

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	171	GLU
1	B	27	GLU
1	B	124	LYS
1	B	171	GLU
1	C	27	GLU
1	C	70	VAL
1	C	171	GLU
1	D	70	VAL
1	D	171	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	107	HIS
1	B	219	ASN
1	C	28	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	EDO	C	302	-	3,3,3	0.37	0	2,2,2	0.45	0
2	4EH	B	301	-	26,28,28	1.06	3 (11%)	28,39,39	1.89	9 (32%)
3	EDO	D	302	-	3,3,3	0.38	0	2,2,2	0.45	0
2	4EH	A	301	-	26,28,28	1.13	2 (7%)	28,39,39	2.00	7 (25%)
2	4EH	C	301	-	26,28,28	1.15	2 (7%)	28,39,39	2.11	11 (39%)
2	4EH	D	301	-	26,28,28	1.14	2 (7%)	28,39,39	2.00	10 (35%)
3	EDO	A	302	-	3,3,3	0.38	0	2,2,2	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	C	302	-	-	0/1/1/1	-
2	4EH	B	301	-	-	1/9/21/21	0/4/4/4
3	EDO	D	302	-	-	0/1/1/1	-
2	4EH	A	301	-	-	1/9/21/21	0/4/4/4
2	4EH	C	301	-	-	1/9/21/21	0/4/4/4
2	4EH	D	301	-	-	1/9/21/21	0/4/4/4
3	EDO	A	302	-	-	0/1/1/1	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	301	4EH	C6'-S	-3.86	1.70	1.76
2	D	301	4EH	C6'-S	-3.83	1.70	1.76
2	C	301	4EH	C6'-S	-3.80	1.70	1.76
2	B	301	4EH	C6'-S	-3.41	1.71	1.76
2	D	301	4EH	C5'-S	-2.46	1.76	1.81
2	C	301	4EH	C5'-S	-2.34	1.76	1.81
2	A	301	4EH	C5'-S	-2.28	1.76	1.81
2	B	301	4EH	C5-N7	-2.07	1.34	1.37
2	B	301	4EH	C5'-S	-2.03	1.76	1.81

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	301	4EH	C5'-S-C6'	4.90	112.60	102.76
2	A	301	4EH	N1-C2-N3	-4.76	121.38	128.58
2	C	301	4EH	N1-C2-N3	-4.57	121.67	128.58
2	B	301	4EH	N1-C2-N3	-4.54	121.71	128.58
2	D	301	4EH	C5'-S-C6'	4.46	111.72	102.76
2	A	301	4EH	C5'-S-C6'	4.42	111.63	102.76
2	D	301	4EH	N1-C2-N3	-3.78	122.86	128.58
2	D	301	4EH	C9'-C6'-N2'	-3.49	118.46	121.70
2	C	301	4EH	C9'-C6'-N2'	-3.34	118.60	121.70
2	D	301	4EH	C10-C9-C4	3.34	132.34	126.58
2	B	301	4EH	C5'-S-C6'	3.27	109.32	102.76
2	C	301	4EH	C4-C9-C8	-3.18	103.12	105.92
2	B	301	4EH	C10-C9-C4	3.15	132.02	126.58
2	D	301	4EH	C4-C9-C8	-3.04	103.23	105.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	301	4EH	C10-N1'-C1'	2.99	119.13	113.39
2	A	301	4EH	C4-C5-N7	2.97	108.09	105.81
2	C	301	4EH	C10-N1'-C1'	2.86	118.88	113.39
2	B	301	4EH	C4-C9-C8	-2.83	103.42	105.92
2	A	301	4EH	C9'-C6'-N2'	-2.80	119.11	121.70
2	C	301	4EH	C10-C9-C4	2.77	131.37	126.58
2	B	301	4EH	C10-N1'-C1'	2.70	118.56	113.39
2	B	301	4EH	C9'-C6'-N2'	-2.68	119.21	121.70
2	C	301	4EH	C7'-C8'-N3'	-2.66	118.65	121.96
2	D	301	4EH	C10-N1'-C1'	2.55	118.27	113.39
2	A	301	4EH	C3'-C2'-N1'	-2.47	99.91	104.47
2	D	301	4EH	C7'-C8'-N3'	-2.44	118.92	121.96
2	B	301	4EH	C4-C5-N7	2.39	107.65	105.81
2	A	301	4EH	C10-C9-C4	2.38	130.69	126.58
2	C	301	4EH	C5-N7-C8	2.27	110.89	108.93
2	C	301	4EH	C4-C5-N7	2.23	107.53	105.81
2	D	301	4EH	C1'-N1'-C2'	2.20	107.32	104.12
2	D	301	4EH	C4-C5-N7	2.19	107.50	105.81
2	C	301	4EH	C8'-N3'-C9'	2.18	120.67	116.85
2	B	301	4EH	C1'-N1'-C2'	2.17	107.28	104.12
2	C	301	4EH	C3'-C2'-N1'	-2.16	100.48	104.47
2	D	301	4EH	C8'-N3'-C9'	2.16	120.63	116.85
2	B	301	4EH	C5-N7-C8	2.02	110.67	108.93

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	4EH	C9-C10-N1'-C1'
2	B	301	4EH	C9-C10-N1'-C1'
2	C	301	4EH	C9-C10-N1'-C1'
2	D	301	4EH	C9-C10-N1'-C1'


There are no ring outliers.

2 monomers are involved in 3 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	4EH	2	0
2	C	301	4EH	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,


bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



Ligand 4EH A 301







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	229/238 (96%)	0.71	24 (10%)	11 11	10, 20, 39, 55	1 (0%)
1	B	229/238 (96%)	1.16	38 (16%)	4 4	11, 25, 45, 51	1 (0%)
1	C	229/238 (96%)	1.17	52 (22%)	2 2	10, 28, 50, 60	1 (0%)
1	D	229/238 (96%)	0.99	39 (17%)	4 3	11, 24, 44, 58	0
All	All	916/952 (96%)	1.01	153 (16%)	4 4	10, 24, 45, 60	3 (0%)

All (153) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	40	ASP	5.7
1	C	0	GLY	5.7
1	C	39	LYS	5.3
1	D	128	LEU	5.1
1	C	199	GLY	4.9
1	A	138	ILE	4.9
1	D	136	LEU	4.7
1	C	38	TYR	4.4
1	C	20	LEU	4.3
1	C	40	ASP	4.2
1	B	128	LEU	3.9
1	D	0	GLY	3.9
1	C	200	GLU	3.9
1	B	137	ASN	3.8
1	C	23	TYR	3.7
1	D	138	ILE	3.7
1	B	133	ALA	3.6
1	D	133	ALA	3.6
1	B	20	LEU	3.6
1	C	201	LYS	3.5
1	C	136	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	156	ALA	3.4
1	B	26	ILE	3.4
1	B	132	VAL	3.4
1	B	131	GLU	3.3
1	A	202	ALA	3.3
1	C	198	ALA	3.3
1	C	21	LYS	3.3
1	D	127	ASN	3.3
1	B	136	LEU	3.3
1	C	82	PRO	3.3
1	C	30	ASN	3.3
1	A	22	ASP	3.2
1	D	132	VAL	3.2
1	C	41	HIS	3.2
1	B	119	ILE	3.2
1	D	41	HIS	3.2
1	C	18	GLU	3.2
1	D	129	ALA	3.1
1	A	209	PHE	3.1
1	B	156	ALA	3.1
1	D	38	TYR	3.1
1	A	155	GLU	3.0
1	D	221	VAL	3.0
1	A	133	ALA	3.0
1	D	224	MET	2.9
1	B	37	LYS	2.9
1	C	134	LYS	2.9
1	B	22	ASP	2.9
1	C	209	PHE	2.9
1	B	183	ALA	2.9
1	C	22	ASP	2.9
1	D	22	ASP	2.9
1	C	1	MET	2.9
1	D	21	LYS	2.8
1	A	128	LEU	2.8
1	D	130	LEU	2.8
1	D	222	LEU	2.8
1	A	201	LYS	2.8
1	C	19	THR	2.8
1	A	40	ASP	2.8
1	C	202	ALA	2.8
1	C	204	PHE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	0	GLY	2.8
1	D	19	THR	2.8
1	D	20	LEU	2.8
1	C	15	PRO	2.8
1	C	35	PHE	2.7
1	D	225	CYS	2.7
1	B	204	PHE	2.7
1	C	26	ILE	2.7
1	C	37	LYS	2.7
1	A	204	PHE	2.7
1	C	14	THR	2.7
1	C	16	LEU	2.7
1	C	85	GLU	2.7
1	D	131	GLU	2.7
1	A	139	LYS	2.7
1	A	205	ASP	2.6
1	B	70	VAL	2.6
1	B	124	LYS	2.6
1	D	227	LYS	2.6
1	A	130	LEU	2.6
1	C	141	ARG	2.6
1	B	142	ALA	2.6
1	C	36	ALA	2.6
1	A	136	LEU	2.6
1	B	42	GLU	2.6
1	B	138	ILE	2.6
1	C	123	GLU	2.6
1	D	218	ALA	2.6
1	C	138	ILE	2.5
1	C	42	GLU	2.5
1	B	225	CYS	2.5
1	C	34	TYR	2.5
1	B	134	LYS	2.5
1	C	130	LEU	2.5
1	D	26	ILE	2.5
1	C	223	LYS	2.5
1	D	223	LYS	2.5
1	B	164	ILE	2.5
1	D	135	GLU	2.5
1	A	135	GLU	2.4
1	C	212	ASN	2.4
1	D	42	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	222	LEU	2.4
1	C	228	LEU	2.4
1	D	84	LEU	2.4
1	D	228	LEU	2.4
1	B	159	ALA	2.4
1	B	116	GLU	2.4
1	A	219	ASN	2.4
1	D	137	ASN	2.4
1	C	153	CYS	2.4
1	A	134	LYS	2.4
1	C	84	LEU	2.4
1	C	227	LYS	2.4
1	D	39	LYS	2.4
1	C	28	HIS	2.4
1	D	139	LYS	2.3
1	B	40	ASP	2.3
1	C	226	GLU	2.3
1	B	121	THR	2.3
1	C	211	ILE	2.3
1	A	131	GLU	2.3
1	A	19	THR	2.3
1	D	1	MET	2.3
1	B	93	THR	2.3
1	D	27	GLU	2.2
1	D	200	GLU	2.2
1	D	85	GLU	2.2
1	B	91	TYR	2.2
1	B	21	LYS	2.2
1	B	14	THR	2.2
1	C	29	ALA	2.2
1	A	21	LYS	2.2
1	B	135	GLU	2.2
1	B	67	GLY	2.1
1	C	83	GLU	2.1
1	B	186	VAL	2.1
1	D	70	VAL	2.1
1	A	225	CYS	2.1
1	B	184	LEU	2.1
1	C	17	LEU	2.1
1	B	125	LEU	2.0
1	B	123	GLU	2.0
1	C	2	LYS	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	2	LYS	2.0
1	A	137	ASN	2.0
1	C	87	GLY	2.0
1	A	15	PRO	2.0
1	B	162	ARG	2.0
1	D	141	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	4EH	C	301	25/25	0.93	0.08	18,19,29,30	0
2	4EH	A	301	25/25	0.95	0.07	15,16,23,24	0
2	4EH	D	301	25/25	0.95	0.07	13,14,25,25	0
3	EDO	C	302	4/4	0.95	0.07	14,14,14,15	0
2	4EH	B	301	25/25	0.96	0.06	14,15,21,22	0
3	EDO	D	302	4/4	0.97	0.06	17,18,18,18	0
3	EDO	A	302	4/4	0.98	0.04	16,17,17,17	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 4EH C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 4EH A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 4EH D 301:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 4EH B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.