



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 07:56 PM UTC

PDB ID : 7AYB / pdb_00007ayb
Title : Crystal Structure of wild type human mitochondrial 2-Enoyl Thioester Reductase (MECR)
Authors : Rahman, M.T.; Koski, M.K.; Autio, K.J.; Kastaniotis, A.J.; Wierenga, R.K.; Hiltunen, J.K.
Deposited on : 2020-11-12
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	8% 86% 10% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2886 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Enoyl-[acyl-carrier-protein] reductase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	340	Total	C	N	O	S	0	9	0
			2657	1679	476	487	15			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	30	MET	-	initiating methionine	UNP Q9BV79
A	96	LEU	PHE	variant	UNP Q9BV79
A	374	LEU	-	expression tag	UNP Q9BV79
A	375	GLU	-	expression tag	UNP Q9BV79
A	376	HIS	-	expression tag	UNP Q9BV79
A	377	HIS	-	expression tag	UNP Q9BV79
A	378	HIS	-	expression tag	UNP Q9BV79
A	379	HIS	-	expression tag	UNP Q9BV79
A	380	HIS	-	expression tag	UNP Q9BV79
A	381	HIS	-	expression tag	UNP Q9BV79

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Cl	0	0
			4	4		

- Molecule 3 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is ACETIC ACID (CCD ID: ACY) (formula: $C_2H_4O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	200	Total	O	0	0
			200	200		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Enoyl-[acyl-carrier-protein] reductase, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	71.47Å 71.47Å 145.09Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	61.89 – 1.85 61.89 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.9 (61.89-1.85) 99.9 (61.89-1.85)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.86Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.189 , 0.223 0.203 , 0.194	Depositor DCC
R_{free} test set	3451 reflections (9.22%)	wwPDB-VP
Wilson B-factor (Å ²)	43.9	Xtriage
Anisotropy	0.001	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 52.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.016 for -h,-k,l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2886	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.08% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PO4, GOL, ACY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2734	0.83	0/3702

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2657	0	2735	20	0
2	A	4	0	0	1	0
3	A	5	0	0	0	0
4	A	12	0	16	3	0
5	A	8	0	6	1	0
6	A	200	0	0	2	0
All	All	2886	0	2757	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:115:VAL:HB	1:A:119:VAL:HG21	1.33	1.10
2:A:401:CL:CL	6:A:686:HOH:O	2.31	0.84
1:A:116:GLY:O	1:A:119:VAL:HG22	1.87	0.73
1:A:44[A]:ARG:NH1	1:A:351:GLN:OE1	2.28	0.66
1:A:263[B]:CYS:SG	6:A:698:HOH:O	2.54	0.65
1:A:311:TRP:HB3	4:A:407:GOL:H11	1.80	0.63
1:A:191:ALA:HA	5:A:408:ACY:H1	1.85	0.57
1:A:115:VAL:HB	1:A:119:VAL:CG2	2.22	0.55
1:A:54:PRO:HG2	1:A:361:MET:HE1	1.91	0.53
1:A:123[B]:LYS:HE2	1:A:126:ASP:OD1	2.13	0.48
1:A:167:ASN:HB2	1:A:168:PRO:HD3	1.95	0.48
1:A:247:MET:HA	1:A:250:PHE:HB2	1.95	0.47
1:A:218:ARG:HD3	1:A:220:ASP:OD1	2.14	0.47
1:A:170:THR:HG22	1:A:174:MET:HE2	1.97	0.47
1:A:115:VAL:CB	1:A:119:VAL:HG21	2.23	0.44
1:A:248:LYS:HA	1:A:275:GLN:OE1	2.17	0.44
1:A:314:GLN:HB3	4:A:407:GOL:H11	2.01	0.42
1:A:127:TRP:CE3	1:A:154:SER:HA	2.55	0.41
1:A:365:ILE:O	1:A:366:SER:C	2.62	0.41
1:A:311:TRP:HB3	4:A:407:GOL:C1	2.51	0.40
1:A:254:MET:HB2	1:A:255:PRO:HD2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/352 (99%)	337 (97%)	10 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/291 (100%)	283 (97%)	9 (3%)	35	20

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	ASN
1	A	93	ASN
1	A	216	ARG
1	A	227	ARG
1	A	247	MET
1	A	303	LYS
1	A	360	SER
1	A	368[A]	LYS
1	A	368[B]	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	91	GLN
1	A	167	ASN
1	A	249	ASN
1	A	314	GLN
1	A	339	GLN
1	A	347	GLN
1	A	376	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	ACY	A	408	-	3,3,3	0.96	0	3,3,3	0.89	0
4	GOL	A	406	-	5,5,5	0.11	0	5,5,5	0.31	0
3	PO4	A	405	-	4,4,4	0.71	0	6,6,6	0.45	0
4	GOL	A	407	-	5,5,5	0.17	0	5,5,5	0.46	0
5	ACY	A	409	-	3,3,3	1.01	0	3,3,3	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	407	-	-	2/4/4/4	-
4	GOL	A	406	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	406	GOL	C1-C2-C3-O3
4	A	407	GOL	O1-C1-C2-C3
4	A	406	GOL	O2-C2-C3-O3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	407	GOL	O1-C1-C2-O2
4	A	406	GOL	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	408	ACY	1	0
4	A	407	GOL	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/352 (96%)	0.61	27 (7%) 18 19	26, 53, 96, 122	9 (2%)

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	365	ILE	5.6
1	A	364	PHE	4.3
1	A	345	CYS	3.9
1	A	247	MET	3.4
1	A	250	PHE	3.1
1	A	227	ARG	2.8
1	A	366	SER	2.7
1	A	363	PRO	2.6
1	A	55	ALA	2.5
1	A	216	ARG	2.4
1	A	367	SER	2.4
1	A	113	VAL	2.3
1	A	43	VAL	2.3
1	A	65	LEU	2.3
1	A	217	ASP	2.2
1	A	380	HIS	2.2
1	A	67	ALA	2.2
1	A	362	LYS	2.2
1	A	381	HIS	2.1
1	A	289	ALA	2.1
1	A	119	VAL	2.1
1	A	93	ASN	2.1
1	A	344	ALA	2.1
1	A	42	ARG	2.0
1	A	221	ILE	2.0
1	A	378	HIS	2.0
1	A	218	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PO4	A	405	5/5	0.50	0.18	108,110,114,120	0
5	ACY	A	409	4/4	0.53	0.29	81,82,85,86	0
5	ACY	A	408	4/4	0.82	0.19	56,65,66,67	0
4	GOL	A	407	6/6	0.87	0.16	55,58,65,66	0
4	GOL	A	406	6/6	0.88	0.17	73,74,76,76	0
2	CL	A	402	1/1	0.96	0.15	51,51,51,51	1
2	CL	A	403	1/1	0.97	0.10	49,49,49,49	0
2	CL	A	401	1/1	0.98	0.11	56,56,56,56	0
2	CL	A	404	1/1	0.99	0.11	52,52,52,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.