



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 05:03 PM UTC

PDB ID : 9AY8 / pdb_00009ay8
Title : Structure of the A type blood alpha-D-galactosamine galactosaminidase from
Flavonifractor plautii
Authors : Worrall, L.J.; Strynadka, N.C.J.
Deposited on : 2024-03-07
Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	672	92% 5% .
1	B	672	90% 7% .
1	C	672	92% 5% .
1	D	672	% 92% 5% .
1	E	672	% 92% 5% .

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called A type blood alpha-D-galactosamine galactosaminidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	654	Total 5067	C 3200	N 832	O 1011	S 24	0	0	0
1	B	654	Total 5067	C 3200	N 832	O 1011	S 24	0	0	0
1	C	654	Total 5067	C 3200	N 832	O 1011	S 24	0	0	0
1	D	654	Total 5067	C 3200	N 832	O 1011	S 24	0	0	0
1	E	654	Total 5072	C 3204	N 832	O 1011	S 25	0	1	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	624	ALA	GLU	conflict	UNP P0DTR5
A	625	ALA	LYS	conflict	UNP P0DTR5
B	624	ALA	GLU	conflict	UNP P0DTR5
B	625	ALA	LYS	conflict	UNP P0DTR5
C	624	ALA	GLU	conflict	UNP P0DTR5
C	625	ALA	LYS	conflict	UNP P0DTR5
D	624	ALA	GLU	conflict	UNP P0DTR5
D	625	ALA	LYS	conflict	UNP P0DTR5
E	624	ALA	GLU	conflict	UNP P0DTR5
E	625	ALA	LYS	conflict	UNP P0DTR5

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0

Continued on next page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0
2	E	1	Total Zn 1 1	0	0

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 3 | A | 1 | Total Cl
1 1 | 0 | 0 |
| 3 | B | 1 | Total Cl
1 1 | 0 | 0 |
| 3 | C | 1 | Total Cl
1 1 | 0 | 0 |
| 3 | D | 1 | Total Cl
1 1 | 0 | 0 |
| 3 | E | 1 | Total Cl
1 1 | 0 | 0 |

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 4 | A | 1 | Total Co
1 1 | 0 | 0 |
| 4 | B | 1 | Total Co
1 1 | 0 | 0 |
| 4 | C | 1 | Total Co
1 1 | 0 | 0 |
| 4 | D | 1 | Total Co
1 1 | 0 | 0 |
| 4 | E | 1 | Total Co
1 1 | 0 | 0 |

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|-----------------|---------|---------|
| 5 | A | 1 | Total Mn
1 1 | 0 | 0 |
| 5 | B | 1 | Total Mn
1 1 | 0 | 0 |



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total Mn 1 1	0	0
5	D	1	Total Mn 1 1	0	0
5	E	1	Total Mn 1 1	0	0

-
- The chemical structure of EPE (1,4-bis(2-hydroxyethyl)piperazine-2,5-dithione) is shown. The molecule consists of a central piperazine ring with two nitrogen atoms (N1 and N4) and four carbon atoms (C2, C3, C5, C6). Each nitrogen atom is substituted with a 2-hydroxyethyl group. The sulfur atoms (S1 and S2) are part of the dithione group, and the hydroxyl groups (OH1 and OH2) are attached to the terminal carbon atoms (C9 and C10). The atom labels are: N1, N4, C2, C3, C5, C6, C9, C10, S1, S2, OH1, OH2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
6	A	1	Total 15	C 8	N 2	O 4	S 1	0	0
6	B	1	Total 15	C 8	N 2	O 4	S 1	0	0

- | Mol | Chain | Residues | Atoms | ZeroOcc | AltConf |
|-----|-------|----------|--------------------|---------|---------|
| 7 | A | 763 | Total O
763 763 | 0 | 0 |
| 7 | B | 776 | Total O
776 776 | 0 | 0 |
| 7 | C | 536 | Total O
536 536 | 0 | 0 |
| 7 | D | 621 | Total O
621 621 | 0 | 0 |



WORLD WIDE PDB
PROTEIN DATA BANK

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	E	641	Total 641	O 641	0	0



● Molecule 1: A type blood alpha-D-galactosamine galactosaminidase



i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	102.69Å 164.57Å 131.65Å 90.00° 90.30° 90.00°	Depositor
Resolution (Å)	39.44 – 2.00 39.44 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.7 (39.44-2.00) 98.7 (39.44-2.00)	Depositor EDS
R _{merge}	0.06	Depositor
R _{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
R, R _{free}	0.168 , 0.193 0.176 , 0.201	Depositor DCC
R _{free} test set	14518 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.229	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.014 for h,-k,-l	Xtriage
F _o ,F _c correlation	0.96	EDS
Total number of atoms	28727	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage’s analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 of 11 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	660	ARG	Sidechain
1	A	684	ARG	Sidechain
1	B	414	ARG	Sidechain
1	B	660	ARG	Sidechain
1	B	684	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5067	0	4765	22	0
1	B	5067	0	4765	29	0
1	C	5067	0	4765	17	0
1	D	5067	0	4764	20	0
1	E	5072	0	4773	27	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
4	A	1	0	0	1	0
4	B	1	0	0	0	0
4	C	1	0	0	1	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
5	E	1	0	0	0	0
6	A	15	0	17	1	0
6	B	15	0	17	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	A	763	0	0	8	0
7	B	776	0	0	11	0
7	C	536	0	0	5	1
7	D	621	0	0	5	0
7	E	641	0	0	8	1
All	All	28727	0	23866	117	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 117 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:626:ASP:HB3	7:D:1186:HOH:O	1.35	1.27
1:A:626:ASP:HB3	7:A:1309:HOH:O	1.58	1.03
1:B:58:ASP:HB3	7:B:1452:HOH:O	1.68	0.92
1:A:513:CYS:HG	4:A:703:CO:CO	0.54	0.88
1:E:178:ASN:HD22	1:E:206:ASN:HD21	1.29	0.80

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:C:842:HOH:O	7:E:1305:HOH:O[1_556]	1.79	0.41

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	652/672 (97%)	629 (96%)	22 (3%)	1 (0%)	43 42
1	B	652/672 (97%)	630 (97%)	21 (3%)	1 (0%)	43 42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	652/672 (97%)	631 (97%)	20 (3%)	1 (0%)	43	42
1	D	652/672 (97%)	629 (96%)	22 (3%)	1 (0%)	43	42
1	E	653/672 (97%)	631 (97%)	21 (3%)	1 (0%)	43	42
All	All	3261/3360 (97%)	3150 (97%)	106 (3%)	5 (0%)	43	42

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	531	ALA
1	B	531	ALA
1	C	531	ALA
1	D	531	ALA
1	E	531	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	536/546 (98%)	534 (100%)	2 (0%)	84	89
1	B	536/546 (98%)	532 (99%)	4 (1%)	76	82
1	C	536/546 (98%)	531 (99%)	5 (1%)	70	78
1	D	536/546 (98%)	528 (98%)	8 (2%)	57	64
1	E	537/546 (98%)	530 (99%)	7 (1%)	61	68
All	All	2681/2730 (98%)	2655 (99%)	26 (1%)	68	75

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	643	THR
1	D	683	THR
1	E	426	ASN
1	D	682	ASN
1	D	684	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	178	ASN
1	E	120	GLN
1	E	444	GLN
1	E	178	ASN
1	D	444	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 20 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EPE	A	705	-	15,15,15	1.09	1 (6%)	19,20,20	1.06	2 (10%)
6	EPE	B	705	-	15,15,15	1.29	1 (6%)	19,20,20	1.13	2 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EPE	A	705	-	-	5/9/19/19	0/1/1/1
6	EPE	B	705	-	-	5/9/19/19	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	705	EPE	O1S-S	4.61	1.58	1.45
6	A	705	EPE	O1S-S	4.02	1.56	1.45

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	A	705	EPE	O1S-S-C10	-3.03	102.15	106.73
6	A	705	EPE	O3S-S-O2S	2.53	117.72	111.40
6	B	705	EPE	O3S-S-O2S	2.22	116.97	111.40
6	B	705	EPE	O1S-S-C10	-2.22	103.37	106.73

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	705	EPE	C10-C9-N1-C2
6	A	705	EPE	C9-C10-S-O2S
6	A	705	EPE	C9-C10-S-O3S
6	B	705	EPE	C10-C9-N1-C2
6	B	705	EPE	C10-C9-N1-C6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	705	EPE	1	0
6	B	705	EPE	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	654/672 (97%)	-0.47	1 (0%) 91 90	22, 32, 56, 74	0
1	B	654/672 (97%)	-0.49	1 (0%) 91 90	21, 30, 52, 68	0
1	C	654/672 (97%)	-0.25	1 (0%) 91 90	26, 39, 68, 89	0
1	D	654/672 (97%)	-0.23	5 (0%) 82 82	26, 40, 66, 95	0
1	E	654/672 (97%)	-0.23	5 (0%) 82 82	23, 36, 65, 89	1 (0%)
All	All	3270/3360 (97%)	-0.33	13 (0%) 88 88	21, 35, 63, 95	1 (0%)

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	697	SER	2.8
1	D	683	THR	2.6
1	E	153	SER	2.6
1	E	267	ILE	2.6
1	E	137	ILE	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EPE	A	705	15/15	0.77	0.19	67,91,117,142	0
6	EPE	B	705	15/15	0.89	0.14	53,65,74,83	0
4	CO	C	703	1/1	0.98	0.13	58,58,58,58	0
4	CO	E	703	1/1	0.98	0.11	52,52,52,52	0
4	CO	A	703	1/1	0.98	0.14	55,55,55,55	0
4	CO	B	703	1/1	0.98	0.15	52,52,52,52	0
2	ZN	D	701	1/1	0.99	0.02	31,31,31,31	0
2	ZN	E	701	1/1	0.99	0.03	33,33,33,33	0
4	CO	D	703	1/1	0.99	0.13	59,59,59,59	0
3	CL	C	702	1/1	0.99	0.06	30,30,30,30	0
5	MN	A	704	1/1	0.99	0.07	36,36,36,36	0
5	MN	B	704	1/1	0.99	0.06	41,41,41,41	0
5	MN	C	704	1/1	0.99	0.10	45,45,45,45	0
5	MN	D	704	1/1	0.99	0.06	43,43,43,43	0
5	MN	E	704	1/1	0.99	0.05	44,44,44,44	0
3	CL	D	702	1/1	0.99	0.04	29,29,29,29	0
2	ZN	C	701	1/1	0.99	0.03	32,32,32,32	0
2	ZN	B	701	1/1	1.00	0.01	28,28,28,28	0
2	ZN	A	701	1/1	1.00	0.01	31,31,31,31	0
3	CL	E	702	1/1	1.00	0.04	26,26,26,26	0
3	CL	A	702	1/1	1.00	0.03	25,25,25,25	0
3	CL	B	702	1/1	1.00	0.03	26,26,26,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.