



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 08:55 PM UTC

PDB ID : 9AYA / pdb_00009aya
Title : Crystal structure of CRAF/MEK complex with NST-628 and active RAF dimer
Authors : Quade, B.; Huang, X.
Deposited on : 2024-03-07
Resolution : 2.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	5% 84% 11% 5%
1	C	280	14% 83% 9% 6%
2	B	310	6% 85% 11% • •
2	D	310	3% 88% 7% 5%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8958 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RAF proto-oncogene serine/threonine-protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2124	1365	364	381	14			
1	C	263	Total	C	N	O	S	0	0	0
			2086	1336	359	377	14			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	336	GLY	-	expression tag	UNP P04049
A	340	ASP	TYR	engineered mutation	UNP P04049
A	341	ASP	TYR	engineered mutation	UNP P04049
C	336	GLY	-	expression tag	UNP P04049
C	340	ASP	TYR	engineered mutation	UNP P04049
C	341	ASP	TYR	engineered mutation	UNP P04049

- Molecule 2 is a protein called Dual specificity mitogen-activated protein kinase kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	299	Total	C	N	O	S	0	1	0
			2255	1433	380	426	16			
2	D	294	Total	C	N	O	S	0	1	0
			2312	1474	396	426	16			

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	36	GLY	-	expression tag	UNP Q02750
B	218	ALA	SER	engineered mutation	UNP Q02750
B	222	ALA	SER	engineered mutation	UNP Q02750
B	264	GLY	-	linker	UNP Q02750
B	303	SER	-	linker	UNP Q02750
B	304	GLY	-	linker	UNP Q02750

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	305	SER	-	linker	UNP Q02750
B	306	GLY	-	linker	UNP Q02750
B	307	SER	-	linker	UNP Q02750
D	36	GLY	-	expression tag	UNP Q02750
D	218	ALA	SER	engineered mutation	UNP Q02750
D	222	ALA	SER	engineered mutation	UNP Q02750
D	264	GLY	-	linker	UNP Q02750
D	303	SER	-	linker	UNP Q02750
D	304	GLY	-	linker	UNP Q02750
D	305	SER	-	linker	UNP Q02750
D	306	GLY	-	linker	UNP Q02750
D	307	SER	-	linker	UNP Q02750

- Molecule 3 is N-[3-fluoro-4-({7-[(3-fluoropyridin-2-yl)oxy]-4-methyl-2-oxo-2H-1-benzopyran-3-yl)methyl}pyridin-2-yl]-N'-methylsulfuric diamide (CCD ID: A1AHE) (formula: $C_{22}H_{18}F_2N_4O_5S$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	B	1	Total	C	F	N	O	S	0	0
			34	22	2	4	5	1		
3	D	1	Total	C	F	N	O	S	0	0
			34	22	2	4	5	1		

- Molecule 4 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
4	D	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		
5	D	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	13	Total	O	0	0
			13	13		
6	B	9	Total	O	0	0
			9	9		
6	C	12	Total	O	0	0
			12	12		
6	D	15	Total	O	0	0
			15	15		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RAF proto-oncogene serine/threonine-protein kinase




- Molecule 1: RAF proto-oncogene serine/threonine-protein kinase




- Molecule 2: Dual specificity mitogen-activated protein kinase kinase 1



- Molecule 2: Dual specificity mitogen-activated protein kinase kinase 1





4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	149.72Å 58.20Å 162.35Å 90.00° 95.59° 90.00°	Depositor
Resolution (Å)	57.64 – 2.59 57.64 – 2.59	Depositor EDS
% Data completeness (in resolution range)	99.8 (57.64-2.59) 99.9 (57.64-2.59)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 2.58Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487, REFMAC 5	Depositor
R, R_{free}	0.212 , 0.257 0.212 , 0.257	Depositor DCC
R_{free} test set	2115 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	49.2	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8958	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, A1AHE, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.17	0/2173	0.35	0/2944
1	C	0.18	0/2133	0.35	0/2885
2	B	0.14	0/2297	0.31	0/3111
2	D	0.17	0/2352	0.32	0/3164
All	All	0.16	0/8955	0.33	0/12104

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2124	0	2107	15	0
1	C	2086	0	2047	20	0
2	B	2255	0	2168	20	0
2	D	2312	0	2331	12	0
3	B	34	0	0	0	0
3	D	34	0	0	0	0
4	B	31	0	13	0	0
4	D	31	0	13	0	0
5	B	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	1	0	0	0	0
6	A	13	0	0	0	0
6	B	9	0	0	0	0
6	C	12	0	0	2	0
6	D	15	0	0	0	0
All	All	8958	0	8679	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:593:LYS:NZ	6:C:701:HOH:O	1.97	0.94
1:C:512:MET:HE3	1:C:516:VAL:HG12	1.71	0.70
1:A:542:MET:HE3	1:A:581:MET:HE2	1.74	0.68
1:C:413:THR:HA	1:C:417:LEU:HD23	1.80	0.64
1:A:396:VAL:HG13	1:A:464:ILE:HD12	1.80	0.63
1:C:515:GLU:OE2	1:C:596:ARG:NH2	2.33	0.62
1:A:597:PRO:HG2	1:A:602:ILE:HD11	1.82	0.61
1:C:364:TYR:CE2	1:C:376:ILE:HG23	2.34	0.61
2:D:188:HIS:CD2	2:D:209:PHE:HB3	2.35	0.61
1:A:512:MET:HE3	1:A:516:VAL:HG12	1.82	0.60
2:B:71:ILE:HD11	2:B:86:SER:HB2	1.83	0.59
2:D:71:ILE:HD11	2:D:86:SER:HB2	1.85	0.58
2:B:251:LEU:HD11	2:B:263:ILE:HD11	1.86	0.57
2:D:308:MET:HG2	2:D:312:GLU:HB3	1.86	0.57
2:B:341:CYS:O	2:B:349:ARG:HD2	2.05	0.56
1:A:465:ILE:HD11	1:A:493:LYS:HD3	1.87	0.56
1:C:553:ASN:HD21	2:D:230:MET:HE2	1.72	0.55
1:A:467:ARG:HG2	1:A:525:PHE:CD1	2.43	0.54
1:A:538:LEU:HB3	1:A:585:VAL:HG22	1.91	0.53
2:B:74:LEU:HD21	2:B:84:LYS:HB2	1.92	0.51
2:D:341:CYS:O	2:D:349:ARG:HD2	2.13	0.49
1:C:374:VAL:HG13	1:C:420:VAL:HG22	1.95	0.49
1:A:543:THR:HG22	1:A:573:LEU:HA	1.94	0.49
1:C:467:ARG:HD2	1:C:525:PHE:CG	2.48	0.48
2:B:158:ALA:HB2	2:B:378:THR:HG21	1.95	0.48
2:B:188:HIS:CD2	2:B:209:PHE:HB3	2.48	0.48
1:C:354:ARG:HA	1:C:363:VAL:O	2.14	0.48
1:A:554:ARG:NH1	2:B:223:PHE:O	2.46	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:350:MET:HE3	1:C:350:MET:HB2	1.82	0.47
1:C:524:PRO:O	6:C:701:HOH:O	2.20	0.47
2:B:134:TYR:HD1	2:B:139:ILE:HG12	1.79	0.46
2:D:134:TYR:OH	2:D:137:GLY:HA2	2.16	0.46
1:C:360:PHE:HB2	1:C:376:ILE:HD11	1.98	0.46
2:B:188:HIS:N	2:B:245:ASP:OD2	2.44	0.45
2:D:175:LYS:HG3	2:D:356:MET:HE1	1.99	0.45
1:C:364:TYR:HB2	1:C:374:VAL:O	2.16	0.45
2:B:233:GLU:OE2	2:B:349:ARG:NH2	2.49	0.45
1:C:474:ILE:HG23	1:C:482:VAL:HG13	1.99	0.45
1:A:474:ILE:HG23	1:A:482:VAL:HG13	1.99	0.44
2:B:343:ILE:O	2:B:349:ARG:HD3	2.17	0.44
1:A:515:GLU:OE2	1:A:596:ARG:NH2	2.48	0.44
1:C:456:MET:SD	1:C:469:MET:HE3	2.58	0.44
1:A:454:GLN:HA	1:A:603:LEU:HD11	1.99	0.44
2:B:60:VAL:HG11	2:B:94:MET:HE1	2.00	0.43
2:B:224:VAL:HG13	2:B:310:ILE:HG21	1.99	0.43
2:D:230:MET:HE1	2:D:234:ARG:CZ	2.48	0.43
1:C:374:VAL:HG12	1:C:418:ALA:HB1	1.98	0.43
1:C:376:ILE:H	1:C:376:ILE:HG13	1.63	0.43
2:D:127:VAL:HG21	2:D:197:LEU:HD12	2.00	0.43
2:B:146:MET:HG3	2:B:197:LEU:HB3	2.01	0.43
1:A:518:ARG:HB3	1:A:520:GLN:HG3	2.00	0.42
2:D:156:LYS:HB2	2:D:156:LYS:HE2	1.78	0.42
1:A:589:VAL:O	1:A:589:VAL:HG12	2.19	0.42
1:C:343:GLU:HG3	1:C:344:ILE:N	2.33	0.42
2:B:105:PRO:HA	2:B:108:ARG:HG2	2.00	0.42
2:D:70:LYS:HE3	2:D:70:LYS:HB2	1.85	0.42
2:B:326:PRO:HD2	2:B:330:PHE:HE2	1.83	0.42
2:B:134:TYR:OH	2:B:137:GLY:HA2	2.20	0.41
1:C:515:GLU:CD	1:C:596:ARG:HH22	2.27	0.41
2:B:202:GLY:HA2	2:B:371:PHE:HD2	1.86	0.41
1:A:382:PRO:HB2	1:A:387:PHE:CZ	2.56	0.41
2:B:112:ILE:O	2:B:116:GLN:HG2	2.21	0.41
2:B:310:ILE:O	2:B:314:LEU:HG	2.20	0.41
1:C:543:THR:HG22	1:C:573:LEU:HA	2.04	0.40
2:D:202:GLY:HA2	2:D:371:PHE:HD2	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	175/280 (62%)	170 (97%)	4 (2%)	1 (1%)	21	42
1	C	257/280 (92%)	243 (95%)	14 (5%)	0	100	100
2	B	225/310 (73%)	220 (98%)	5 (2%)	0	100	100
2	D	287/310 (93%)	283 (99%)	3 (1%)	1 (0%)	36	58
All	All	944/1180 (80%)	916 (97%)	26 (3%)	2 (0%)	43	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	486	ASP
2	D	262	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/249 (93%)	226 (98%)	5 (2%)	45	72
1	C	224/249 (90%)	218 (97%)	6 (3%)	39	67
2	B	235/265 (89%)	230 (98%)	5 (2%)	47	73
2	D	253/265 (96%)	252 (100%)	1 (0%)	84	93
All	All	943/1028 (92%)	926 (98%)	17 (2%)	51	76

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	376	ILE
1	A	415	ASP
1	A	493	LYS
1	A	495	ARG
1	A	613	LEU
2	B	50	LEU
2	B	58	GLN
2	B	84	LYS
2	B	224	VAL
2	B	382	ASN
1	C	343	GLU
1	C	350	MET
1	C	353	THR
1	C	374	VAL
1	C	376	ILE
1	C	609	LEU
2	D	64	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	369	HIS
1	C	556	GLN
2	D	116	GLN
2	D	145	HIS
2	D	236	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	A1AHE	B	401	-	36,37,37	1.29	2 (5%)	47,54,54	1.70	9 (19%)
3	A1AHE	D	401	-	36,37,37	1.29	1 (2%)	47,54,54	1.69	9 (19%)
4	ANP	B	402	5	33,33,33	0.95	3 (9%)	45,52,52	0.67	0
4	ANP	D	402	5	33,33,33	0.90	3 (9%)	45,52,52	0.83	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	A1AHE	B	401	-	-	0/15/16/16	0/4/4/4
3	A1AHE	D	401	-	-	0/15/16/16	0/4/4/4
4	ANP	B	402	5	-	4/18/38/38	0/3/3/3
4	ANP	D	402	5	-	1/18/38/38	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	401	A1AHE	S03-N02	5.08	1.67	1.61
3	D	401	A1AHE	S03-N02	4.78	1.67	1.61
4	D	402	ANP	PG-O1G	2.92	1.50	1.46
4	B	402	ANP	PG-O1G	2.82	1.50	1.46
3	B	401	A1AHE	C28-C29	2.72	1.43	1.38
4	B	402	ANP	PG-N3B	2.47	1.69	1.63
4	B	402	ANP	PB-O1B	2.36	1.49	1.46
4	D	402	ANP	PG-N3B	2.22	1.69	1.63
4	D	402	ANP	PB-O1B	2.19	1.49	1.46

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	401	A1AHE	C31-C13-C14	-6.37	115.92	120.35
3	B	401	A1AHE	C31-C13-C14	-5.56	116.48	120.35
3	B	401	A1AHE	C11-C12-C13	3.79	120.34	114.05
3	D	401	A1AHE	O30-C31-O32	-3.67	111.39	116.22
3	B	401	A1AHE	O30-C31-O32	-3.56	111.53	116.22
3	D	401	A1AHE	C11-C12-C13	3.46	119.78	114.05
4	D	402	ANP	O1B-PB-N3B	-3.41	106.74	111.77
3	D	401	A1AHE	O30-C31-C13	3.36	122.04	117.86
3	B	401	A1AHE	O30-C31-C13	3.30	121.96	117.86
3	B	401	A1AHE	O20-C21-C26	2.73	121.08	117.57
3	D	401	A1AHE	C17-C16-C29	-2.56	114.85	118.21
3	B	401	A1AHE	C17-C16-C29	-2.42	115.03	118.21
3	D	401	A1AHE	C10-C11-C33	-2.32	114.75	116.46
3	B	401	A1AHE	C10-C11-C33	-2.23	114.82	116.46
3	B	401	A1AHE	C33-C07-N08	2.19	120.97	119.50
3	D	401	A1AHE	C33-C07-N08	2.08	120.89	119.50
3	D	401	A1AHE	C29-O30-C31	-2.03	120.32	122.26
3	B	401	A1AHE	C29-O30-C31	-2.01	120.34	122.26
3	D	401	A1AHE	O20-C21-C26	2.00	120.14	117.57

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	402	ANP	PB-N3B-PG-O1G
4	B	402	ANP	PG-N3B-PB-O1B
4	D	402	ANP	PB-N3B-PG-O1G
4	B	402	ANP	O4'-C4'-C5'-O5'
4	B	402	ANP	C3'-C4'-C5'-O5'


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/280 (95%)	0.26	13 (4%) 35 29	35, 47, 85, 123	0
1	C	263/280 (93%)	0.58	38 (14%) 6 5	34, 50, 102, 140	0
2	B	299/310 (96%)	0.62	18 (6%) 27 22	24, 63, 96, 107	1 (0%)
2	D	294/310 (94%)	0.06	10 (3%) 48 42	19, 47, 77, 98	1 (0%)
All	All	1123/1180 (95%)	0.38	79 (7%) 22 18	19, 53, 92, 140	2 (0%)

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	376	ILE	5.8
1	C	416	ASN	5.7
1	C	362	THR	5.4
1	C	382	PRO	4.8
1	C	360	PHE	4.7
2	D	264	GLY	4.7
2	B	381	LEU	4.6
2	D	76	ALA	4.4
1	C	415	ASP	4.1
1	C	413	THR	3.9
1	A	380	VAL	3.9
1	C	354	ARG	3.8
1	C	383	THR	3.6
1	C	412	MET	3.6
1	C	364	TYR	3.5
2	B	264	GLY	3.5
1	A	381	ASP	3.5
1	C	496	TRP	3.5
2	B	42	LEU	3.4
1	C	613	LEU	3.3
1	C	384	PRO	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	355	ILE	3.2
1	C	504	GLN	3.0
1	C	361	GLY	3.0
1	C	417	LEU	2.8
2	B	307	SER	2.8
2	B	382	ASN	2.8
2	B	379	ILE	2.7
1	C	390	PHE	2.7
1	C	495	ARG	2.7
1	C	414	LYS	2.7
2	D	221	ASN	2.6
1	C	576	ASN	2.6
2	B	316	TYR	2.6
1	C	505	PRO	2.5
2	D	239	HIS	2.5
1	A	506	THR	2.5
2	B	134	TYR	2.5
1	C	391	ARG	2.5
1	C	387	PHE	2.5
2	D	238	THR	2.5
1	A	376	ILE	2.4
1	C	363	VAL	2.4
1	C	498	GLY	2.4
1	A	379	VAL	2.4
2	D	75	GLY	2.3
2	B	263	ILE	2.3
1	A	479	GLY	2.3
2	D	379	ILE	2.3
2	B	44	GLU	2.3
2	B	63	LEU	2.3
1	C	385	GLU	2.3
2	D	382	ASN	2.3
1	C	508	SER	2.3
1	A	413	THR	2.2
2	B	164	GLN	2.2
1	C	350	MET	2.2
1	C	352	SER	2.2
2	D	225	GLY	2.2
1	A	614	PRO	2.2
1	C	411	TYR	2.2
1	C	341	ASP	2.2
1	A	500	GLN	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	506	THR	2.1
1	C	494	SER	2.1
2	B	239	HIS	2.1
2	B	92	LEU	2.1
1	C	353	THR	2.1
1	C	386	GLN	2.1
1	A	576	ASN	2.1
2	B	353	LYS	2.1
2	D	316	TYR	2.1
1	A	384	PRO	2.1
2	B	130	TYR	2.0
1	A	387	PHE	2.0
2	B	133	PHE	2.0
1	C	418	ALA	2.0
2	B	328	GLY	2.0
1	C	612	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ANP	B	402	31/31	0.94	0.08	56,65,81,109	0
4	ANP	D	402	31/31	0.94	0.09	30,48,99,131	0
3	A1AHE	B	401	34/34	0.95	0.09	40,48,57,58	0
5	MG	B	403	1/1	0.95	0.05	51,51,51,51	0
3	A1AHE	D	401	34/34	0.97	0.06	29,37,45,46	0
5	MG	D	403	1/1	0.98	0.05	51,51,51,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around ANP D 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around A1AHE B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1AHE D 401:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.