



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 20, 2026 – 04:16 AM UTC

PDB ID : 3AZF / pdb_00003azf
Title : Crystal Structure of Human Nucleosome Core Particle Containing H3K79Q mutation
Authors : Iwasaki, W.; Tachiwana, H.; Kawaguchi, K.; Shibata, T.; Kagawa, W.; Kurumizaka, H.
Deposited on : 2011-05-25
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	3% 49% 19% • 30%
1	E	139	2% 45% 22% • • 29%
2	B	106	• 56% 17% • 26%
2	F	106	3% 60% 17% • 21%
3	C	133	53% 23% 6% 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	133	% 59%16%•22%
4	D	129	% 43%23%6%27%
4	H	129	% 55%12%•29%
5	I	146	16% 14%84%••
5	J	146	18% 12%86%••

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 12152 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	97	Total	C	N	O	S	0	0	0
			801	504	155	138	4			
1	E	99	Total	C	N	O	S	0	0	0
			816	513	158	141	4			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P68431
A	-2	SER	-	expression tag	UNP P68431
A	-1	HIS	-	expression tag	UNP P68431
A	79	GLN	LYS	engineered mutation	UNP P68431
E	-3	GLY	-	expression tag	UNP P68431
E	-2	SER	-	expression tag	UNP P68431
E	-1	HIS	-	expression tag	UNP P68431
E	79	GLN	LYS	engineered mutation	UNP P68431

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	78	Total	C	N	O	S	0	0	0
			619	391	120	107	1			
2	F	84	Total	C	N	O	S	0	0	0
			673	424	133	115	1			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P62805
B	-2	SER	-	expression tag	UNP P62805
B	-1	HIS	-	expression tag	UNP P62805
F	-3	GLY	-	expression tag	UNP P62805

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	-2	SER	-	expression tag	UNP P62805
F	-1	HIS	-	expression tag	UNP P62805

- Molecule 3 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	108	Total	C	N	O	0	0	0
			835	526	165	144			
3	G	104	Total	C	N	O	0	0	0
			805	508	157	140			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	expression tag	UNP P04908
C	-2	SER	-	expression tag	UNP P04908
C	-1	HIS	-	expression tag	UNP P04908
G	-3	GLY	-	expression tag	UNP P04908
G	-2	SER	-	expression tag	UNP P04908
G	-1	HIS	-	expression tag	UNP P04908

- Molecule 4 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	94	Total	C	N	O	S	0	0	0
			736	462	134	138	2			
4	H	91	Total	C	N	O	S	0	0	0
			714	450	128	134	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	GLY	-	expression tag	UNP P06899
D	-2	SER	-	expression tag	UNP P06899
D	-1	HIS	-	expression tag	UNP P06899
H	-3	GLY	-	expression tag	UNP P06899
H	-2	SER	-	expression tag	UNP P06899
H	-1	HIS	-	expression tag	UNP P06899

- Molecule 5 is a DNA chain called 146-MER DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	I	145	Total 2970	C 1421	N 538	O 867	P 144	0	0	0
5	J	145	Total 2969	C 1421	N 535	O 869	P 144	0	0	0

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total 1	Cl 1	0	0
6	C	1	Total 1	Cl 1	0	0
6	E	1	Total 1	Cl 1	0	0
6	G	1	Total 1	Cl 1	0	0

- Molecule 7 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	1	Total 1	Mn 1	0	0
7	I	6	Total 6	Mn 6	0	0
7	J	5	Total 5	Mn 5	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	22	Total 22	O 22	0	0
8	B	19	Total 19	O 19	0	0
8	C	29	Total 29	O 29	0	0
8	D	13	Total 13	O 13	0	0
8	E	40	Total 40	O 40	0	0
8	F	28	Total 28	O 28	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	G	14	Total 14	O 14	0	0
8	H	15	Total 15	O 15	0	0
8	I	8	Total 8	O 8	0	0
8	J	10	Total 10	O 10	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Histone H3.1




• Molecule 1: Histone H3.1



• Molecule 2: Histone H4




• Molecule 2: Histone H4




- Molecule 3: Histone H2A type 1-B/E

Chain C:  53% 23% 6% 19%




- Molecule 3: Histone H2A type 1-B/E

Chain G:  59% 16% 22%




- Molecule 4: Histone H2B type 1-J

Chain D:  43% 23% 6% 27%





- Molecule 4: Histone H2B type 1-J


Chain H:  55% 12% 29%




- Molecule 5: 146-MER DNA

Chain I:  16% 14% 84%





● Molecule 5: 146-MER DNA



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	106.55Å 109.78Å 182.22Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.80 – 2.70 48.80 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.6 (48.80-2.70) 99.8 (48.80-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.99 (at 2.69Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.213 , 0.263 0.212 , 0.264	Depositor DCC
R_{free} test set	3000 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtriage
Anisotropy	0.326	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.014 for k,h,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	12152	wwPDB-VP
Average B, all atoms (Å ²)	70.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/813	0.94	2/1091 (0.2%)
1	E	0.58	0/828	0.93	3/1110 (0.3%)
2	B	0.51	0/626	0.97	2/837 (0.2%)
2	F	0.57	0/680	1.03	3/908 (0.3%)
3	C	0.53	0/845	0.95	1/1139 (0.1%)
3	G	0.51	0/815	0.95	1/1100 (0.1%)
4	D	0.53	0/747	0.96	4/1004 (0.4%)
4	H	0.53	0/725	0.91	1/975 (0.1%)
5	I	0.31	0/3332	0.77	2/5141 (0.0%)
5	J	0.30	0/3330	0.78	3/5138 (0.1%)
All	All	0.43	0/12741	0.86	22/18443 (0.1%)

There are no bond length outliers.

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	30	THR	N-CA-C	7.01	120.02	110.55
4	D	52	THR	N-CA-C	6.28	120.51	109.96
1	A	117	VAL	N-CA-C	-6.14	107.30	113.20
2	F	30	THR	N-CA-C	6.11	118.59	110.53
5	J	227	DG	C5'-C4'-C3'	-5.95	105.97	114.90

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	801	0	834	36	0
1	E	816	0	851	35	0
2	B	619	0	659	20	0
2	F	673	0	722	20	0
3	C	835	0	897	49	0
3	G	805	0	861	27	0
4	D	736	0	758	41	0
4	H	714	0	735	27	0
5	I	2970	0	1640	236	0
5	J	2969	0	1641	200	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
6	E	1	0	0	1	0
6	G	1	0	0	0	0
7	D	1	0	0	0	0
7	I	6	0	0	0	0
7	J	5	0	0	0	0
8	A	22	0	0	2	0
8	B	19	0	0	0	0
8	C	29	0	0	4	0
8	D	13	0	0	0	0
8	E	40	0	0	0	0
8	F	28	0	0	2	0
8	G	14	0	0	2	0
8	H	15	0	0	2	0
8	I	8	0	0	0	0
8	J	10	0	0	0	0
All	All	12152	0	9598	603	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 603 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:J:242:DT:H2''	5:J:243:DG:H5'	1.25	1.16
5:I:9:DC:H2''	5:I:10:DC:H5'	1.29	1.11
5:J:197:DA:H2''	5:J:198:DT:H5'	1.27	1.11
5:J:231:DA:H2''	5:J:232:DT:H5'	1.28	1.10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:J:190:DC:H2''	5:J:191:DT:H5''	1.32	1.09

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/139 (68%)	93 (98%)	2 (2%)	0	100	100
1	E	97/139 (70%)	94 (97%)	2 (2%)	1 (1%)	12	32
2	B	76/106 (72%)	74 (97%)	2 (3%)	0	100	100
2	F	82/106 (77%)	80 (98%)	2 (2%)	0	100	100
3	C	106/133 (80%)	101 (95%)	5 (5%)	0	100	100
3	G	102/133 (77%)	98 (96%)	3 (3%)	1 (1%)	12	32
4	D	92/129 (71%)	89 (97%)	2 (2%)	1 (1%)	11	29
4	H	89/129 (69%)	86 (97%)	2 (2%)	1 (1%)	11	29
All	All	739/1014 (73%)	715 (97%)	20 (3%)	4 (0%)	24	48

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	133	GLU
4	H	104	GLY
4	D	104	GLY
3	G	26	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/113 (75%)	81 (95%)	4 (5%)	23	51
1	E	86/113 (76%)	80 (93%)	6 (7%)	14	34
2	B	63/81 (78%)	61 (97%)	2 (3%)	34	64
2	F	69/81 (85%)	67 (97%)	2 (3%)	37	67
3	C	85/102 (83%)	77 (91%)	8 (9%)	8	21
3	G	83/102 (81%)	80 (96%)	3 (4%)	31	60
4	D	80/107 (75%)	72 (90%)	8 (10%)	7	19
4	H	78/107 (73%)	75 (96%)	3 (4%)	29	58
All	All	629/806 (78%)	593 (94%)	36 (6%)	18	43

5 of 36 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	19	ARG
4	H	106	LEU
2	F	47	SER
3	G	114	VAL
3	C	84	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	113	HIS
3	G	31	HIS
2	F	93	GLN
3	G	73	ASN
3	C	38	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 16 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	97/139 (69%)	-0.28	4 (4%) 41 37	26, 39, 64, 96	0
1	E	99/139 (71%)	-0.51	3 (3%) 52 49	16, 32, 68, 86	0
2	B	78/106 (73%)	-0.46	1 (1%) 75 73	27, 38, 60, 80	0
2	F	84/106 (79%)	-0.57	3 (3%) 46 42	21, 32, 56, 101	0
3	C	108/133 (81%)	-0.38	0 100 100	21, 38, 75, 125	0
3	G	104/133 (78%)	-0.34	1 (0%) 79 78	26, 43, 72, 98	0
4	D	94/129 (72%)	-0.26	1 (1%) 78 76	22, 40, 75, 105	0
4	H	91/129 (70%)	-0.26	1 (1%) 78 76	28, 42, 67, 87	0
5	I	145/146 (99%)	1.17	23 (15%) 5 4	43, 99, 141, 165	0
5	J	145/146 (99%)	1.22	26 (17%) 3 3	46, 101, 145, 160	0
All	All	1045/1306 (80%)	0.06	63 (6%) 27 24	16, 44, 129, 165	0

The worst 5 of 63 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	102	GLY	5.6
2	F	19	ARG	3.7
4	D	124	ALA	3.6
5	I	136	DT	3.5
4	H	123	SER	3.4

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	MN	I	1006	1/1	0.70	0.19	119,119,119,119	0
7	MN	I	1001	1/1	0.74	0.17	125,125,125,125	0
7	MN	I	1002	1/1	0.78	0.14	116,116,116,116	0
7	MN	I	1005	1/1	0.84	0.11	114,114,114,114	0
7	MN	J	1001	1/1	0.91	0.08	112,112,112,112	0
7	MN	J	1002	1/1	0.92	0.12	87,87,87,87	0
7	MN	I	1003	1/1	0.93	0.22	85,85,85,85	0
7	MN	I	1004	1/1	0.95	0.12	85,85,85,85	0
7	MN	J	1005	1/1	0.95	0.08	121,121,121,121	0
7	MN	J	1004	1/1	0.96	0.14	71,71,71,71	0
6	CL	C	1001	1/1	0.97	0.05	45,45,45,45	0
6	CL	G	1001	1/1	0.97	0.06	49,49,49,49	0
7	MN	J	1003	1/1	0.98	0.06	72,72,72,72	0
6	CL	E	1001	1/1	0.98	0.04	51,51,51,51	0
6	CL	A	1001	1/1	0.98	0.06	53,53,53,53	0
7	MN	D	201	1/1	0.99	0.08	36,36,36,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.