



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:48 PM UTC

PDB ID : 3AZR / pdb_00003azr
Title : Diverse Substrates Recognition Mechanism Revealed by Thermotoga maritima Cel5A Structures in Complex with Cellobiose
Authors : Wu, T.H.; Huang, C.H.; Ko, T.P.; Lai, H.L.; Ma, Y.; Chen, C.C.; Cheng, Y.S.; Liu, J.R.; Guo, R.T.
Deposited on : 2011-05-30
Resolution : 1.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1039 (1.72-1.72)
Clashscore	190562	1049 (1.72-1.72)
Ramachandran outliers	187476	1041 (1.72-1.72)
Sidechain outliers	187428	1041 (1.72-1.72)
RSRZ outliers	180081	1039 (1.72-1.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	317	3% 76% 20% ..
1	B	317	2% 71% 24% ..
2	C	2	50% 50%
2	D	2	50% 50%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5852 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Endoglucanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	311	Total	C	N	O	S	0	0	0
			2610	1699	445	461	5			
1	B	308	Total	C	N	O	S	0	0	0
			2591	1687	442	458	4			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	253	ALA	GLU	engineered mutation	UNP Q9X273
B	253	ALA	GLU	engineered mutation	UNP Q9X273

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is beta-D-glucopyranose (CCD ID: BGC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			12	6	6		
3	B	1	Total	C	O	0	0
			12	6	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	296	Total	O	0	0
			296	296		
4	B	285	Total	O	0	0
			285	285		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endoglucanase



- Molecule 1: Endoglucanase



- Molecule 2: beta-D-glucopyranose-(1-4)-alpha-D-glucopyranose



- Molecule 2: beta-D-glucopyranose-(1-4)-alpha-D-glucopyranose





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.42Å 77.14Å 62.62Å 90.00° 97.50° 90.00°	Depositor
Resolution (Å)	24.98 – 1.71 24.98 – 1.71	Depositor EDS
% Data completeness (in resolution range)	(Not available) (24.98-1.71) 95.8 (24.98-1.71)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.15 (at 1.72Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.201 , 0.246 0.198 , 0.240	Depositor DCC
R_{free} test set	3062 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.441	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.027 for l,-k,h	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5852	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.42% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.10	11/2691 (0.4%)	1.13	18/3640 (0.5%)
1	B	0.95	6/2672 (0.2%)	1.10	16/3615 (0.4%)
All	All	1.03	17/5363 (0.3%)	1.11	34/7255 (0.5%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	137	PRO	C-O	-15.39	1.04	1.23
1	A	138	HIS	C-O	-15.34	1.06	1.23
1	B	138	HIS	C-O	-13.11	1.09	1.23
1	A	94	ILE	C-O	-12.45	1.10	1.24
1	A	95	HIS	C-O	-12.10	1.09	1.23
1	B	137	PRO	C-O	-10.53	1.10	1.23
1	A	204	THR	C-O	-9.26	1.12	1.24
1	B	206	GLN	C-O	-8.18	1.13	1.23
1	A	276	MET	SD-CE	6.82	1.96	1.79
1	A	52	ILE	CA-CB	6.43	1.60	1.54
1	B	206	GLN	CD-NE2	-6.26	1.20	1.33
1	B	95	HIS	C-O	-5.95	1.16	1.23
1	B	205	HIS	C-O	-5.90	1.15	1.24
1	A	205	HIS	C-O	-5.87	1.16	1.24
1	A	95	HIS	CA-C	5.56	1.59	1.52
1	A	204	THR	CB-OG1	-5.33	1.35	1.43
1	A	34	ILE	CA-CB	5.32	1.60	1.53

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	288	PHE	N-CA-C	9.07	120.85	110.97
1	B	94	ILE	N-CA-C	-8.36	96.47	108.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	133	ILE	N-CA-C	7.89	118.65	110.36
1	B	304	LYS	N-CA-C	7.32	119.26	111.28
1	B	133	ILE	N-CA-C	6.85	117.55	110.36
1	A	204	THR	N-CA-C	6.74	119.46	111.71
1	B	288	PHE	N-CA-C	6.66	118.42	111.03
1	B	89	ALA	N-CA-C	-6.30	102.41	110.53
1	A	304	LYS	N-CA-C	6.29	118.14	111.28
1	A	52	ILE	CA-C-N	6.20	126.40	119.90
1	A	52	ILE	C-N-CA	6.20	126.40	119.90
1	A	177	SER	N-CA-C	6.17	118.81	111.71
1	B	294	VAL	N-CA-C	-6.16	106.72	113.43
1	A	70	ASP	N-CA-C	6.09	118.41	111.11
1	A	52	ILE	N-CA-C	5.92	114.01	108.15
1	A	303	ASN	N-CA-C	-5.85	98.18	108.23
1	B	289	CYS	N-CA-C	5.54	119.84	112.30
1	A	98	GLU	N-CA-C	5.53	117.31	111.28
1	B	303	ASN	N-CA-C	-5.51	98.75	108.23
1	A	31	GLY	N-CA-C	5.47	122.03	114.92
1	A	138	HIS	N-CA-C	5.43	115.71	108.38
1	B	160	ILE	CB-CA-C	-5.38	105.64	110.91
1	B	159	SER	N-CA-C	-5.36	106.70	113.18
1	A	282	SER	N-CA-C	-5.36	102.82	110.59
1	B	98	GLU	N-CA-C	5.34	117.52	111.11
1	B	136	GLU	N-CA-C	5.33	121.59	109.81
1	A	89	ALA	N-CA-C	-5.27	103.73	110.53
1	B	52	ILE	CA-C-N	5.24	125.40	119.90
1	B	52	ILE	C-N-CA	5.24	125.40	119.90
1	B	282	SER	N-CA-C	-5.22	103.03	110.59
1	B	32	VAL	N-CA-C	5.20	115.34	107.75
1	A	121	ARG	N-CA-C	5.18	116.93	111.28
1	A	292	PHE	N-CA-C	-5.08	106.32	112.88
1	A	252	GLY	N-CA-C	5.05	120.01	114.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2610	0	2551	44	0
1	B	2591	0	2527	55	0
2	C	23	0	21	1	0
2	D	23	0	21	1	0
3	A	12	0	12	1	0
3	B	12	0	12	2	0
4	A	296	0	0	7	0
4	B	285	0	0	10	0
All	All	5852	0	5144	101	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (101) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:VAL:HG12	1:A:306:LEU:HD23	1.46	0.95
1:A:225:PRO:O	1:A:229:LYS:HG3	1.69	0.93
1:B:65:PRO:HG2	1:B:67:LYS:HG3	1.54	0.89
1:A:8:ARG:HD2	1:A:165:THR:OG1	1.75	0.86
1:B:93:ASN:C	1:B:93:ASN:HD22	1.93	0.77
1:A:1:MET:HE2	4:A:567:HOH:O	1.85	0.75
1:A:244:LYS:HB2	1:A:244:LYS:NZ	2.01	0.74
1:A:303:ASN:HB2	4:A:533:HOH:O	1.87	0.74
1:B:158:ARG:O	1:B:162:LYS:HE3	1.88	0.73
1:A:294:VAL:HG12	1:A:306:LEU:CD2	2.20	0.72
1:A:258:ARG:HH11	1:A:258:ARG:HG3	1.55	0.70
1:A:1:MET:HG2	1:A:2:GLY:N	2.06	0.68
1:A:231:LEU:HD11	1:A:272:VAL:HG22	1.76	0.67
1:A:8:ARG:CD	1:A:165:THR:OG1	2.44	0.65
1:B:202:GLU:H	1:B:202:GLU:CD	2.03	0.65
1:A:244:LYS:HB2	1:A:244:LYS:HZ2	1.62	0.64
3:B:402:BGC:H6C2	4:B:382:HOH:O	1.97	0.63
1:A:308:GLU:O	1:A:311:ILE:HG13	1.98	0.62
1:B:229:LYS:HE3	4:B:460:HOH:O	1.98	0.62
1:B:186:LYS:HD2	1:B:186:LYS:N	2.14	0.60
1:A:206:GLN:NE2	1:A:219:GLY:H	1.99	0.60
1:B:163:LYS:NZ	4:B:586:HOH:O	2.33	0.60
1:B:280:ARG:HG2	1:B:280:ARG:NH1	2.17	0.60
1:A:8:ARG:NH1	1:A:127:GLU:HB3	2.17	0.59
1:B:8:ARG:HB3	1:B:130:PHE:HE1	1.67	0.59
1:A:271:PHE:HD1	1:A:274:ARG:NH2	2.01	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:216:LYS:NZ	4:B:400:HOH:O	2.36	0.59
1:B:65:PRO:CG	1:B:67:LYS:HG3	2.31	0.59
1:B:280:ARG:HG2	1:B:280:ARG:HH11	1.69	0.58
1:B:299:ARG:O	1:B:301:THR:HG23	2.04	0.58
1:A:8:ARG:HD2	1:A:165:THR:CB	2.34	0.58
1:A:272:VAL:HG12	1:A:276:MET:HE2	1.86	0.57
1:B:8:ARG:HG3	1:B:165:THR:OG1	2.06	0.56
1:B:61:TYR:HB3	1:B:63:PHE:CE1	2.41	0.56
1:A:233:GLU:HB2	4:A:393:HOH:O	2.05	0.56
1:B:93:ASN:C	1:B:93:ASN:ND2	2.60	0.55
1:B:14:ARG:NH1	1:B:311:ILE:HG22	2.21	0.55
1:B:4:ASP:N	4:B:552:HOH:O	2.39	0.55
1:B:8:ARG:HB3	1:B:130:PHE:CE1	2.42	0.54
1:B:272:VAL:HG12	1:B:276:MET:HE2	1.87	0.54
1:B:148:GLU:HG2	4:B:388:HOH:O	2.07	0.54
1:B:274:ARG:HD3	4:B:484:HOH:O	2.08	0.54
1:A:258:ARG:HG3	1:A:258:ARG:NH1	2.19	0.54
1:B:202:GLU:CD	1:B:202:GLU:N	2.66	0.54
1:B:15:GLY:HA3	1:B:49:HIS:CE1	2.43	0.53
1:B:55:ARG:O	1:B:58:THR:HG22	2.08	0.53
1:B:54:ILE:H	1:B:93:ASN:HD21	1.57	0.52
1:B:114:LEU:O	1:B:118:ILE:HG13	2.10	0.52
1:B:58:THR:HG23	1:B:59:HIS:ND1	2.25	0.52
1:B:58:THR:HG23	1:B:59:HIS:CE1	2.44	0.51
1:B:194:THR:HA	1:B:250:TYR:O	2.10	0.51
1:A:59:HIS:HE1	4:A:332:HOH:O	1.93	0.51
1:A:8:ARG:HH12	1:A:127:GLU:HB3	1.75	0.50
1:A:194:THR:HA	1:A:250:TYR:O	2.12	0.50
1:A:243:LYS:HG3	4:A:493:HOH:O	2.12	0.50
1:B:308:GLU:O	1:B:311:ILE:HG13	2.11	0.49
1:A:8:ARG:NE	1:A:165:THR:OG1	2.46	0.49
1:A:202:GLU:HG3	4:A:461:HOH:O	2.13	0.49
1:A:271:PHE:HD1	1:A:274:ARG:HH22	1.60	0.49
1:A:59:HIS:HD2	4:B:330:HOH:O	1.96	0.49
1:B:195:ILE:O	1:B:251:ILE:HA	2.13	0.49
1:B:65:PRO:HG3	1:B:67:LYS:HE2	1.95	0.48
1:B:124:ASP:O	4:B:509:HOH:O	2.20	0.47
1:A:136:GLU:OE2	2:C:1:GLC:H1	2.15	0.47
1:A:3:VAL:HG12	1:A:8:ARG:CZ	2.45	0.46
1:A:5:PRO:HB2	1:A:248:PRO:HG2	1.98	0.46
1:A:195:ILE:O	1:A:251:ILE:HA	2.15	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:136:GLU:O	1:B:138:HIS:HD2	1.98	0.46
3:B:402:BGC:C6	4:B:382:HOH:O	2.58	0.46
1:A:172:GLU:O	1:A:173:TRP:HB2	2.16	0.45
1:B:26:ASN:ND2	1:B:29:ASP:OD2	2.48	0.45
1:B:266:ILE:HD13	1:B:305:ASP:HB3	1.98	0.45
1:A:263:GLU:O	1:A:263:GLU:HG3	2.16	0.45
1:B:274:ARG:O	1:B:278:LYS:HG3	2.16	0.45
1:A:249:ILE:HB	1:A:281:TRP:CD1	2.51	0.45
1:B:54:ILE:H	1:B:93:ASN:ND2	2.14	0.45
1:B:136:GLU:OE2	2:D:1:GLC:H1	2.17	0.45
1:A:178:ALA:HA	1:A:181:LYS:HE3	1.99	0.44
1:B:116:LYS:HG3	1:B:160:ILE:HD11	1.99	0.44
1:B:287:GLU:OE2	1:B:290:SER:HB3	2.18	0.44
1:B:306:LEU:HD23	1:B:306:LEU:HA	1.80	0.44
1:B:161:ASP:OD1	1:B:161:ASP:C	2.60	0.44
1:B:93:ASN:HB3	1:B:132:GLU:HB3	2.00	0.43
1:B:150:LEU:HD11	1:B:185:PRO:HD3	2.00	0.43
1:A:243:LYS:CG	4:A:493:HOH:O	2.65	0.43
1:B:63:PHE:CD2	1:B:64:PRO:O	2.72	0.43
1:B:280:ARG:HH11	1:B:280:ARG:CG	2.30	0.43
1:A:201:PHE:CZ	3:A:401:BGC:H5	2.53	0.42
1:A:235:PHE:HZ	1:A:272:VAL:HG13	1.84	0.42
1:B:285:TYR:CE1	1:B:294:VAL:HG13	2.54	0.42
1:A:93:ASN:HB3	1:A:132:GLU:HB3	2.01	0.42
1:B:22:LEU:HD23	1:B:27:GLU:HB2	2.02	0.42
1:A:244:LYS:NZ	1:A:244:LYS:CB	2.77	0.41
1:B:273:VAL:O	1:B:277:GLU:HG3	2.20	0.41
1:A:142:THR:O	1:A:143:PRO:C	2.62	0.41
1:B:85:LYS:HG2	1:B:86:ARG:HD2	2.03	0.41
1:B:186:LYS:N	1:B:186:LYS:CD	2.82	0.41
1:A:58:THR:HG23	1:A:59:HIS:CE1	2.55	0.41
1:B:286:TRP:CD1	1:B:287:GLU:HG3	2.56	0.41
1:A:93:ASN:OD1	1:A:93:ASN:C	2.64	0.40
1:B:272:VAL:O	1:B:276:MET:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/317 (98%)	300 (97%)	9 (3%)	0	100	100
1	B	306/317 (96%)	297 (97%)	9 (3%)	0	100	100
All	All	615/634 (97%)	597 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	272/276 (99%)	263 (97%)	9 (3%)	33	10
1	B	270/276 (98%)	265 (98%)	5 (2%)	50	27
All	All	542/552 (98%)	528 (97%)	14 (3%)	40	18

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	8	ARG
1	A	73	PHE
1	A	143	PRO
1	A	144	GLU
1	A	220	ARG
1	A	244	LYS
1	A	263	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	287	GLU
1	B	8	ARG
1	B	44	GLU
1	B	93	ASN
1	B	244	LYS
1	B	287	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	59	HIS
1	A	96	HIS
1	A	140	ASN
1	A	190	ASN
1	A	206	GLN
1	B	81	ASN
1	B	93	ASN
1	B	96	HIS
1	B	138	HIS
1	B	140	ASN
1	B	190	ASN
1	B	206	GLN
1	B	230	HIS
1	B	245	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GLC	C	1	2	12,12,12	0.50	0	17,17,17	0.75	0
2	BGC	C	2	2	11,11,12	0.38	0	15,15,17	0.87	0
2	GLC	D	1	2	12,12,12	0.49	0	17,17,17	0.84	1 (5%)
2	BGC	D	2	2	11,11,12	0.34	0	15,15,17	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	BGC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	BGC	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	D	1	GLC	C3-C4-C5	-2.08	106.46	110.23

There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1	GLC	1	0
2	D	1	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BGC	A	401	-	12,12,12	0.49	0	17,17,17	0.66	0
3	BGC	B	402	-	12,12,12	0.48	0	17,17,17	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BGC	A	401	-	-	0/2/22/22	0/1/1/1
3	BGC	B	402	-	-	0/2/22/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	401	BGC	1	0
3	B	402	BGC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	311/317 (98%)	0.13	9 (2%) 53 62	7, 19, 32, 40	0
1	B	308/317 (97%)	0.29	6 (1%) 66 73	11, 21, 33, 42	0
All	All	619/634 (97%)	0.21	15 (2%) 59 67	7, 20, 33, 42	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	63	PHE	5.2
1	B	311	ILE	3.5
1	A	8	ARG	3.2
1	A	240	GLU	3.1
1	B	8	ARG	2.5
1	A	3	VAL	2.5
1	B	61	TYR	2.5
1	A	274	ARG	2.4
1	A	217	TRP	2.3
1	A	311	ILE	2.3
1	B	67	LYS	2.2
1	B	301	THR	2.2
1	A	230	HIS	2.1
1	A	186	LYS	2.1
1	A	215	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BGC	D	2	11/12	0.87	0.10	18,20,24,27	0
2	GLC	C	1	12/12	0.92	0.07	17,19,21,21	0
2	BGC	C	2	11/12	0.94	0.07	17,19,22,24	0
2	GLC	D	1	12/12	0.96	0.05	17,19,21,22	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BGC	A	401	12/12	0.74	0.16	64,65,65,65	0
3	BGC	B	402	12/12	0.80	0.16	48,49,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.