



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 02:53 PM UTC

PDB ID : 5AZ9 / pdb_00005az9
Title : Crystal structure of (5-residue deleted)MBP-Tom20 fusion protein tethered with ALDH presequence via a disulfide bond
Authors : Matsuoka, R.; Kohda, D.
Deposited on : 2015-09-27
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	430	9% 84% 12% ...

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3498 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltose-binding periplasmic protein,Mitochondrial import receptor subunit TOM20 homolog.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	425	Total	C	N	O	S	0	0	0
			3326	2151	539	628	8			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P0AEX9
A	?	-	ALA	deletion	UNP P0AEX9
A	?	-	ALA	deletion	UNP P0AEX9
A	?	-	THR	deletion	UNP P0AEX9
A	?	-	GLY	deletion	UNP P0AEX9
A	?	-	ASP	deletion	UNP P0AEX9
A	308	VAL	ALA	engineered mutation	UNP P0AEX9
A	365	LYS	-	linker	UNP P0AEX9
A	366	GLU	-	linker	UNP P0AEX9
A	367	ALA	-	linker	UNP P0AEX9
A	368	LEU	-	linker	UNP P0AEX9


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	172	Total	O	0	0
			172	172		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Maltose-binding periplasmic protein,Mitochondrial import receptor subunit TOM20 homolog



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	160.48Å 71.40Å 47.54Å 90.00° 104.91° 90.00°	Depositor
Resolution (Å)	39.55 – 1.82 39.55 – 1.82	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.55-1.82) 99.8 (39.55-1.82)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.192 , 0.227 0.202 , 0.232	Depositor DCC
R_{free} test set	2351 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	28.8	Xtriage
Anisotropy	0.057	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.020 for -h-2*1,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3498	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	1/3403 (0.0%)	1.13	7/4618 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	404	CYS	CA-CB	7.72	1.64	1.53

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	394	VAL	N-CA-C	-6.37	106.33	111.81
1	A	363	ARG	CG-CD-NE	6.01	125.23	112.00
1	A	358	LYS	N-CA-C	5.47	117.67	111.11
1	A	345	ASN	N-CA-C	5.37	117.14	111.28
1	A	52	GLY	CA-C-N	5.32	126.67	120.98
1	A	52	GLY	C-N-CA	5.32	126.67	120.98
1	A	392	LYS	N-CA-C	5.10	125.28	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	391	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3326	0	3330	35	1
2	A	172	0	0	2	0
All	All	3498	0	3330	35	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:363:ARG:NH1	1:A:366:GLU:OE2	1.99	0.94
1:A:352:THR:HG23	1:A:355:GLU:H	1.55	0.71
1:A:394:VAL:O	1:A:398:THR:HG23	1.92	0.70
1:A:422:VAL:HG22	1:A:425:MET:CE	2.24	0.68
1:A:422:VAL:HG22	1:A:425:MET:HE2	1.84	0.60
1:A:417:THR:HG22	1:A:418:LEU:HD13	1.86	0.58
1:A:390:TYR:CE1	1:A:391:GLU:HB3	2.40	0.56
1:A:11:TRP:CG	1:A:53:PRO:HG3	2.42	0.55
1:A:40:HIS:HD2	2:A:650:HOH:O	1.90	0.53
1:A:415:GLN:HG2	1:A:416:GLN:HE22	1.72	0.53
1:A:191:LEU:CD1	1:A:200:MET:HE1	2.39	0.52
1:A:415:GLN:HG2	1:A:416:GLN:NE2	2.25	0.52
1:A:64:GLY:HA3	1:A:328:ASN:O	2.12	0.50
1:A:392:LYS:HD2	1:A:392:LYS:N	2.29	0.48
1:A:168:GLU:HG2	1:A:169:ASN:ND2	2.29	0.47
1:A:398:THR:HG22	1:A:426:LEU:HD12	1.96	0.47
1:A:412:GLN:O	1:A:416:GLN:NE2	2.48	0.46
1:A:352:THR:CG2	1:A:355:GLU:HB2	2.45	0.46
1:A:370:VAL:CG1	1:A:403:VAL:HG13	2.47	0.45
1:A:390:TYR:OH	1:A:422:VAL:HG21	2.16	0.45
1:A:345:ASN:HD22	1:A:351:GLN:NE2	2.14	0.45
1:A:35:LYS:HG3	2:A:551:HOH:O	2.16	0.45
1:A:398:THR:HG21	1:A:429:LYS:HB2	1.99	0.44
1:A:342:ALA:HB2	1:A:360:ALA:HB2	2.00	0.43
1:A:423:PHE:O	1:A:426:LEU:HB3	2.18	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:THR:HG22	1:A:355:GLU:HB2	2.00	0.42
1:A:391:GLU:CD	1:A:392:LYS:N	2.77	0.42
1:A:381:GLY:O	1:A:384:LEU:HG	2.20	0.42
1:A:77:PRO:HA	1:A:277:GLU:OE2	2.20	0.42
1:A:105:ALA:O	1:A:257:VAL:HA	2.20	0.42
1:A:420:PRO:N	1:A:421:PRO:CD	2.83	0.41
1:A:48:PHE:HB3	1:A:49:PRO:HD3	2.02	0.41
1:A:416:GLN:N	1:A:416:GLN:CD	2.79	0.41
1:A:419:PRO:HG2	1:A:422:VAL:HB	2.02	0.40
1:A:363:ARG:CZ	1:A:366:GLU:OE2	2.66	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:363:ARG:NH1	1:A:363:ARG:NH1[2_755]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	421/430 (98%)	407 (97%)	9 (2%)	5 (1%)	10 3

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	391	GLU
1	A	392	LYS
1	A	418	LEU
1	A	416	GLN
1	A	419	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	349/352 (99%)	324 (93%)	25 (7%)	13 2

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	GLN
1	A	75	ILE
1	A	79	LYS
1	A	147	LEU
1	A	171	LYS
1	A	198	LYS
1	A	291	LYS
1	A	293	LYS
1	A	326	MET
1	A	358	LYS
1	A	372	LYS
1	A	384	LEU
1	A	391	GLU
1	A	403	VAL
1	A	404	CYS
1	A	409	GLN
1	A	411	LEU
1	A	413	VAL
1	A	414	LEU
1	A	415	GLN
1	A	416	GLN
1	A	417	THR
1	A	418	LEU
1	A	424	GLN
1	A	428	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	GLN
1	A	169	ASN
1	A	345	ASN
1	A	399	ASN
1	A	406	GLN
1	A	416	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	425/430 (98%)	0.37	40 (9%) 14 15	20, 31, 68, 99	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	418	LEU	6.6
1	A	393	GLY	5.9
1	A	385	LEU	5.5
1	A	417	THR	5.4
1	A	390	TYR	4.7
1	A	384	LEU	4.2
1	A	337	TYR	4.2
1	A	421	PRO	4.2
1	A	419	PRO	3.9
1	A	392	LYS	3.8
1	A	380	LEU	3.7
1	A	394	VAL	3.5
1	A	404	CYS	3.4
1	A	422	VAL	3.3
1	A	373	PHE	3.1
1	A	396	HIS	3.1
1	A	344	ILE	3.0
1	A	375	LEU	2.9
1	A	413	VAL	2.9
1	A	51	VAL	2.8
1	A	420	PRO	2.8
1	A	423	PHE	2.7
1	A	430	LEU	2.6
1	A	378	ILE	2.6
1	A	426	LEU	2.5
1	A	2	LYS	2.4
1	A	3	ILE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	352	THR	2.3
1	A	395	ASP	2.2
1	A	397	LEU	2.2
1	A	198	LYS	2.2
1	A	415	GLN	2.2
1	A	429	LYS	2.2
1	A	424	GLN	2.2
1	A	412	GLN	2.1
1	A	349	GLY	2.1
1	A	382	GLU	2.1
1	A	425	MET	2.0
1	A	408	GLN	2.0
1	A	345	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.