



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:49 AM UTC

PDB ID : 7AZF / pdb_00007azf
Title : DNA polymerase sliding clamp from Escherichia coli with peptide 8 bound
Authors : Monsarrat, C.; Compain, G.; Andre, C.; Martiel, I.; Engilberge, S.; Olieric, V.;
Wolff, P.; Brillet, K.; Landolfo, M.; Silva da Veiga, C.; Wagner, J.; Guichard,
G.; Burnouf, D.Y.
Deposited on : 2020-11-16
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	369	4% 92% 7%
1	B	369	4% 94% 6%
1	C	369	13% 90% 10%
1	D	369	18% 92% 7%
2	H	6	67% 33%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	I	6	 83% 17%
2	J	6	 83% 17%
2	K	6	 17% 83% 17%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 12840 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta sliding clamp.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	369	Total	C	N	O	S	0	8	0
			2905	1825	509	550	21			
1	B	369	Total	C	N	O	S	0	7	0
			2900	1821	515	545	19			
1	C	368	Total	C	N	O	S	0	3	0
			2879	1807	506	547	19			
1	D	368	Total	C	N	O	S	0	1	0
			2828	1776	496	537	19			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP A0A073FMV0
A	-1	SER	-	expression tag	UNP A0A073FMV0
A	0	HIS	-	expression tag	UNP A0A073FMV0
B	-2	GLY	-	expression tag	UNP A0A073FMV0
B	-1	SER	-	expression tag	UNP A0A073FMV0
B	0	HIS	-	expression tag	UNP A0A073FMV0
C	-2	GLY	-	expression tag	UNP A0A073FMV0
C	-1	SER	-	expression tag	UNP A0A073FMV0
C	0	HIS	-	expression tag	UNP A0A073FMV0
D	-2	GLY	-	expression tag	UNP A0A073FMV0
D	-1	SER	-	expression tag	UNP A0A073FMV0
D	0	HIS	-	expression tag	UNP A0A073FMV0

- Molecule 2 is a protein called Peptide 8.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	H	6	Total	C	N	O	0	0	0
			59	41	8	10			
2	I	6	Total	C	N	O	0	1	0
			70	49	10	11			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	J	6	Total	C	N	O	0	0	0
			51	35	6	10			
2	K	6	Total	C	N	O	0	0	0
			51	35	6	10			

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	312	Total	O	0	7
			315	315		
5	B	321	Total	O	0	10
			329	329		
5	C	193	Total	O	0	10
			203	203		
5	D	201	Total	O	0	3
			203	203		
5	H	7	Total	O	0	0
			7	7		
5	I	9	Total	O	0	0
			9	9		
5	J	9	Total	O	0	0
			9	9		
5	K	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 1: Beta sliding clamp




- Molecule 2: Peptide 8




- Molecule 2: Peptide 8



- Molecule 2: Peptide 8



- Molecule 2: Peptide 8



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	70.42Å 82.03Å 82.39Å 116.87° 100.26° 95.47°	Depositor
Resolution (Å)	37.75 – 1.93 37.75 – 1.93	Depositor EDS
% Data completeness (in resolution range)	70.8 (37.75-1.93) 71.2 (37.75-1.93)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.92Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.204 , 0.251 0.206 , 0.253	Depositor DCC
R_{free} test set	4137 reflections (3.44%)	wwPDB-VP
Wilson B-factor (Å ²)	34.4	Xtriage
Anisotropy	0.007	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 66.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.000 for -h,-l,-k	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	12840	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PEG, SGK, GOL, ALC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.91	2/2973 (0.1%)	0.93	2/4020 (0.0%)
1	B	0.88	2/2969 (0.1%)	0.90	0/4017
1	C	0.79	1/2938 (0.0%)	0.91	0/3975
1	D	0.81	1/2880 (0.0%)	0.94	4/3902 (0.1%)
2	H	0.70	0/36	0.76	0/44
2	I	0.65	0/36	0.76	0/44
2	J	0.54	0/36	0.87	0/44
2	K	0.63	0/36	0.81	0/44
All	All	0.85	6/11904 (0.1%)	0.92	6/16090 (0.0%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	146	MET	SD-CE	-10.61	1.53	1.79
1	C	146	MET	SD-CE	-7.10	1.61	1.79
1	A	53	MET	SD-CE	-6.78	1.62	1.79
1	B	146	MET	SD-CE	-6.71	1.62	1.79
1	B	315	MET	SD-CE	-5.74	1.65	1.79
1	D	146	MET	SD-CE	-5.05	1.67	1.79

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	345[A]	SER	CA-C-N	5.73	128.98	120.90
1	D	345[A]	SER	C-N-CA	5.73	128.98	120.90
1	D	345[B]	SER	CA-C-N	5.73	128.98	120.90
1	D	345[B]	SER	C-N-CA	5.73	128.98	120.90
1	A	45	THR	CA-C-N	5.13	125.72	122.18
1	A	45	THR	C-N-CA	5.13	125.72	122.18

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2905	0	2929	22	0
1	B	2900	0	2918	15	0
1	C	2879	0	2898	20	0
1	D	2828	0	2824	13	0
2	H	59	0	45	1	0
2	I	70	0	45	0	0
2	J	51	0	46	0	0
2	K	51	0	45	1	0
3	A	6	0	8	0	0
4	A	7	0	10	0	0
5	A	315	0	0	3	0
5	B	329	0	0	1	0
5	C	203	0	0	0	0
5	D	203	0	0	2	0
5	H	7	0	0	0	0
5	I	9	0	0	0	0
5	J	9	0	0	0	0
5	K	9	0	0	0	0
All	All	12840	0	11768	66	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (66) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:214:LEU:HD12	1:B:227:VAL:CG2	2.11	0.79
1:B:184:ILE:HD11	1:B:188:LEU:HD11	1.67	0.76
1:B:184:ILE:HD11	1:B:188:LEU:CD1	2.16	0.75
1:B:214:LEU:HD12	1:B:227:VAL:HG22	1.73	0.71
1:D:177:LEU:HD21	1:D:179:VAL:HG23	1.81	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:51:MET:HE1	1:A:198:LYS:HB3	1.83	0.61
1:A:151:VAL:HG13	1:D:175:HIS:CE1	2.38	0.58
1:C:184:ILE:HD11	1:C:188:LEU:HD11	1.84	0.58
1:C:16:GLN:HG3	1:C:230:PHE:CZ	2.39	0.58
1:A:144:PHE:CD2	1:A:326:ASP:HB3	2.39	0.58
1:A:17:VAL:HG12	1:A:33:LEU:HD22	1.87	0.56
1:D:159:LEU:HB3	1:D:170:VAL:CG1	2.35	0.56
1:B:206:MET:HE3	1:B:230:PHE:CD2	2.40	0.56
1:D:258:ALA:HB2	1:D:308:VAL:HG22	1.88	0.56
1:C:132:GLN:HB2	1:C:207:LEU:HD22	1.87	0.56
1:C:150:ASP:OD1	1:C:152:ARG:HD3	2.05	0.55
1:B:2:LYS:NZ	1:B:91:GLN:OE1	2.39	0.54
1:A:315:MET:HE1	1:A:340:LEU:HD22	1.89	0.54
1:A:151:VAL:HG13	1:D:175:HIS:NE2	2.21	0.53
1:A:51:MET:HE1	1:A:198:LYS:CB	2.38	0.53
1:C:134:THR:HG23	1:C:182:MET:HG2	1.90	0.52
1:C:139:ILE:HG21	1:C:204:MET:HG2	1.91	0.52
1:C:82:LEU:HD13	1:C:88:ILE:CD1	2.41	0.51
1:A:365:ARG:HD2	2:H:403:SGK:C7	2.41	0.51
1:D:159:LEU:HB3	1:D:170:VAL:HG13	1.92	0.50
1:B:144:PHE:CD2	1:B:326:ASP:HB3	2.47	0.49
1:B:184:ILE:CD1	1:B:188:LEU:HD11	2.41	0.48
1:C:144:PHE:CD2	1:C:326:ASP:HB3	2.48	0.48
1:A:2:LYS:NZ	5:A:502:HOH:O	2.47	0.48
1:A:19:GLY:HA2	5:A:515:HOH:O	2.13	0.47
1:A:13:PRO:HA	1:A:230:PHE:HE1	1.80	0.46
1:D:356:SER:HB3	5:D:599:HOH:O	2.14	0.46
1:B:338:MET:HG2	1:B:349:ILE:HG12	1.97	0.46
1:C:139:ILE:HG21	1:C:204:MET:CG	2.45	0.46
1:A:338:MET:HG2	1:A:349:ILE:HG12	1.98	0.46
1:A:175:HIS:CE1	1:D:151:VAL:HB	2.52	0.45
1:C:284:TYR:HB2	1:C:291:LYS:HB3	1.98	0.45
1:B:151:VAL:HB	1:C:175:HIS:CE1	2.51	0.45
1:C:126:VAL:HG22	1:C:189:PRO:HG2	1.99	0.45
1:D:144:PHE:CD2	1:D:326:ASP:HB3	2.52	0.45
1:D:365:ARG:HE	2:K:403:ALC:HZ3	1.82	0.45
1:C:11:LEU:HD11	1:C:80:ARG:HA	1.98	0.44
1:A:33:LEU:O	1:A:69:THR:HA	2.17	0.44
1:D:184:ILE:HD11	1:D:188:LEU:HD11	2.00	0.44
1:A:243:ASP:OD1	1:A:245:ARG:HG2	2.18	0.44
1:B:21:LEU:HD21	5:B:526:HOH:O	2.16	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:16:GLN:HB3	1:A:53:MET:HE2	1.99	0.43
1:C:33:LEU:O	1:C:69:THR:HA	2.18	0.43
1:C:282:ARG:HD2	1:C:284:TYR:OH	2.19	0.42
1:B:130:LEU:HD21	1:B:214:LEU:HD23	2.01	0.42
1:A:19:GLY:C	1:A:21:LEU:H	2.27	0.42
1:D:263:LEU:HD21	1:D:338:MET:CE	2.50	0.42
1:C:12:LYS:HB3	1:C:13:PRO:HD3	2.01	0.42
1:C:132:GLN:CB	1:C:207:LEU:HD22	2.50	0.42
1:A:17:VAL:CG1	1:A:33:LEU:HD22	2.50	0.42
1:A:165:GLU:HG3	5:A:748:HOH:O	2.20	0.41
1:B:206:MET:HE3	1:B:230:PHE:HD2	1.84	0.41
1:C:2:LYS:HG2	1:C:64:GLU:HG2	2.03	0.41
1:B:197[A]:ARG:HH22	1:B:198:LYS:HE3	1.84	0.41
1:C:13:PRO:HA	1:C:230:PHE:HE1	1.86	0.41
1:D:20:PRO:HG2	5:D:672:HOH:O	2.20	0.41
1:A:53:MET:CE	1:A:230:PHE:HB3	2.51	0.41
1:A:146:MET:HE3	1:A:157:GLY:C	2.46	0.40
1:A:203:LEU:O	1:A:206[B]:MET:HG2	2.20	0.40
1:C:244:TYR:O	1:C:248:LEU:HG	2.22	0.40
1:B:20:PRO:HG2	1:B:53:MET:SD	2.62	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/369 (102%)	364 (97%)	11 (3%)	0	100	100
1	B	374/369 (101%)	366 (98%)	8 (2%)	0	100	100
1	C	369/369 (100%)	361 (98%)	8 (2%)	0	100	100
1	D	367/369 (100%)	357 (97%)	9 (2%)	1 (0%)	36	30

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	H	3/6 (50%)	3 (100%)	0	0	100	100
2	I	3/6 (50%)	3 (100%)	0	0	100	100
2	J	3/6 (50%)	3 (100%)	0	0	100	100
2	K	3/6 (50%)	3 (100%)	0	0	100	100
All	All	1497/1500 (100%)	1460 (98%)	36 (2%)	1 (0%)	48	41

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	252	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	321/315 (102%)	317 (99%)	4 (1%)	63	57
1	B	318/315 (101%)	316 (99%)	2 (1%)	78	78
1	C	318/315 (101%)	315 (99%)	3 (1%)	70	67
1	D	308/315 (98%)	299 (97%)	9 (3%)	37	24
2	H	4/4 (100%)	4 (100%)	0	100	100
2	I	4/4 (100%)	4 (100%)	0	100	100
2	J	4/4 (100%)	4 (100%)	0	100	100
2	K	4/4 (100%)	4 (100%)	0	100	100
All	All	1281/1276 (100%)	1263 (99%)	18 (1%)	59	52

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	THR
1	A	118	ASN
1	A	119	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	151	VAL
1	B	12	LYS
1	B	118	ASN
1	C	124	SER
1	C	138	LEU
1	C	305	ILE
1	D	11	LEU
1	D	12	LYS
1	D	120	ASP
1	D	121	ASP
1	D	188	LEU
1	D	286	SER
1	D	308	VAL
1	D	332	LYS
1	D	346	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	GLN
1	A	118	ASN
1	A	186	GLN
1	B	15	GLN
1	B	118	ASN
1	B	265	GLN
1	C	148	HIS
1	C	186	GLN
1	C	191	HIS
1	D	118	ASN
1	D	148	HIS
1	D	186	GLN
1	D	335	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ALC	J	403	2	9,11,12	0.42	0	11,13,15	1.83	1 (9%)
2	ALC	K	403	2	9,11,12	0.41	0	11,13,15	0.72	1 (9%)
2	ALC	I	403	2	9,11,12	0.39	0	11,13,15	1.82	1 (9%)
2	ALC	H	405	2	9,11,12	0.51	0	11,13,15	1.03	1 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALC	J	403	2	-	2/5/14/16	0/1/1/1
2	ALC	K	403	2	-	0/5/14/16	0/1/1/1
2	ALC	I	403	2	-	4/5/14/16	0/1/1/1
2	ALC	H	405	2	-	0/5/14/16	0/1/1/1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	403	ALC	CG-CB-CA	5.84	122.37	114.52
2	I	403	ALC	CG-CB-CA	5.63	122.08	114.52
2	H	405	ALC	CG-CB-CA	-3.23	110.17	114.52
2	K	403	ALC	CG-CB-CA	-2.18	111.58	114.52

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	I	403	ALC	O-C-CA-CB
2	I	403	ALC	CA-CB-CG-CD2
2	J	403	ALC	CA-CB-CG-CD2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	I	403	ALC	CA-CB-CG-CD1
2	J	403	ALC	CA-CB-CG-CD1
2	I	403	ALC	N-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	K	403	ALC	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	A	401	-	5,5,5	0.21	0	5,5,5	0.31	0
4	PEG	A	402	-	6,6,6	0.67	0	5,5,5	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	401	-	-	0/4/4/4	-
4	PEG	A	402	-	-	1/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	402	PEG	O2-C3-C4-O4

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	H	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	H	403:SGK	C	404:GLN	N	1.19

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	369/369 (100%)	0.28	13 (3%)	47	53	17, 39, 75, 89	8 (2%)
1	B	369/369 (100%)	0.26	13 (3%)	47	53	14, 39, 74, 89	7 (1%)
1	C	368/369 (99%)	0.93	48 (13%)	7	8	22, 59, 96, 114	3 (0%)
1	D	368/369 (99%)	1.17	68 (18%)	3	3	26, 63, 118, 145	1 (0%)
2	H	4/6 (66%)	0.23	0	100	100	40, 40, 46, 48	0
2	I	4/6 (66%)	0.52	0	100	100	35, 39, 43, 45	0
2	J	4/6 (66%)	0.51	0	100	100	38, 44, 50, 53	0
2	K	4/6 (66%)	1.36	1 (25%)	2	1	49, 57, 60, 67	0
All	All	1490/1500 (99%)	0.66	143 (9%)	13	15	14, 49, 95, 145	19 (1%)

All (143) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	256	LEU	4.9
1	A	20	PRO	4.9
1	D	353	ALA	4.8
1	C	284	TYR	4.4
1	D	357	ALA	4.2
1	D	285	VAL	4.0
1	D	336	VAL	4.0
1	A	366	LEU	3.9
1	C	129	THR	3.9
1	D	310	TYR	3.8
1	B	103[A]	ARG	3.8
1	D	313	ALA	3.6
1	D	312	GLY	3.6
1	D	340	LEU	3.6
1	D	339	MET	3.5
1	C	164	GLY	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	306	LEU	3.4
1	D	360	VAL	3.4
1	D	247	VAL	3.4
1	D	308	VAL	3.4
1	A	21	LEU	3.3
1	D	290	LEU	3.3
1	D	338	MET	3.3
1	D	347	VAL	3.3
1	C	128	PHE	3.3
1	D	259	GLY	3.3
1	C	21	LEU	3.3
1	D	184	ILE	3.2
1	D	345[A]	SER	3.2
1	C	352	ALA	3.2
1	D	341	THR	3.2
1	D	252	PRO	3.2
1	C	60	VAL	3.2
1	C	167	LEU	3.2
1	C	59	LEU	3.1
1	C	38	ALA	3.1
1	C	10	LEU	3.1
1	D	309	THR	3.0
1	C	184	ILE	3.0
1	C	126	VAL	3.0
1	D	258	ALA	3.0
1	D	358	ALA	3.0
1	C	256	LEU	2.9
1	D	253	ASP	2.9
1	D	352	ALA	2.9
1	C	11	LEU	2.9
1	C	245[A]	ARG	2.9
1	D	283	LEU	2.9
1	D	305	ILE	2.8
1	D	317	ILE	2.8
1	C	191	HIS	2.8
1	C	99	VAL	2.8
1	A	119	LEU	2.8
1	B	188	LEU	2.8
1	D	248	LEU	2.8
1	D	349	ILE	2.8
1	D	59	LEU	2.8
1	D	343	SER	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	244	TYR	2.7
1	C	187	SER	2.7
1	D	251	ASN	2.7
1	D	337	ARG	2.7
1	D	354	SER	2.7
1	D	287	GLU	2.7
1	C	259	GLY	2.6
1	C	188	LEU	2.6
2	K	405	LEU	2.6
1	C	189	PRO	2.6
1	D	288	ASN	2.6
1	B	253	ASP	2.5
1	D	356	SER	2.5
1	C	5	VAL	2.5
1	D	284	TYR	2.5
1	C	92	LEU	2.5
1	D	263	LEU	2.5
1	A	126	VAL	2.5
1	D	361	VAL	2.5
1	D	333	CYS	2.4
1	D	38	ALA	2.4
1	D	179	VAL	2.4
1	D	237	VAL	2.4
1	A	120	ASP	2.4
1	C	131	PRO	2.4
1	D	255	HIS	2.4
1	A	338	MET	2.4
1	B	21	LEU	2.4
1	D	286	SER	2.4
1	C	331	LEU	2.3
1	D	331	LEU	2.3
1	D	60	VAL	2.3
1	D	188	LEU	2.3
1	A	118	ASN	2.3
1	D	315	MET	2.3
1	A	-2	GLY	2.3
1	D	355	GLN	2.3
1	C	122	TRP	2.3
1	C	138	LEU	2.3
1	C	41	THR	2.3
1	C	338	MET	2.3
1	C	336	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	344	VAL	2.3
1	A	148	HIS	2.3
1	C	255	HIS	2.3
1	C	130	LEU	2.3
1	B	252	PRO	2.2
1	C	285	VAL	2.2
1	B	119	LEU	2.2
1	D	262	LEU	2.2
1	B	184	ILE	2.2
1	C	218	ILE	2.2
1	C	219	GLY	2.2
1	C	354	SER	2.2
1	D	189	PRO	2.2
1	D	209	GLY	2.2
1	C	9	HIS	2.1
1	B	124	SER	2.1
1	C	185	GLY	2.1
1	C	209	GLY	2.1
1	A	188	LEU	2.1
1	D	20	PRO	2.1
1	C	353	ALA	2.1
1	D	58	ALA	2.1
1	C	244	TYR	2.1
1	B	117	PRO	2.1
1	D	363	PRO	2.1
1	B	118	ASN	2.1
1	C	118	ASN	2.1
1	B	255	HIS	2.1
1	D	26	THR	2.0
1	D	182	MET	2.0
1	A	210	GLY	2.0
1	B	249	PRO	2.0
1	C	227	VAL	2.0
1	D	170	VAL	2.0
1	D	119	LEU	2.0
1	C	313	ALA	2.0
1	A	253	ASP	2.0
1	D	254	LYS	2.0
1	C	230	PHE	2.0
1	B	239	GLY	2.0
1	C	102	GLY	2.0
1	C	65	PRO	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	246	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ALC	H	405	11/12	0.90	0.09	33,37,41,42	0
2	ALC	K	403	11/12	0.90	0.11	51,55,59,61	0
2	ALC	I	403	11/12	0.94	0.09	34,40,50,53	0
2	ALC	J	403	11/12	0.95	0.08	40,44,48,50	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PEG	A	402	7/7	0.86	0.10	20,20,21,32	7
3	GOL	A	401	6/6	0.87	0.10	28,29,30,30	6

6.5 Other polymers [i](#)

There are no such residues in this entry.