



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 12:04 AM UTC

PDB ID : 3B0U / pdb_00003b0u
Title : tRNA-dihydrouridine synthase from *Thermus thermophilus* in complex with tRNA fragment
Authors : Yu, F.; Tanaka, Y.; Yamashita, K.; Nakamura, A.; Yao, M.; Tanaka, I.
Deposited on : 2011-06-14
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)
RNA backbone	3983	1010 (2.30-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	4	75% 25% 50% 25%
1	B	4	25% 25% 50% 25%
2	X	350	% 81% 9% 11%
2	Y	350	78% 11% 11%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
1	H2U	B	20	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5640 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA (5'-R(*GP*GP*(H2U)P*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	4	Total	C	N	O	P	0	0	0
			85	39	17	26	3			
1	B	4	Total	C	N	O	P	0	0	0
			85	39	17	26	3			


- Molecule 2 is a protein called tRNA-dihydrouridine synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	X	312	Total	C	N	O	S	0	2	0
			2495	1574	474	435	12			
2	Y	312	Total	C	N	O	S	0	2	0
			2495	1574	474	435	12			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	343	LEU	-	expression tag	UNP Q5SMC7
X	344	GLU	-	expression tag	UNP Q5SMC7
X	345	HIS	-	expression tag	UNP Q5SMC7
X	346	HIS	-	expression tag	UNP Q5SMC7
X	347	HIS	-	expression tag	UNP Q5SMC7
X	348	HIS	-	expression tag	UNP Q5SMC7
X	349	HIS	-	expression tag	UNP Q5SMC7
X	350	HIS	-	expression tag	UNP Q5SMC7
Y	343	LEU	-	expression tag	UNP Q5SMC7
Y	344	GLU	-	expression tag	UNP Q5SMC7
Y	345	HIS	-	expression tag	UNP Q5SMC7
Y	346	HIS	-	expression tag	UNP Q5SMC7
Y	347	HIS	-	expression tag	UNP Q5SMC7
Y	348	HIS	-	expression tag	UNP Q5SMC7
Y	349	HIS	-	expression tag	UNP Q5SMC7
Y	350	HIS	-	expression tag	UNP Q5SMC7

- Molecule 3 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: $C_{17}H_{21}N_4O_9P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	X	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
3	Y	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	4	Total	O	0	0
			4	4		
4	B	7	Total	O	0	0
			7	7		
4	X	198	Total	O	0	0
			198	198		
4	Y	209	Total	O	0	0
			209	209		


- Molecule 1: RNA (5'-R(*GP*GP*(H2U)P*A)-3')





- Chain B: 



- Chain X:  81% 9% 11%



- Chain Y:  78% 11% 11%



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	126.64Å 126.64Å 112.67Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.80 – 1.95 19.80 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.7 (19.80-1.95) 99.6 (19.80-1.95)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.14 (at 1.94Å)	Xtriage
Refinement program	PHENIX dev_708	Depositor
R, R_{free}	0.181 , 0.220 0.175 , 0.217	Depositor DCC
R_{free} test set	2159 reflections (4.38%)	wwPDB-VP
Wilson B-factor (Å ²)	25.1	Xtriage
Anisotropy	0.340	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.028 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5640	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.45 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1424e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, H2U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.06	0/73	0.18	0/113
1	B	0.11	0/73	0.25	0/113
2	X	0.29	0/2543	0.65	0/3428
2	Y	0.29	0/2543	0.67	2/3428 (0.1%)
All	All	0.28	0/5232	0.65	2/7082 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	Y	68	ASP	CA-C-N	5.25	124.70	119.24
2	Y	68	ASP	C-N-CA	5.25	124.70	119.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	85	0	46	8	0
1	B	85	0	47	13	0
2	X	2495	0	2570	25	0
2	Y	2495	0	2571	40	0
3	X	31	0	19	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	Y	31	0	19	2	0
4	A	4	0	0	0	0
4	B	7	0	0	0	0
4	X	198	0	0	0	0
4	Y	209	0	0	1	0
All	All	5640	0	5272	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:20:H2U:H51	2:Y:93:CYS:SG	1.11	1.64
1:A:20:H2U:C5	2:X:93:CYS:SG	2.05	1.44
1:B:20:H2U:C5	2:Y:93:CYS:SG	2.08	1.39
1:B:20:H2U:H51	2:Y:93:CYS:HG	1.32	0.93
1:B:18:G:H1	2:Y:49:ARG:HH12	1.27	0.82
1:B:18:G:H22	2:Y:49:ARG:HH22	1.30	0.79
2:X:134:ARG:HD2	2:X:168:ALA:HB2	1.69	0.74
1:B:20:H2U:H51	2:Y:93:CYS:CB	2.21	0.69
1:A:20:H2U:C4	2:X:93:CYS:SG	2.81	0.69
2:Y:134:ARG:CD	2:Y:168:ALA:HB2	2.27	0.65
2:Y:153:MET:HE3	2:Y:161:PHE:HE1	1.63	0.64
2:Y:134:ARG:HD2	2:Y:168:ALA:HB2	1.82	0.62
2:X:134:ARG:CD	2:X:168:ALA:HB2	2.32	0.60
1:B:20:H2U:C4	2:Y:93:CYS:SG	2.90	0.59
1:A:20:H2U:H4'	1:A:21:A:OP2	2.04	0.58
2:X:310:ARG:O	2:X:313:GLU:HG2	2.06	0.56
1:B:18:G:N2	2:Y:49:ARG:HH22	2.01	0.55
2:Y:265:GLU:O	2:Y:268:LYS:HG3	2.06	0.55
2:Y:138:GLU:HB2	2:Y:140:LYS:HE2	1.88	0.53
1:A:19:G:H5''	2:X:38:VAL:HG21	1.89	0.53
2:Y:265:GLU:HG3	2:Y:270:THR:CB	2.38	0.53
1:A:18:G:C3'	1:A:19:G:H5'	2.39	0.52
2:Y:133:MET:SD	2:Y:153:MET:HE1	2.50	0.52
2:X:292:TRP:CE3	2:X:308:ALA:HB1	2.46	0.51
2:Y:310:ARG:O	2:Y:313:GLU:HG2	2.11	0.50
2:X:16:ASP:O	2:X:20:ARG:HG3	2.12	0.50
2:X:267:LEU:C	2:X:267:LEU:HD23	2.37	0.50
2:Y:11:MET:HA	3:Y:400:FMN:C5A	2.42	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:X:22:LEU:C	2:X:22:LEU:HD23	2.36	0.49
2:Y:115:ARG:HG3	2:Y:156:ALA:HB2	1.94	0.49
1:A:20:H2U:C6	2:X:93:CYS:SG	2.96	0.48
2:X:136:GLY:HA3	2:X:141:GLU:HB3	1.95	0.48
2:Y:144:ARG:HD2	4:Y:406:HOH:O	2.12	0.48
2:Y:265:GLU:HG3	2:Y:270:THR:HB	1.97	0.47
2:Y:153:MET:HE3	2:Y:161:PHE:CE1	2.47	0.47
2:X:11:MET:HA	3:X:400:FMN:C5A	2.45	0.47
1:B:20:H2U:C6	2:Y:93:CYS:SG	2.95	0.47
1:B:18:G:H1	2:Y:49:ARG:NH1	2.03	0.47
2:Y:22:LEU:C	2:Y:22:LEU:HD23	2.40	0.47
2:Y:16:ASP:O	2:Y:20:ARG:HG3	2.14	0.46
2:X:108:LEU:HB2	2:X:137:LEU:HD21	1.97	0.46
2:Y:306:ASP:O	2:Y:310:ARG:HG3	2.15	0.46
2:Y:152:ALA:O	2:Y:155:GLU:HG2	2.15	0.45
1:B:20:H2U:C5	2:Y:93:CYS:CB	2.89	0.45
2:Y:8:VAL:HG21	2:Y:23:VAL:HG11	1.98	0.45
2:X:260:ALA:O	2:X:264:GLU:HG3	2.17	0.45
1:B:18:G:H22	2:Y:49:ARG:NH2	2.06	0.44
2:Y:136:GLY:HA3	2:Y:141:GLU:HB3	1.99	0.44
2:X:136:GLY:C	2:X:137:LEU:HD22	2.43	0.44
2:Y:140:LYS:HB2	2:Y:140:LYS:HE3	1.78	0.44
2:Y:262:LEU:O	2:Y:266:VAL:HG23	2.18	0.44
1:A:20:H2U:OP1	2:X:97:LYS:HD2	2.18	0.44
2:Y:11:MET:HA	3:Y:400:FMN:N5	2.33	0.44
2:Y:49:ARG:CZ	2:Y:49:ARG:HB3	2.48	0.44
1:A:20:H2U:C5	2:X:93:CYS:CB	2.94	0.43
2:X:8:VAL:HG21	2:X:23:VAL:HG11	1.99	0.43
2:X:108:LEU:HB2	2:X:137:LEU:CD2	2.48	0.43
2:X:12:VAL:HG23	3:X:400:FMN:O4	2.19	0.43
2:X:11:MET:HA	3:X:400:FMN:N5	2.34	0.42
2:X:209:LEU:HD12	2:X:209:LEU:HA	1.93	0.41
2:X:4:PRO:O	2:X:221:GLY:HA3	2.21	0.41
2:X:130:THR:HG22	2:X:160:VAL:HB	2.01	0.41
2:Y:159:LYS:HE3	2:Y:197:GLN:HE22	1.85	0.41
2:Y:141:GLU:CD	2:Y:141:GLU:H	2.29	0.41
2:Y:102:GLY:HA2	2:Y:106[A]:CYS:SG	2.60	0.40
2:Y:130:THR:HG22	2:Y:160:VAL:HB	2.03	0.40
1:B:20:H2U:C5	2:Y:93:CYS:HG	2.08	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	X	312/350 (89%)	299 (96%)	13 (4%)	0	100	100
2	Y	312/350 (89%)	301 (96%)	11 (4%)	0	100	100
All	All	624/700 (89%)	600 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	X	260/287 (91%)	258 (99%)	2 (1%)	73	71
2	Y	260/287 (91%)	257 (99%)	3 (1%)	63	57
All	All	520/574 (91%)	515 (99%)	5 (1%)	68	64

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	X	170	LEU
2	X	263	GLU
2	Y	137	LEU
2	Y	170	LEU
2	Y	250	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	X	203	ASN
2	Y	203	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	3/4 (75%)	3 (100%)	0
1	B	3/4 (75%)	2 (66%)	0
All	All	6/8 (75%)	5 (83%)	0

All (5) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	19	G
1	A	20	H2U
1	A	21	A
1	B	20	H2U
1	B	21	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	H2U	B	20	1	18,21,22	0.64	0	19,30,33	1.49	3 (15%)
1	H2U	A	20	1	18,21,22	0.69	0	19,30,33	1.54	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	H2U	B	20	1	-	2/7/38/39	0/2/2/2
1	H2U	A	20	1	-	2/7/38/39	0/2/2/2

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	20	H2U	N3-C2-N1	4.60	121.27	116.65
1	B	20	H2U	N3-C2-N1	4.29	120.96	116.65
1	A	20	H2U	O2-C2-N1	-3.09	119.39	123.10
1	A	20	H2U	C5-C4-N3	2.92	119.80	116.69
1	B	20	H2U	C5-C4-N3	2.88	119.75	116.69
1	B	20	H2U	O2-C2-N1	-2.86	119.67	123.10

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	20	H2U	O4'-C4'-C5'-O5'
1	A	20	H2U	C3'-C4'-C5'-O5'
1	B	20	H2U	C3'-C4'-C5'-O5'
1	B	20	H2U	O4'-C4'-C5'-O5'

There are no ring outliers.

2 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	20	H2U	8	0
1	A	20	H2U	6	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FMN	X	400	-	33,33,33	1.05	2 (6%)	48,50,50	1.23	7 (14%)
3	FMN	Y	400	-	33,33,33	1.03	2 (6%)	48,50,50	1.18	6 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FMN	X	400	-	-	1/18/18/18	0/3/3/3
3	FMN	Y	400	-	-	1/18/18/18	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	X	400	FMN	C4A-N5	3.66	1.38	1.30
3	Y	400	FMN	C4A-N5	3.66	1.38	1.30
3	Y	400	FMN	C10-N1	2.62	1.38	1.33
3	X	400	FMN	C10-N1	2.56	1.38	1.33

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	X	400	FMN	C4-N3-C2	-3.06	120.22	125.64
3	Y	400	FMN	C4-N3-C2	-2.77	120.73	125.64
3	X	400	FMN	C4A-C4-N3	2.62	119.92	113.25
3	Y	400	FMN	C9A-C5A-N5	-2.61	119.68	122.45
3	Y	400	FMN	C4A-C4-N3	2.55	119.75	113.25
3	X	400	FMN	C4A-C10-N10	2.50	120.06	116.48
3	X	400	FMN	C9A-C5A-N5	-2.46	119.85	122.45
3	X	400	FMN	C10-C4A-N5	-2.45	119.81	124.81
3	Y	400	FMN	C10-C4A-N5	-2.43	119.85	124.81
3	Y	400	FMN	C4A-C10-N10	2.42	119.95	116.48
3	X	400	FMN	O4-C4-C4A	-2.32	120.41	126.53
3	X	400	FMN	C5A-C9A-N10	2.17	119.93	117.97
3	Y	400	FMN	O4-C4-C4A	-2.14	120.89	126.53

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	X	400	FMN	C4'-C5'-O5'-P
3	Y	400	FMN	C4'-C5'-O5'-P


There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	X	400	FMN	3	0
3	Y	400	FMN	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	3/4 (75%)	3.07	3 (100%) 0 0	95, 95, 101, 105	0
1	B	3/4 (75%)	1.97	1 (33%) 1 0	53, 53, 60, 98	0
2	X	312/350 (89%)	-0.29	5 (1%) 70 77	11, 27, 49, 78	2 (0%)
2	Y	312/350 (89%)	-0.25	1 (0%) 90 93	11, 27, 49, 70	2 (0%)
All	All	630/708 (88%)	-0.24	10 (1%) 70 77	11, 27, 51, 105	4 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	X	267	LEU	3.7
1	B	21	A	3.2
1	A	19	G	3.2
1	A	21	A	3.0
1	A	18	G	3.0
2	X	246	PRO	2.6
2	Y	267	LEU	2.6
2	X	140	LYS	2.5
2	X	137	LEU	2.3
2	X	197	GLN	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	H2U	A	20	20/21	0.91	0.09	18,29,87,88	0
1	H2U	B	20	20/21	0.95	0.07	17,25,32,35	0

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	FMN	X	400	31/31	0.97	0.06	10,17,21,23	0
3	FMN	Y	400	31/31	0.97	0.06	13,18,21,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around FMN Y 400:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.