



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 10:03 PM UTC

PDB ID : 6B0G / pdb_00006b0g
Title : Crystal structure of Pfs25 in complex with the transmission blocking antibody 1245
Authors : McLeod, B.; Scally, S.W.; Bosch, A.; King, C.R.; Julien, J.P.
Deposited on : 2017-09-14
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	C	219	 91% 8%
2	D	228	 94% 5%
3	E	183	 4% 78% 7% 16%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5121 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 1245 Antibody, Light Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	C	218	Total	C	N	O	S	0	0	0
			1685	1051	289	339	6			

- Molecule 2 is a protein called 1245 Antibody, Heavy Chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	226	Total	C	N	O	S	0	1	0
			1706	1072	289	336	9			


- Molecule 3 is a protein called 25 kDa ookinete surface antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	E	154	Total	C	N	O	S	0	2	0
			1154	704	195	230	25			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	THR	-	expression tag	UNP P13829
E	0	GLY	-	expression tag	UNP P13829
E	91	GLN	ASN	conflict	UNP P13829
E	144	GLN	ASN	conflict	UNP P13829
E	166	GLN	ASN	conflict	UNP P13829
E	173	GLY	-	expression tag	UNP P13829
E	174	THR	-	expression tag	UNP P13829
E	175	LYS	-	expression tag	UNP P13829
E	176	HIS	-	expression tag	UNP P13829
E	177	HIS	-	expression tag	UNP P13829
E	178	HIS	-	expression tag	UNP P13829
E	179	HIS	-	expression tag	UNP P13829
E	180	HIS	-	expression tag	UNP P13829
E	181	HIS	-	expression tag	UNP P13829

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	D	1	Total	C	O	0	0
			6	3	3		
4	E	1	Total	C	O	0	0
			6	3	3		
4	E	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	C	191	Total	O	0	0
			191	191		
5	D	213	Total	O	0	0
			213	213		
5	E	142	Total	O	0	0
			142	142		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 1245 Antibody, Light Chain

Chain C:  91% 8%




- Molecule 2: 1245 Antibody, Heavy Chain

Chain D:  94% 5%



- Molecule 3: 25 kDa ookinete surface antigen

Chain E:  4% 78% 7% 16%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.77Å 83.69Å 76.21Å 90.00° 93.65° 90.00°	Depositor
Resolution (Å)	38.99 – 1.90 38.99 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (38.99-1.90) 93.2 (38.99-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 1.89Å)	Xtriage
Refinement program	PHENIX (1.11.1 _2575: ???)	Depositor
R, R_{free}	0.164 , 0.200 0.164 , 0.200	Depositor DCC
R_{free} test set	2003 reflections (3.92%)	wwPDB-VP
Wilson B-factor (Å ²)	17.9	Xtriage
Anisotropy	0.164	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5121	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	C	0.44	0/1723	0.64	0/2340
2	D	0.49	0/1750	0.68	0/2379
3	E	0.42	0/1169	0.64	0/1570
All	All	0.46	0/4642	0.65	0/6289

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1685	0	1633	13	0
2	D	1706	0	1667	7	0
3	E	1154	0	1133	7	0
4	D	18	0	24	2	0
4	E	12	0	16	1	0
5	C	191	0	0	1	1
5	D	213	0	0	2	2
5	E	142	0	0	1	0
All	All	5121	0	4473	24	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 24 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:123:PRO:HD3	2:D:214:LYS:HE2	1.72	0.69
3:E:73:CYS:SG	3:E:79:MET:HG2	2.38	0.62
1:C:13:VAL:HG21	1:C:78:VAL:HG11	1.80	0.62
3:E:142:LYS:NZ	4:E:202:GOL:O2	2.25	0.59
1:C:116:PHE:HD1	2:D:135:SER:HA	1.68	0.58

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:C:367:HOH:O	5:D:489:HOH:O[2_1558]	2.15	0.05
5:D:451:HOH:O	5:D:538:HOH:O[1_455]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	216/219 (99%)	213 (99%)	3 (1%)	0	100	100
2	D	225/228 (99%)	221 (98%)	4 (2%)	0	100	100
3	E	152/183 (83%)	149 (98%)	3 (2%)	0	100	100
All	All	593/630 (94%)	583 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	194/195 (100%)	193 (100%)	1 (0%)	81	84
2	D	190/191 (100%)	187 (98%)	3 (2%)	55	54
3	E	141/166 (85%)	140 (99%)	1 (1%)	76	78
All	All	525/552 (95%)	520 (99%)	5 (1%)	68	70

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	105	GLU
2	D	143	LEU
2	D	154	PRO
2	D	183	LEU
3	E	50	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	160	GLN
2	D	43	GLN
2	D	79	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	E	202	-	5,5,5	0.32	0	5,5,5	0.58	0
4	GOL	E	201	-	5,5,5	0.35	0	5,5,5	0.44	0
4	GOL	D	302	-	5,5,5	0.38	0	5,5,5	0.57	0
4	GOL	D	301	-	5,5,5	0.38	0	5,5,5	0.36	0
4	GOL	D	303	-	5,5,5	0.57	0	5,5,5	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	E	202	-	-	0/4/4/4	-
4	GOL	E	201	-	-	2/4/4/4	-
4	GOL	D	302	-	-	3/4/4/4	-
4	GOL	D	301	-	-	2/4/4/4	-
4	GOL	D	303	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	302	GOL	O1-C1-C2-C3
4	D	303	GOL	O1-C1-C2-O2
4	D	303	GOL	O1-C1-C2-C3
4	E	201	GOL	O1-C1-C2-C3
4	D	301	GOL	O1-C1-C2-C3

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	E	202	GOL	1	0
4	D	303	GOL	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	C	218/219 (99%)	-0.37	1 (0%)	87 89	12, 24, 47, 77	0
2	D	226/228 (99%)	-0.48	1 (0%)	88 90	11, 20, 38, 62	1 (0%)
3	E	154/183 (84%)	0.12	8 (5%)	33 35	11, 32, 63, 80	2 (1%)
All	All	598/630 (94%)	-0.28	10 (1%)	69 72	11, 24, 52, 80	3 (0%)

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	E	67	VAL	4.3
3	E	62	ILE	4.3
2	D	220	SER	2.7
3	E	69	TYR	2.5
3	E	66	PRO	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	E	202	6/6	0.75	0.18	56,61,66,72	0
4	GOL	D	302	6/6	0.84	0.15	41,53,55,56	0
4	GOL	E	201	6/6	0.87	0.14	50,59,61,64	0
4	GOL	D	301	6/6	0.88	0.13	51,58,59,60	0
4	GOL	D	303	6/6	0.89	0.13	24,31,35,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.