



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:19 PM UTC

PDB ID : 8B0D / pdb_00008b0d
Title : Crystal structure of beta-glucuronidase from *Acidobacterium capsulatum* in complex with covalent inhibitor VB151
Authors : Armstrong, Z.; Davies, G.J.
Deposited on : 2022-09-07
Resolution : 1.62 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.62 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6728 (1.64-1.60)
Clashscore	190562	7023 (1.64-1.60)
Ramachandran outliers	187476	6898 (1.64-1.60)
Sidechain outliers	187428	6896 (1.64-1.60)
RSRZ outliers	180081	6727 (1.64-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	475	9% 81% 10% 9%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 6784 atoms, of which 3216 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-glucuronidase from *Acidobacterium capsulatum*.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	AAA	433	6493	2101	3192	565	627	8	156	1	0

- Molecule 2 is ALANINE (CCD ID: ALA) (formula: $C_3H_7NO_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	AAA	1	12	3	7	1	1	0	0

- Molecule 3 is (1 {S},2 {R},3 {R},4 {S},6 {S})-2-(2-acetamidoethoxy)-3,4,6-tris(oxidanyl)cyclohexane-1-carboxylic acid (CCD ID: OUU) (formula: $C_{11}H_{19}NO_7$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	AAA	1	Total	C	H	N	O	3	0
			36	11	17	1	7		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	AAA	1	Total	O S	0	0
			5	4 1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	AAA	238	Total 238	O 238	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: beta-glucuronidase from *Acidobacterium capsulatum*



4 Data and refinement statistics

Property	Value	Source
Space group	I 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	82.46Å 43.60Å 136.20Å 90.00° 98.69° 90.00°	Depositor
Resolution (Å)	74.92 – 1.62 74.92 – 1.62	Depositor EDS
% Data completeness (in resolution range)	99.6 (74.92-1.62) 99.6 (74.92-1.62)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 1.62Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.198 , 0.226 0.203 , 0.232	Depositor DCC
R_{free} test set	2970 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	23.7	Xtriage
Anisotropy	0.258	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6784	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, OUU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	1.16	4/3387 (0.1%)	1.31	9/4609 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	AAA	474	PHE	C-N	25.67	1.93	1.34
1	AAA	173	GLU	N-CA	5.24	1.50	1.46
1	AAA	287	GLU	CD-OE2	5.13	1.35	1.25
1	AAA	191	ALA	C-O	5.04	1.29	1.24

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	474	PHE	O-C-N	-10.30	106.22	122.70
1	AAA	83	GLU	CB-CG-CD	7.13	124.72	112.60
1	AAA	18	SER	CA-C-N	5.30	125.76	120.14
1	AAA	18	SER	C-N-CA	5.30	125.76	120.14
1	AAA	474	PHE	CA-C-O	5.27	129.76	120.80
1	AAA	161	GLY	CA-C-O	-5.25	118.04	122.29
1	AAA	256	GLU	CA-C-N	5.17	127.21	120.28
1	AAA	256	GLU	C-N-CA	5.17	127.21	120.28
1	AAA	218	ALA	CA-C-O	-5.13	116.11	121.55

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	3301	3192	3178	23	0
2	AAA	5	7	4	1	0
3	AAA	19	17	0	0	0
4	AAA	5	0	0	0	0
5	AAA	238	0	0	1	0
All	All	3568	3216	3182	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:474:PHE:C	2:AAA:501:ALA:N	1.93	1.25
1:AAA:83:GLU:HG3	1:AAA:142:GLY:HA3	1.71	0.73
1:AAA:48:GLN:HG3	1:AAA:54:TYR:CD1	2.40	0.57
1:AAA:465:ARG:NH1	5:AAA:604:HOH:O	2.37	0.56
1:AAA:225:VAL:HB	1:AAA:226:PRO:HD3	1.89	0.54
1:AAA:217:THR:O	1:AAA:240:SER:HA	2.07	0.54
1:AAA:232:LYS:HA	1:AAA:281:LEU:HD11	1.92	0.51
1:AAA:48:GLN:CG	1:AAA:54:TYR:CD1	2.96	0.49
1:AAA:259:MET:O	1:AAA:398[B]:LYS:NZ	2.42	0.48
1:AAA:85:THR:HG22	1:AAA:116:ILE:HG13	1.97	0.47
1:AAA:164:ARG:HA	1:AAA:164:ARG:HD2	1.76	0.45
1:AAA:274:GLN:HE21	1:AAA:278:ASP:CG	2.25	0.45
1:AAA:300:SER:HB3	1:AAA:334:TYR:HB3	1.99	0.43
1:AAA:165:LEU:HD21	1:AAA:168:PHE:CE2	2.54	0.43
1:AAA:128:LEU:O	1:AAA:164:ARG:NH2	2.52	0.43
1:AAA:418:VAL:HG23	1:AAA:472:LEU:HD12	2.01	0.42
1:AAA:425:ARG:NH1	1:AAA:427:ASP:OD1	2.47	0.42
1:AAA:85:THR:CG2	1:AAA:116:ILE:HG13	2.49	0.42
1:AAA:135:LEU:HD22	1:AAA:160:ILE:HD13	2.02	0.42
1:AAA:274:GLN:NE2	1:AAA:278:ASP:OD1	2.52	0.42
1:AAA:69:LEU:HD22	1:AAA:354:MET:HB3	2.03	0.41
1:AAA:224:LEU:C	1:AAA:224:LEU:HD23	2.46	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:AAA:39:TYR:CD2	1:AAA:323:GLY:HA2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	AAA	429/475 (90%)	418 (97%)	10 (2%)	1 (0%)	43 25

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	454	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	AAA	331/357 (93%)	323 (98%)	8 (2%)	43 18

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	AAA	71	ARG
1	AAA	83	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	AAA	180	ASN
1	AAA	334	TYR
1	AAA	346	ILE
1	AAA	413	SER
1	AAA	457	SER
1	AAA	459	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ALA	AAA	501	-	3,4,5	0.68	0	2,4,6	2.56	1 (50%)
4	SO4	AAA	503	-	4,4,4	0.32	0	6,6,6	0.11	0
3	OUU	AAA	502	1	19,19,19	0.66	1 (5%)	23,26,26	1.09	2 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALA	AAA	501	-	-	0/1/2/4	-
3	OUU	AAA	502	1	-	2/11/31/31	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	AAA	502	OUU	O6B-C6	-2.34	1.23	1.30

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	AAA	501	ALA	O-C-CA	-3.59	112.73	124.23
3	AAA	502	OUU	C7-C1-C2	-2.75	108.30	111.50
3	AAA	502	OUU	O6A-C6-C5	-2.56	116.30	122.84

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	AAA	502	OUU	C4-C5-C6-O6A
3	AAA	502	OUU	C4-C5-C6-O6B


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	AAA	501	ALA	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	433/475 (91%)	0.66	41 (9%) 14 14	17, 35, 59, 87	3 (0%)

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	458	GLY	5.1
1	AAA	30	ALA	4.3
1	AAA	454	PRO	4.0
1	AAA	455	GLY	3.7
1	AAA	456	HIS	3.7
1	AAA	450	GLN	3.6
1	AAA	247	PRO	3.5
1	AAA	18	SER	3.4
1	AAA	250	ASP	3.3
1	AAA	178	TYR	3.3
1	AAA	474	PHE	3.3
1	AAA	84	TYR	3.2
1	AAA	219	TYR	3.2
1	AAA	181	GLY	3.1
1	AAA	412	ALA	3.1
1	AAA	90	HIS	3.0
1	AAA	22	VAL	2.8
1	AAA	414	PRO	2.7
1	AAA	453	VAL	2.6
1	AAA	180	ASN	2.6
1	AAA	472	LEU	2.5
1	AAA	218	ALA	2.5
1	AAA	72	GLN	2.4
1	AAA	221	THR	2.4
1	AAA	452	TYR	2.4
1	AAA	449	VAL	2.4
1	AAA	86	PHE	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	AAA	418	VAL	2.4
1	AAA	389	GLY	2.3
1	AAA	457	SER	2.3
1	AAA	31	LEU	2.3
1	AAA	411	VAL	2.3
1	AAA	223	TRP	2.2
1	AAA	71	ARG	2.2
1	AAA	224	LEU	2.2
1	AAA	179	ARG	2.1
1	AAA	365	LEU	2.1
1	AAA	35	ILE	2.1
1	AAA	292	TYR	2.1
1	AAA	369	LYS	2.1
1	AAA	470	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ALA	AAA	501	5/6	0.84	0.20	50,56,58,60	0
4	SO4	AAA	503	5/5	0.87	0.09	68,68,72,72	0
3	OUU	AAA	502	19/19	0.96	0.07	26,30,39,44	3

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around OUU AAA 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.