



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 05:55 PM UTC

PDB ID : 7B1X / pdb_00007b1x
Title : Crystal structure of cold-active esterase PMGL3 from permafrost metagenomic library
Authors : Boyko, K.M.; Nikolaeva, A.Y.; Petrovskaya, L.E.; Kryukova, M.V.; Kryukova, E.A.; Korzhenevsky, D.A.; Lomakina, G.Y.; Novototskaya-Vlasova, K.A.; Rivkina, E.M.; Dolgikh, D.A.; Kirpichnikov, M.P.; Popov, V.O.
Deposited on : 2020-11-25
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	
1	B	312	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4716 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called esterase PMGL3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	302	Total	C	N	O	S	0	1	0
			2259	1430	396	423	10			
1	B	302	Total	C	N	O	S	0	1	0
			2260	1429	397	424	10			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	98	Total	O	0	0
			98	98		
2	B	99	Total	O	0	0
			99	99		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: esterase PMGL3



• Molecule 1: esterase PMGL3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	103.80Å 98.40Å 75.20Å 90.00° 128.10° 90.00°	Depositor
Resolution (Å)	37.28 – 2.30 37.28 – 2.30	Depositor EDS
% Data completeness (in resolution range)	89.2 (37.28-2.30) 89.2 (37.28-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.12 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.162 , 0.231 (Not available) , 0.203	Depositor DCC
R_{free} test set	1494 reflections (4.30%)	wwPDB-VP
Wilson B-factor (Å ²)	37.5	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4716	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	1/2319 (0.0%)	1.44	23/3158 (0.7%)
1	B	0.85	2/2322 (0.1%)	1.44	15/3164 (0.5%)
All	All	0.87	3/4641 (0.1%)	1.44	38/6322 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	83	HIS	CE1-NE2	6.08	1.38	1.32
1	B	40	HIS	CE1-NE2	5.54	1.38	1.32
1	B	91	HIS	CE1-NE2	5.28	1.37	1.32

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	228	ASP	CB-CA-C	8.92	124.44	109.80
1	B	254	ARG	CG-CD-NE	-8.75	92.75	112.00
1	B	270	ASN	CA-CB-CG	-8.31	104.28	112.60
1	A	270	ASN	CA-CB-CG	-8.19	104.41	112.60
1	A	62	GLU	CB-CA-C	-7.97	96.42	109.89
1	A	254	ARG	CG-CD-NE	-7.91	94.59	112.00
1	B	62	GLU	CB-CA-C	-7.80	96.71	109.89
1	B	202	ASN	CB-CA-C	6.81	122.10	110.79
1	B	4	ASN	CA-CB-CG	-6.66	105.94	112.60
1	A	23	GLU	CB-CG-CD	6.63	123.87	112.60
1	A	165	GLN	CB-CA-C	-6.59	99.05	110.72

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	254	ARG	NE-CZ-NH2	-6.27	113.55	119.20
1	A	270	ASN	CB-CA-C	-6.18	98.59	110.65
1	B	124	ASP	CA-CB-CG	6.13	118.73	112.60
1	A	52	GLU	CB-CG-CD	5.94	122.70	112.60
1	A	13	GLN	CB-CA-C	-5.89	101.02	110.79
1	B	270	ASN	CB-CA-C	-5.85	99.24	110.65
1	A	202	ASN	CB-CA-C	5.80	119.99	110.88
1	A	151	GLY	CA-C-N	5.80	126.42	119.98
1	A	151	GLY	C-N-CA	5.80	126.42	119.98
1	B	5	ASP	CB-CA-C	5.75	115.86	110.17
1	A	255	ARG	NE-CZ-NH1	-5.64	115.86	121.50
1	A	49	CYS	CA-C-O	-5.43	114.97	121.11
1	A	216	SER	CA-C-N	5.43	128.10	120.28
1	A	216	SER	C-N-CA	5.43	128.10	120.28
1	A	73	LYS	CB-CA-C	5.42	119.72	109.37
1	A	282	LEU	N-CA-CB	-5.41	102.44	110.99
1	A	18	MET	CB-CG-SD	-5.38	96.54	112.70
1	A	65	THR	CA-CB-OG1	-5.37	101.55	109.60
1	B	18	MET	CA-C-N	5.36	127.46	120.28
1	B	18	MET	C-N-CA	5.36	127.46	120.28
1	A	18	MET	CA-C-N	5.23	127.29	120.28
1	A	18	MET	C-N-CA	5.23	127.29	120.28
1	A	264	ARG	CB-CG-CD	-5.21	99.33	111.30
1	B	270	ASN	OD1-CG-ND2	5.20	127.80	122.60
1	A	116	GLU	CB-CA-C	-5.17	102.20	110.79
1	B	25	GLN	CB-CG-CD	5.05	121.18	112.60
1	B	264	ARG	CB-CG-CD	5.03	122.87	111.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	301	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2259	0	2198	24	0
1	B	2260	0	2195	31	0
2	A	98	0	0	0	0
2	B	99	0	0	1	0
All	All	4716	0	4393	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:TRP:CH2	1:A:182:MET:HE3	2.17	0.79
1:B:1:MET:CE	1:B:6:PRO:HG3	2.13	0.79
1:B:275:TRP:CD2	1:B:276:PRO:HD3	2.25	0.72
1:A:179:TRP:CZ2	1:A:182:MET:HE3	2.25	0.71
1:A:275:TRP:CD2	1:A:276:PRO:HD3	2.26	0.70
1:A:251:GLU:OE2	1:A:255:ARG:HD2	1.93	0.68
1:A:28:SER:HB3	1:A:31:ILE:HB	1.75	0.67
1:B:28:SER:HB3	1:B:31:ILE:HB	1.79	0.64
1:B:182:MET:HE1	1:B:211:TYR:CE2	2.34	0.62
1:B:1:MET:HE1	1:B:6:PRO:HG3	1.83	0.60
1:A:264:ARG:HH12	1:A:299[B]:ARG:NH1	2.02	0.58
1:A:246:LEU:HD21	1:B:257:ALA:HB1	1.86	0.58
1:B:232:LEU:HB3	1:B:233:PRO:HD2	1.86	0.57
1:B:1:MET:N	1:B:1:MET:HE3	2.20	0.57
1:A:257:ALA:HB1	1:B:246:LEU:HD21	1.88	0.56
1:B:1:MET:HE3	1:B:1:MET:H3	1.71	0.55
1:A:165:GLN:O	1:A:167:LEU:HD13	2.08	0.53
1:A:126:ALA:HB2	1:A:154:LEU:HD13	1.92	0.52
1:A:73:LYS:HG2	1:A:301:LEU:HB3	1.93	0.51
1:B:73:LYS:HG2	1:B:301:LEU:HB3	1.93	0.51
1:B:60:ARG:HD2	1:B:110:GLU:HG3	1.93	0.50
1:A:232:LEU:HB3	1:A:233:PRO:HD2	1.94	0.49
1:A:40:HIS:HB3	1:A:90:SER:O	2.13	0.48
1:A:62:GLU:OE2	1:A:92:ARG:NH2	2.43	0.48
1:B:62:GLU:OE2	1:B:92:ARG:NH2	2.33	0.48
1:B:232:LEU:HB2	1:B:261:LEU:HD21	1.95	0.48
1:B:22:ARG:HB3	1:B:22:ARG:NH2	2.30	0.47
1:A:195:HIS:O	1:A:197:PRO:HD3	2.15	0.47
1:A:182:MET:HE2	1:A:208:SER:HA	1.98	0.46
1:B:126:ALA:HB2	1:B:154:LEU:HD13	1.98	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:35:MET:HE2	1:B:35:MET:HB3	1.73	0.46
1:B:40:HIS:HB3	1:B:90:SER:O	2.16	0.46
1:A:275:TRP:CG	1:A:276:PRO:HD3	2.52	0.45
1:B:195:HIS:O	1:B:197:PRO:HD3	2.16	0.45
1:B:128:GLN:OE1	1:B:128:GLN:HA	2.16	0.45
1:B:275:TRP:CG	1:B:276:PRO:HD3	2.51	0.44
1:A:180:LEU:HD13	1:A:252:PHE:CG	2.52	0.44
1:B:60:ARG:CD	1:B:110:GLU:HG3	2.47	0.44
1:A:79:HIS:HB2	1:A:91:HIS:CE1	2.53	0.43
1:B:180:LEU:HD22	1:B:224:PRO:HB2	2.00	0.43
1:A:243:GLU:CD	1:A:273:HIS:HA	2.43	0.43
1:B:254:ARG:NH2	2:B:403:HOH:O	2.39	0.43
1:A:182:MET:CE	1:A:208:SER:HA	2.48	0.43
1:A:250:VAL:HG11	1:B:254:ARG:HG3	2.01	0.42
1:A:221:LEU:HD12	1:A:221:LEU:HA	1.91	0.42
1:B:180:LEU:HD13	1:B:252:PHE:CG	2.55	0.42
1:B:29:ILE:HD12	1:B:84:LEU:HD13	2.02	0.41
1:A:232:LEU:HB2	1:A:261:LEU:HD21	2.01	0.41
1:B:22:ARG:HB3	1:B:22:ARG:CZ	2.50	0.41
1:B:243:GLU:CD	1:B:273:HIS:HA	2.46	0.41
1:B:212:ARG:HG2	1:B:222:ILE:HD13	2.03	0.40
1:B:154:LEU:HD23	1:B:154:LEU:HA	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	301/312 (96%)	288 (96%)	13 (4%)	0	100	100
1	B	301/312 (96%)	286 (95%)	15 (5%)	0	100	100
All	All	602/624 (96%)	574 (95%)	28 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	224/237 (94%)	208 (93%)	16 (7%)	13	19
1	B	226/237 (95%)	205 (91%)	21 (9%)	8	11
All	All	450/474 (95%)	413 (92%)	37 (8%)	10	14

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	MET
1	A	52	GLU
1	A	55	SER
1	A	73	LYS
1	A	84	LEU
1	A	89	LEU
1	A	154	LEU
1	A	162	LEU
1	A	165	GLN
1	A	190	GLU
1	A	204	LEU
1	A	221	LEU
1	A	235	LEU
1	A	258	LEU
1	A	261	LEU
1	A	282	LEU
1	B	1	MET
1	B	25	GLN
1	B	31	ILE
1	B	52	GLU
1	B	55	SER
1	B	56	THR
1	B	89	LEU
1	B	98	LEU
1	B	128	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	134	LEU
1	B	154	LEU
1	B	162	LEU
1	B	167	LEU
1	B	168	PRO
1	B	221	LEU
1	B	235	LEU
1	B	258	LEU
1	B	261	LEU
1	B	264	ARG
1	B	282	LEU
1	B	299	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	165	GLN
1	A	201	HIS
1	B	201	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	302/312 (96%)	-0.63	0 100 100	24, 38, 67, 101	1 (0%)
1	B	302/312 (96%)	-0.58	0 100 100	17, 40, 66, 100	1 (0%)
All	All	604/624 (96%)	-0.60	0 100 100	17, 39, 67, 101	2 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.