



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:10 AM UTC

PDB ID : 3B2P / pdb_00003b2p
Title : Crystal structure of E. coli Aminopeptidase N in complex with arginine
Authors : Anthony, A.; Leslie, G.; Matthews, B.W.
Deposited on : 2007-10-18
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	891	 88% 9% ..

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8117 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopeptidase N.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	866	Total	C	N	O	S	0	36	0
			7252	4572	1270	1382	28			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	MET	-	expression tag	UNP P04825
A	-19	GLY	-	expression tag	UNP P04825
A	-18	SER	-	expression tag	UNP P04825
A	-17	SER	-	expression tag	UNP P04825
A	-16	HIS	-	expression tag	UNP P04825
A	-15	HIS	-	expression tag	UNP P04825
A	-14	HIS	-	expression tag	UNP P04825
A	-13	HIS	-	expression tag	UNP P04825
A	-12	HIS	-	expression tag	UNP P04825
A	-11	HIS	-	expression tag	UNP P04825
A	-10	SER	-	expression tag	UNP P04825
A	-9	SER	-	expression tag	UNP P04825
A	-8	GLY	-	expression tag	UNP P04825
A	-7	GLU	-	expression tag	UNP P04825
A	-6	ASN	-	expression tag	UNP P04825
A	-5	LEU	-	expression tag	UNP P04825
A	-4	TYR	-	expression tag	UNP P04825
A	-3	PHE	-	expression tag	UNP P04825
A	-2	GLN	-	expression tag	UNP P04825
A	-1	GLY	-	expression tag	UNP P04825
A	0	HIS	-	expression tag	UNP P04825


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Zn		0	0
			1	1			

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	2	Total	Na		0	0
			2	2			

- Molecule 4 is ARGinine (CCD ID: ARG) (formula: C₆H₁₅N₄O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			12	6	4	2		

- Molecule 5 is MALONATE ION (CCD ID: MLI) (formula: C₃H₂O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	1	0
			7	3	4		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	1
			12	6	6		
6	A	1	Total	C	O	0	1
			12	6	6		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		
6	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	765	Total	O	0	0
			765	765		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aminopeptidase N



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	120.20Å 120.20Å 169.66Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.37 – 2.00 44.37 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (44.37-2.00) 99.7 (44.37-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.153 , 0.188 0.155 , 0.189	Depositor DCC
R_{free} test set	2855 reflections (2.98%)	wwPDB-VP
Wilson B-factor (Å ²)	21.4	Xtriage
Anisotropy	0.037	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 48.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8117	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.98% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI, NA, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.86	0/7407	0.84	2/10046 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	693	THR	N-CA-C	5.37	116.81	111.07
1	A	259	ASN	N-CA-C	5.00	116.82	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7252	0	7056	70	0
2	A	1	0	0	0	0
3	A	2	0	0	0	0
4	A	12	0	12	2	0
5	A	7	0	2	0	0
6	A	78	0	104	7	0
7	A	765	0	0	29	1
All	All	8117	0	7174	73	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:423[A]:GLN:NE2	7:A:1723:HOH:O	1.57	1.33
6:A:981:GOL:H11	7:A:1286:HOH:O	1.68	0.94
1:A:107[A]:GLU:HG3	7:A:1636:HOH:O	1.70	0.90
1:A:343:ASN:OD1	1:A:346[A]:ARG:NH1	2.04	0.90
1:A:468[A]:ARG:HA	6:A:978:GOL:H2	1.54	0.86
1:A:442:GLN:HE22	1:A:479:GLN:H	1.26	0.80
1:A:138[B]:LEU:HD12	1:A:183:PRO:HD3	1.63	0.79
1:A:514[B]:GLN:H	1:A:517[B]:GLN:HE22	1.30	0.76
1:A:346[B]:ARG:NH1	7:A:1594:HOH:O	2.21	0.74
1:A:651:GLU:OE1	7:A:1729:HOH:O	2.07	0.72
1:A:476:ALA:HB2	6:A:981:GOL:H2	1.73	0.70
1:A:842[A]:GLU:OE2	1:A:845:ARG:NH1	2.24	0.70
1:A:748[B]:LYS:NZ	7:A:1076:HOH:O	2.08	0.69
1:A:627[A]:GLU:OE2	7:A:1706:HOH:O	2.11	0.69
1:A:346[A]:ARG:NH2	1:A:615:GLU:OE1	2.22	0.68
1:A:279:ARG:NH1	7:A:1616:HOH:O	2.28	0.66
1:A:695:LEU:HD22	1:A:699:LEU:HD22	1.78	0.65
1:A:213[B]:LEU:HG	1:A:227:MET:HE2	1.81	0.63
1:A:107[A]:GLU:OE2	7:A:1405:HOH:O	2.15	0.63
1:A:235:LYS:HE2	1:A:239:GLU:OE2	2.00	0.61
1:A:55:ASN:HD22	1:A:133:ASP:H	1.49	0.61
1:A:753:GLN:HE21	1:A:753:GLN:HA	1.67	0.59
1:A:140[B]:ARG:HD3	7:A:1338:HOH:O	2.03	0.58
1:A:665:GLN:HE22	1:A:674:ASP:HA	1.68	0.58
1:A:71:TRP:O	6:A:973:GOL:H2	2.04	0.57
1:A:138[B]:LEU:HD12	1:A:183:PRO:CD	2.31	0.56
1:A:213[A]:LEU:HD23	1:A:223:ALA:HB1	1.86	0.56
1:A:488[B]:GLU:OE2	1:A:559:ARG:NH2	2.36	0.56
1:A:627[A]:GLU:HG3	7:A:1231:HOH:O	2.06	0.55
1:A:442:GLN:NE2	1:A:479:GLN:H	1.99	0.55
1:A:741:GLN:NE2	1:A:741:GLN:H	2.05	0.54
1:A:468[B]:ARG:HD3	7:A:1196:HOH:O	2.07	0.53
1:A:275:TYR:OH	6:A:972[B]:GOL:H31	2.09	0.53
1:A:825:ARG:HD2	7:A:1753:HOH:O	2.09	0.52
1:A:709:ASN:ND2	1:A:712:ASP:H	2.09	0.51
1:A:260:MET:HB2	7:A:1681:HOH:O	2.10	0.51
1:A:492:ASN:HB3	1:A:526:PHE:CE2	2.47	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:776:MET:O	1:A:782:ILE:HD11	2.13	0.49
1:A:293[B]:ARG:NH2	7:A:1752:HOH:O	2.30	0.49
1:A:28:ASP:OD1	1:A:171[A]:ARG:NH2	2.45	0.48
1:A:350:GLY:C	7:A:1231:HOH:O	2.56	0.47
1:A:195:ASP:HB3	1:A:220:LEU:HD13	1.95	0.47
1:A:517[B]:GLN:HG3	1:A:519:PHE:CE2	2.50	0.46
1:A:168:GLU:HG2	7:A:1418:HOH:O	2.16	0.46
1:A:326:ARG:HG2	1:A:326:ARG:HH11	1.79	0.45
1:A:423[B]:GLN:HG3	7:A:1434:HOH:O	2.16	0.45
1:A:714:LEU:HD22	1:A:748[B]:LYS:HD3	1.99	0.45
4:A:900:ARG:HG2	7:A:1681:HOH:O	2.17	0.45
1:A:65:HIS:HD2	7:A:1048:HOH:O	1.99	0.45
1:A:121:GLU:OE1	4:A:900:ARG:N	2.49	0.45
1:A:709:ASN:ND2	1:A:711:THR:HB	2.32	0.44
1:A:213[B]:LEU:HD22	1:A:252:ILE:HB	1.99	0.44
1:A:264:GLU:HG3	1:A:301:HIS:HB3	1.99	0.44
1:A:138[B]:LEU:CD1	1:A:183:PRO:HD3	2.42	0.43
1:A:423[A]:GLN:HE21	1:A:437:ARG:HH21	1.67	0.43
1:A:473:PRO:CD	7:A:1432:HOH:O	2.65	0.43
1:A:739:TRP:CG	1:A:745:VAL:HG11	2.53	0.43
1:A:339:VAL:HG21	1:A:611:ALA:HB1	2.01	0.43
1:A:349[B]:ARG:O	1:A:627[B]:GLU:HG3	2.18	0.43
1:A:825:ARG:HD2	7:A:1248:HOH:O	2.18	0.43
1:A:249:ILE:HG22	7:A:1358:HOH:O	2.18	0.43
1:A:682:ASN:ND2	1:A:703:GLN:HE22	2.17	0.42
1:A:849:GLU:OE2	7:A:1648:HOH:O	2.22	0.42
1:A:468[B]:ARG:HD3	7:A:1123:HOH:O	2.19	0.42
1:A:840:ARG:HG2	7:A:1617:HOH:O	2.18	0.41
1:A:87:PRO:HG2	1:A:90:PHE:HB3	2.02	0.41
1:A:235:LYS:HE3	7:A:1250:HOH:O	2.19	0.41
1:A:260:MET:HE3	1:A:260:MET:HB3	2.00	0.41
1:A:430:ASN:ND2	7:A:1413:HOH:O	2.53	0.41
1:A:825:ARG:CZ	7:A:1753:HOH:O	2.69	0.41
1:A:235:LYS:O	1:A:239:GLU:HG3	2.21	0.41
1:A:62:VAL:O	6:A:974:GOL:H11	2.21	0.40
6:A:971[A]:GOL:O3	6:A:972[A]:GOL:H12	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:1348:HOH:O	7:A:1470:HOH:O[4_565]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	900/891 (101%)	886 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	777/763 (102%)	760 (98%)	17 (2%)	45	50

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	LEU
1	A	220	LEU
1	A	260	MET
1	A	327	ASP
1	A	356	ASP
1	A	423[A]	GLN
1	A	423[B]	GLN
1	A	477	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	559	ARG
1	A	572	LEU
1	A	589	LEU
1	A	695	LEU
1	A	699	LEU
1	A	741	GLN
1	A	753	GLN
1	A	842[A]	GLU
1	A	842[B]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	HIS
1	A	55	ASN
1	A	65	HIS
1	A	219	ASN
1	A	306	ASN
1	A	430	ASN
1	A	442	GLN
1	A	511	ASN
1	A	661	ASN
1	A	663	ASN
1	A	665	GLN
1	A	682	ASN
1	A	709	ASN
1	A	741	GLN
1	A	753	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 18 ligands modelled in this entry, 3 are monoatomic - leaving 15 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GOL	A	972[B]	-	5,5,5	0.33	0	5,5,5	0.36	0
6	GOL	A	975	-	5,5,5	0.33	0	5,5,5	0.71	0
6	GOL	A	973	-	5,5,5	0.30	0	5,5,5	1.69	2 (40%)
6	GOL	A	980	-	5,5,5	0.46	0	5,5,5	0.56	0
4	ARG	A	900	2	10,11,11	0.84	1 (10%)	9,13,13	1.51	1 (11%)
6	GOL	A	971[B]	-	5,5,5	0.36	0	5,5,5	0.16	0
6	GOL	A	972[A]	-	5,5,5	0.46	0	5,5,5	0.22	0
6	GOL	A	981	-	5,5,5	0.29	0	5,5,5	0.78	0
6	GOL	A	982	-	5,5,5	0.28	0	5,5,5	0.68	0
6	GOL	A	974	-	5,5,5	0.47	0	5,5,5	0.85	0
6	GOL	A	978	-	5,5,5	0.92	1 (20%)	5,5,5	0.59	0
6	GOL	A	970	-	5,5,5	0.70	0	5,5,5	1.87	1 (20%)
6	GOL	A	971[A]	-	5,5,5	0.29	0	5,5,5	0.49	0
5	MLI	A	990	-	6,6,6	2.45	1 (16%)	7,7,7	4.83	2 (28%)
6	GOL	A	976	-	5,5,5	0.35	0	5,5,5	0.91	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	972[B]	-	-	4/4/4/4	-
6	GOL	A	975	-	-	2/4/4/4	-
6	GOL	A	973	-	-	3/4/4/4	-
6	GOL	A	980	-	-	2/4/4/4	-
4	ARG	A	900	2	-	1/11/11/11	-
6	GOL	A	971[B]	-	-	0/4/4/4	-
6	GOL	A	972[A]	-	-	4/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	A	981	-	-	2/4/4/4	-
6	GOL	A	982	-	-	1/4/4/4	-
6	GOL	A	974	-	-	4/4/4/4	-
6	GOL	A	978	-	-	2/4/4/4	-
6	GOL	A	970	-	-	2/4/4/4	-
6	GOL	A	971[A]	-	-	2/4/4/4	-
5	MLI	A	990	-	-	3/4/4/4	-
6	GOL	A	976	-	-	3/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	990	MLI	O6-C2	-5.28	1.04	1.22
4	A	900	ARG	OXT-C	-2.35	1.23	1.30
6	A	978	GOL	O2-C2	-2.02	1.37	1.43

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	990	MLI	O7-C2-O6	-9.64	98.55	123.33
5	A	990	MLI	O6-C2-C1	7.85	144.46	122.11
4	A	900	ARG	OXT-C-O	-3.80	115.46	124.08
6	A	970	GOL	C3-C2-C1	-3.66	98.39	111.80
6	A	973	GOL	O2-C2-C1	2.95	121.38	109.18
6	A	973	GOL	O1-C1-C2	2.01	119.44	110.38

There are no chirality outliers.

All (35) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	970	GOL	O1-C1-C2-C3
6	A	971[A]	GOL	C1-C2-C3-O3
6	A	972[A]	GOL	C1-C2-C3-O3
6	A	972[B]	GOL	C1-C2-C3-O3
6	A	973	GOL	O1-C1-C2-C3
6	A	974	GOL	O1-C1-C2-O2
6	A	974	GOL	O1-C1-C2-C3
6	A	974	GOL	C1-C2-C3-O3
6	A	976	GOL	C1-C2-C3-O3
6	A	978	GOL	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	A	980	GOL	O1-C1-C2-C3
6	A	981	GOL	O1-C1-C2-C3
4	A	900	ARG	NE-CD-CG-CB
6	A	972[B]	GOL	O2-C2-C3-O3
6	A	978	GOL	O1-C1-C2-O2
6	A	972[A]	GOL	O1-C1-C2-C3
6	A	972[B]	GOL	O1-C1-C2-C3
6	A	975	GOL	O1-C1-C2-C3
6	A	976	GOL	O1-C1-C2-C3
6	A	970	GOL	O1-C1-C2-O2
6	A	972[A]	GOL	O1-C1-C2-O2
6	A	972[A]	GOL	O2-C2-C3-O3
6	A	972[B]	GOL	O1-C1-C2-O2
6	A	973	GOL	O1-C1-C2-O2
6	A	974	GOL	O2-C2-C3-O3
6	A	976	GOL	O1-C1-C2-O2
6	A	980	GOL	O1-C1-C2-O2
6	A	981	GOL	O1-C1-C2-O2
6	A	971[A]	GOL	O2-C2-C3-O3
6	A	973	GOL	O2-C2-C3-O3
6	A	982	GOL	O2-C2-C3-O3
5	A	990	MLI	C3-C1-C2-O6
5	A	990	MLI	C2-C1-C3-O9
5	A	990	MLI	C2-C1-C3-O8
6	A	975	GOL	O1-C1-C2-O2

There are no ring outliers.

8 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	972[B]	GOL	1	0
6	A	973	GOL	1	0
4	A	900	ARG	2	0
6	A	972[A]	GOL	1	0
6	A	981	GOL	2	0
6	A	974	GOL	1	0
6	A	978	GOL	1	0
6	A	971[A]	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	866/891 (97%)	-0.64	4 (0%) 87 87	6, 18, 30, 49	36 (4%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	504	HIS	3.5
1	A	169	ASN	2.5
1	A	213[A]	LEU	2.5
1	A	855	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	A	972[A]	6/6	0.69	0.20	31,32,32,33	6
6	GOL	A	972[B]	6/6	0.69	0.20	30,32,33,35	6
6	GOL	A	981	6/6	0.70	0.30	28,28,28,30	6

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	MLI	A	990	7/7	0.81	0.17	36,38,39,41	1
6	GOL	A	971[B]	6/6	0.84	0.19	34,36,36,36	6
6	GOL	A	971[A]	6/6	0.84	0.19	40,41,42,43	6
6	GOL	A	976	6/6	0.85	0.18	45,46,47,48	0
6	GOL	A	982	6/6	0.85	0.32	15,16,19,22	6
6	GOL	A	974	6/6	0.86	0.18	34,43,47,52	0
6	GOL	A	970	6/6	0.87	0.13	22,28,29,29	0
6	GOL	A	978	6/6	0.87	0.24	15,18,19,20	6
6	GOL	A	980	6/6	0.89	0.30	15,18,20,20	6
6	GOL	A	975	6/6	0.91	0.15	40,43,44,44	0
6	GOL	A	973	6/6	0.93	0.10	25,28,31,31	0
4	ARG	A	900	12/12	0.93	0.11	18,25,39,39	0
3	NA	A	952	1/1	0.98	0.06	33,33,33,33	0
3	NA	A	951	1/1	0.98	0.04	23,23,23,23	0
2	ZN	A	950	1/1	1.00	0.01	14,14,14,14	0

6.5 Other polymers [i](#)

There are no such residues in this entry.