



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:55 PM UTC

PDB ID : 5B2R / pdb_00005b2r
Title : Crystal structure of the Streptococcus pyogenes Cas9 VQR variant in complex with sgRNA and target DNA (TGA PAM)
Authors : Hirano, S.; Nishimasu, H.; Ishitani, R.; Nureki, O.
Deposited on : 2016-02-02
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)
RNA backbone	3983	1000 (2.36-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	81	12% 80% 15% 5%
2	B	1372	12% 86% 9% ••
3	C	28	71% 29%
4	D	8	88% 12%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 13799 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called Guide RNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	81	Total	C	N	O	P	0	0	0
			1739	778	319	561	81			

- Molecule 2 is a protein called CRISPR-associated endonuclease Cas9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1319	Total	C	N	O	S	0	1	0
			10547	6734	1826	1967	20			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP Q99ZW2
B	-2	SER	-	expression tag	UNP Q99ZW2
B	-1	GLY	-	expression tag	UNP Q99ZW2
B	0	HIS	-	expression tag	UNP Q99ZW2
B	10	ALA	ASP	engineered mutation	UNP Q99ZW2
B	80	LEU	CYS	engineered mutation	UNP Q99ZW2
B	574	GLU	CYS	engineered mutation	UNP Q99ZW2
B	840	ALA	HIS	engineered mutation	UNP Q99ZW2
B	1135	VAL	ASP	engineered mutation	UNP Q99ZW2
B	1335	GLN	ARG	engineered mutation	UNP Q99ZW2
B	1337	ARG	THR	engineered mutation	UNP Q99ZW2

- Molecule 3 is a DNA chain called Target DNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	28	Total	C	N	O	P	0	0	0
			560	269	100	164	27			

- Molecule 4 is a DNA chain called Non-target DNA, DNA (5'-D(*TP*GP*AP*GP*AP*TP*TP*G)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	8	Total	C	N	O	P	0	0	0
			165	80	31	47	7			


- Molecule 5 is POTASSIUM ION (CCD ID: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	4	Total	K	0	0
			4	4		
5	B	7	Total	K	0	0
			7	7		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	2	Total	Mg	0	0
			2	2		
6	B	1	Total	Mg	0	0
			1	1		

- Molecule 7 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	B	1	Total	C	O	0	0
			4	2	2		
7	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 8 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	199	Total	O	0	0
			199	199		
9	B	492	Total	O	0	0
			492	492		
9	C	50	Total	O	0	0
			50	50		
9	D	13	Total	O	0	0
			13	13		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Guide RNA




• Molecule 2: CRISPR-associated endonuclease Cas9



● Molecule 3: Target DNA



● Molecule 4: Non-target DNA, DNA (5'-D(*TP*GP*AP*GP*AP*TP*TP*G)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	177.76Å 67.74Å 187.59Å 90.00° 111.23° 90.00°	Depositor
Resolution (Å)	47.94 – 2.00 47.94 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.2 (47.94-2.00) 97.3 (47.94-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.03 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.10_2155: ???	Depositor
R, R_{free}	0.201 , 0.230 0.203 , 0.231	Depositor DCC
R_{free} test set	6819 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.586	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13799	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, MG, K, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.25	0/1949	0.46	0/3037
2	B	0.23	0/10738	0.40	0/14474
3	C	0.33	0/626	0.55	0/961
4	D	0.28	0/185	0.53	0/285
All	All	0.24	0/13498	0.42	0/18757

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1739	0	870	5	0
2	B	10547	0	10509	72	0
3	C	560	0	316	7	0
4	D	165	0	93	1	0
5	A	4	0	0	0	0
5	B	7	0	0	0	0
6	A	2	0	0	0	0
6	B	1	0	0	0	0
7	A	4	0	6	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	B	12	0	18	0	0
8	A	4	0	3	0	0
9	A	199	0	0	0	0
9	B	492	0	0	6	0
9	C	50	0	0	2	0
9	D	13	0	0	0	0
All	All	13799	0	11815	82	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (82) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:2:DA:H2'	3:C:3:DA:C8	2.19	0.78
3:C:19:DC:OP2	9:C:101:HOH:O	2.07	0.72
2:B:962:LEU:HD11	2:B:1043:MET:HE3	1.75	0.68
2:B:586:ARG:NH1	9:B:1501:HOH:O	2.26	0.67
2:B:342:GLN:HB2	2:B:383:MET:HE3	1.78	0.65
2:B:558:LYS:HD2	2:B:586:ARG:HH21	1.62	0.64
2:B:241:LEU:O	2:B:244:LEU:N	2.24	0.64
2:B:468:LYS:NZ	9:B:1504:HOH:O	2.32	0.62
2:B:222:LEU:HD23	2:B:234:LYS:HE3	1.83	0.61
2:B:253:LYS:NZ	2:B:261:ASP:OD1	2.30	0.59
2:B:1105:PHE:CD2	2:B:1169:MET:HE2	2.38	0.59
1:A:71:U:H2'	1:A:72:U:C6	2.37	0.58
2:B:809:GLU:OE2	2:B:855:LYS:NZ	2.32	0.58
2:B:790:GLU:HG2	2:B:889:ALA:HA	1.85	0.57
2:B:427:GLU:OE1	2:B:437[B]:ARG:NH1	2.37	0.57
2:B:842:VAL:HB	2:B:847:LEU:HD22	1.87	0.56
2:B:784:ILE:HD12	2:B:806:LEU:HD13	1.88	0.56
2:B:302:LEU:HD12	2:B:414:ILE:HD11	1.88	0.56
3:C:19:DC:H5''	3:C:19:DC:H6	1.71	0.55
2:B:787:GLY:HA3	2:B:891:LEU:HD21	1.88	0.55
2:B:1267:ASP:OD1	2:B:1294:TYR:OH	2.22	0.54
2:B:1179:ILE:HD11	2:B:1192:LYS:HG3	1.90	0.53
2:B:1256:GLN:O	2:B:1260:GLU:HG2	2.08	0.53
2:B:1303:ARG:NH2	2:B:1307:GLU:OE2	2.42	0.53
2:B:999:LYS:HB3	2:B:1073:VAL:HG22	1.91	0.52
2:B:243:ALA:HB1	2:B:248:LEU:HB3	1.92	0.51
2:B:1114:ARG:NH1	4:D:9:DA:OP1	2.42	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:302:LEU:HD22	2:B:302:LEU:H	1.76	0.51
2:B:826:GLN:NE2	2:B:859:ARG:HD2	2.25	0.51
2:B:776:ASN:N	2:B:776:ASN:OD1	2.43	0.50
2:B:817:GLN:CD	2:B:822:MET:HG2	2.36	0.50
2:B:411:PRO:HD2	2:B:414:ILE:HG13	1.94	0.50
2:B:627:GLU:HG3	9:C:102:HOH:O	2.11	0.49
2:B:524:LEU:HD13	2:B:545:LYS:HG2	1.93	0.49
2:B:822:MET:HG3	2:B:856:VAL:HB	1.93	0.49
2:B:249:THR:HG22	2:B:263:LYS:HB2	1.93	0.49
2:B:846:PHE:O	2:B:920:GLN:NE2	2.44	0.48
2:B:1339:THR:O	2:B:1342:VAL:HG22	2.14	0.48
2:B:195:LEU:HD21	2:B:286:TYR:HA	1.94	0.48
3:C:1:DC:H2'	3:C:2:DA:C8	2.48	0.48
2:B:1177:ASN:ND2	2:B:1180:ASP:OD2	2.48	0.47
2:B:901:THR:O	2:B:904:GLU:HG2	2.16	0.46
2:B:185:PHE:HD2	2:B:296:LEU:HD11	1.80	0.46
2:B:1042:ILE:HG23	2:B:1043:MET:HE2	1.97	0.46
2:B:1215:ALA:HB2	2:B:1221:GLN:HG3	1.98	0.46
2:B:253:LYS:HG3	2:B:261:ASP:HA	1.97	0.46
2:B:302:LEU:HG	2:B:306:LEU:HD12	1.97	0.46
2:B:972:PHE:HE1	2:B:1084:ARG:HG2	1.81	0.45
2:B:1085:LYS:O	2:B:1089:MET:HG3	2.17	0.45
1:A:14:G:OP2	2:B:63:ARG:NH1	2.39	0.45
2:B:756:PRO:O	2:B:953:VAL:HG22	2.17	0.45
2:B:1169:MET:HE3	9:B:1966:HOH:O	2.16	0.45
2:B:387:GLU:O	2:B:391:VAL:HG23	2.16	0.45
2:B:1004:LEU:HD11	2:B:1042:ILE:HD11	1.98	0.45
1:A:45:U:H5'	2:B:402:GLN:HE21	1.82	0.44
2:B:788:ILE:HG23	2:B:793:SER:HB3	2.00	0.44
2:B:511:HIS:HD2	9:B:1610:HOH:O	2.00	0.44
2:B:336:LYS:NZ	9:B:1511:HOH:O	2.41	0.44
2:B:1194:LEU:HD13	2:B:1365:LEU:HD22	2.00	0.44
2:B:248:LEU:HD12	2:B:249:THR:H	1.83	0.43
2:B:288:ASP:HA	2:B:291:LEU:HB3	2.00	0.43
2:B:870:VAL:HG11	2:B:902:LYS:HB3	2.01	0.43
1:A:46:A:H2'	1:A:47:A:C8	2.54	0.43
2:B:1041:ASN:OD1	2:B:1044:ASN:ND2	2.49	0.43
2:B:187:GLN:O	2:B:191:THR:N	2.50	0.43
2:B:390:LEU:HD23	2:B:390:LEU:HA	1.87	0.43
2:B:841:ILE:HD13	2:B:900:LEU:HG	2.01	0.43
2:B:1231:LYS:HE3	2:B:1232:TYR:CZ	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:A:H3'	1:A:75:A:H8	1.84	0.42
2:B:195:LEU:HD23	2:B:289:LEU:HB2	2.01	0.42
2:B:302:LEU:H	2:B:302:LEU:CD2	2.31	0.42
2:B:1063:ILE:HG23	2:B:1072:ILE:HG13	2.02	0.42
2:B:1119:LEU:HB3	2:B:1128:PRO:HB3	2.02	0.42
3:C:20:DC:H2''	3:C:21:DT:H5'	2.02	0.42
3:C:16:DG:H2'	3:C:17:DC:C6	2.54	0.42
2:B:381:GLU:HG2	2:B:390:LEU:HD11	2.02	0.41
2:B:15:SER:HA	2:B:51:LEU:O	2.21	0.41
2:B:302:LEU:HG	2:B:306:LEU:CD1	2.50	0.41
3:C:2:DA:H2''	3:C:3:DA:O5'	2.20	0.41
2:B:812:TYR:CZ	2:B:816:LEU:HD11	2.55	0.41
2:B:338:LEU:C	2:B:383:MET:HE1	2.46	0.40
2:B:104:SER:HA	9:B:1774:HOH:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	B	1308/1372 (95%)	1269 (97%)	38 (3%)	1 (0%)	48 46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	1216	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	1121/1227 (91%)	1090 (97%)	31 (3%)	38 41

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	67	THR
2	B	101	LEU
2	B	123	VAL
2	B	216	LEU
2	B	219	SER
2	B	253	LYS
2	B	263	LYS
2	B	264	LEU
2	B	279	LEU
2	B	301	LEU
2	B	302	LEU
2	B	313	THR
2	B	388	GLU
2	B	419	LEU
2	B	490	SER
2	B	524	LEU
2	B	571	LYS
2	B	638	THR
2	B	642	LEU
2	B	718	ASP
2	B	776	ASN
2	B	801	VAL
2	B	833	LEU
2	B	847	LEU
2	B	853	ASP
2	B	870	VAL
2	B	917	ILE
2	B	947	ASP
2	B	1119	LEU
2	B	1154	SER
2	B	1250	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	402	GLN
2	B	412	HIS
2	B	511	HIS
2	B	726	ASN
2	B	739	GLN
2	B	926	GLN
2	B	982	HIS
2	B	1177	ASN
2	B	1261	GLN
2	B	1262	HIS
2	B	1264	HIS
2	B	1317	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	80/81 (98%)	12 (15%)	1 (1%)

All (12) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	17	U
1	A	28	A
1	A	40	C
1	A	51	A
1	A	56	U
1	A	58	G
1	A	59	U
1	A	68	A
1	A	71	U
1	A	72	U
1	A	74	A
1	A	75	A

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
1	A	71	U

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 14 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	EDO	B	1411	-	3,3,3	0.42	0	2,2,2	0.38	0
7	EDO	A	106	-	3,3,3	0.44	0	2,2,2	0.29	0
8	ACT	A	107	-	3,3,3	0.80	0	3,3,3	1.31	0
7	EDO	B	1409	-	3,3,3	0.53	0	2,2,2	0.21	0
7	EDO	B	1410	-	3,3,3	0.50	0	2,2,2	0.22	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	B	1411	-	-	0/1/1/1	-
7	EDO	B	1410	-	-	0/1/1/1	-
7	EDO	A	106	-	-	0/1/1/1	-
7	EDO	B	1409	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	81/81 (100%)	-0.69	0 100 100	26, 41, 122, 188	0
2	B	1319/1372 (96%)	0.68	161 (12%) 8 7	24, 52, 95, 123	1 (0%)
3	C	28/28 (100%)	-0.22	0 100 100	34, 48, 74, 84	0
4	D	8/8 (100%)	0.17	0 100 100	37, 52, 98, 119	0
All	All	1436/1489 (96%)	0.58	161 (11%) 10 9	24, 51, 96, 188	1 (0%)

All (161) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	299	ALA	8.0
2	B	246	LEU	7.5
2	B	264	LEU	7.3
2	B	296	LEU	7.1
2	B	292	ALA	6.7
2	B	245	SER	6.2
2	B	275	LEU	6.2
2	B	244	LEU	6.0
2	B	189	VAL	5.8
2	B	287	ALA	5.7
2	B	186	ILE	5.6
2	B	247	GLY	5.6
2	B	188	LEU	5.5
2	B	271	TYR	5.5
2	B	194	GLN	5.5
2	B	243	ALA	5.3
2	B	262	ALA	5.2
2	B	297	SER	5.2
2	B	250	PRO	5.1
2	B	286	TYR	5.1
2	B	187	GLN	5.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	301	LEU	4.9
2	B	178	ASN	4.8
2	B	300	ILE	4.7
2	B	284	ASP	4.7
2	B	185	PHE	4.6
2	B	309	ASN	4.6
2	B	291	LEU	4.5
2	B	911	LEU	4.5
2	B	248	LEU	4.4
2	B	184	LEU	4.4
2	B	176	PRO	4.4
2	B	1299	ASP	4.2
2	B	266	LEU	4.2
2	B	289	LEU	4.2
2	B	295	ASN	4.2
2	B	1062	LEU	4.1
2	B	195	LEU	4.1
2	B	729	GLY	4.0
2	B	290	PHE	4.0
2	B	279	LEU	4.0
2	B	249	THR	4.0
2	B	193	ASN	3.9
2	B	215	ARG	3.9
2	B	273	ASP	3.8
2	B	201	ILE	3.8
2	B	803	ASN	3.8
2	B	1242	TYR	3.7
2	B	242	ILE	3.7
2	B	191	THR	3.7
2	B	270	THR	3.6
2	B	285	GLN	3.6
2	B	312	ILE	3.6
2	B	1055	GLY	3.6
2	B	216	LEU	3.5
2	B	214	ALA	3.5
2	B	1037	PHE	3.4
2	B	192	TYR	3.4
2	B	1051	THR	3.4
2	B	200	PRO	3.4
2	B	174	LEU	3.4
2	B	1052	LEU	3.4
2	B	199	ASN	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	804	THR	3.3
2	B	765	ARG	3.3
2	B	1243	GLU	3.2
2	B	890	LYS	3.2
2	B	1057	ILE	3.2
2	B	1257	LEU	3.2
2	B	190	GLN	3.2
2	B	252	PHE	3.1
2	B	1366	GLY	3.1
2	B	258	LEU	3.1
2	B	731	PRO	3.1
2	B	390	LEU	3.1
2	B	341	GLN	3.1
2	B	259	ALA	3.0
2	B	219	SER	3.0
2	B	1011	GLY	3.0
2	B	269	ASP	3.0
2	B	268	LYS	3.0
2	B	1054	ASN	3.0
2	B	1072	ILE	3.0
2	B	1241	HIS	3.0
2	B	260	GLU	2.9
2	B	278	LEU	2.9
2	B	870	VAL	2.9
2	B	251	ASN	2.9
2	B	1264	HIS	2.8
2	B	728	ALA	2.8
2	B	730	SER	2.8
2	B	868	ASP	2.8
2	B	387	GLU	2.8
2	B	302	LEU	2.8
2	B	847	LEU	2.8
2	B	274	ASP	2.8
2	B	796	LEU	2.8
2	B	1250	GLU	2.7
2	B	824	VAL	2.7
2	B	1031	LYS	2.7
2	B	1050	ILE	2.7
2	B	782	LYS	2.7
2	B	310	THR	2.6
2	B	293	ALA	2.6
2	B	1115	ASN	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	280	ALA	2.5
2	B	1116	SER	2.5
2	B	211	ILE	2.5
2	B	795	ILE	2.5
2	B	1070	GLY	2.5
2	B	288	ASP	2.5
2	B	386	THR	2.5
2	B	778	ARG	2.5
2	B	208	ALA	2.4
2	B	283	GLY	2.4
2	B	1085	LYS	2.4
2	B	1056	GLU	2.4
2	B	1258	PHE	2.4
2	B	476	TRP	2.4
2	B	263	LYS	2.4
2	B	210	ALA	2.4
2	B	1306	ALA	2.4
2	B	213	SER	2.4
2	B	272	ASP	2.3
2	B	218	LYS	2.3
2	B	1249	PRO	2.3
2	B	343	LEU	2.3
2	B	1251	ASP	2.3
2	B	852	ILE	2.3
2	B	230	PRO	2.3
2	B	265	GLN	2.3
2	B	261	ASP	2.3
2	B	1260	GLU	2.2
2	B	753	ARG	2.2
2	B	783	ARG	2.2
2	B	866	LYS	2.2
2	B	921	LEU	2.2
2	B	281	GLN	2.2
2	B	276	ASP	2.2
2	B	435	ASP	2.2
2	B	1293	ALA	2.2
2	B	222	LEU	2.2
2	B	1069	THR	2.2
2	B	786	GLU	2.1
2	B	1034	ALA	2.1
2	B	209	LYS	2.1
2	B	177	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	205	GLY	2.1
2	B	867	SER	2.1
2	B	175	ASN	2.1
2	B	235	ASN	2.1
2	B	836	TYR	2.1
2	B	779	GLU	2.0
2	B	812	TYR	2.0
2	B	815	TYR	2.0
2	B	1036	TYR	2.0
2	B	916	PHE	2.0
2	B	889	ALA	2.0
2	B	225	LEU	2.0
2	B	791	LEU	2.0
2	B	1058	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	K	B	1407	1/1	0.38	0.35	210,210,210,210	0
5	K	B	1406	1/1	0.76	0.29	118,118,118,118	0
5	K	A	108	1/1	0.82	0.24	114,114,114,114	0
5	K	B	1405	1/1	0.83	0.13	83,83,83,83	0
5	K	B	1404	1/1	0.84	0.15	94,94,94,94	0
8	ACT	A	107	4/4	0.88	0.16	49,52,53,55	0
7	EDO	B	1410	4/4	0.90	0.15	45,47,47,48	0
7	EDO	A	106	4/4	0.94	0.09	33,34,34,37	0
6	MG	B	1408	1/1	0.96	0.11	46,46,46,46	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	MG	A	105	1/1	0.97	0.06	46,46,46,46	0
7	EDO	B	1411	4/4	0.97	0.09	34,41,45,48	0
7	EDO	B	1409	4/4	0.97	0.08	33,34,35,35	0
5	K	B	1403	1/1	0.98	0.06	58,58,58,58	0
5	K	B	1402	1/1	0.98	0.04	48,48,48,48	0
5	K	A	102	1/1	0.99	0.11	34,34,34,34	0
5	K	A	103	1/1	0.99	0.08	34,34,34,34	0
6	MG	A	104	1/1	0.99	0.03	42,42,42,42	0
5	K	A	101	1/1	0.99	0.03	30,30,30,30	0
5	K	B	1401	1/1	0.99	0.10	39,39,39,39	0

6.5 Other polymers [i](#)

There are no such residues in this entry.