



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 08:02 PM EDT

PDB ID : 7B2O / pdb_00007b2o
Title : Crystal structure of Chlamydomonas reinhardtii chloroplastic sedoheptulose-1,7-bisphosphatase
Authors : Le Moigne, T.; Lemaire, S.D.; Henri, J.
Deposited on : 2020-11-27
Resolution : 3.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtrriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	86% 8% 6%
1	B	331	84% 9% 7%
1	C	331	85% 9% 7%
1	D	331	82% 11% 6%
1	E	331	80% 13% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	331	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into four segments: a small red segment at the beginning labeled '2%', a large green segment labeled '78%', a yellow segment labeled '15%', and a small grey segment at the end labeled '7%'.</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 14194 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called FB Pase domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	311	2372	1509	393	449	21	0	0	0
1	B	309	2363	1503	391	448	21	0	1	0
1	C	309	2359	1501	390	447	21	0	0	0
1	D	310	2362	1504	391	446	21	0	0	0
1	E	311	2371	1509	392	449	21	0	0	0
1	F	309	2363	1504	392	446	21	0	1	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	61	MET	-	initiating methionine	UNP A0A2K3DY10
A	62	HIS	-	expression tag	UNP A0A2K3DY10
A	63	HIS	-	expression tag	UNP A0A2K3DY10
A	64	HIS	-	expression tag	UNP A0A2K3DY10
A	65	HIS	-	expression tag	UNP A0A2K3DY10
A	66	HIS	-	expression tag	UNP A0A2K3DY10
A	67	HIS	-	expression tag	UNP A0A2K3DY10
A	68	HIS	-	expression tag	UNP A0A2K3DY10
A	69	MET	-	expression tag	UNP A0A2K3DY10
B	64	MET	-	initiating methionine	UNP A0A2K3DY10
B	65	HIS	-	expression tag	UNP A0A2K3DY10
B	66	HIS	-	expression tag	UNP A0A2K3DY10
B	67	HIS	-	expression tag	UNP A0A2K3DY10
B	68	HIS	-	expression tag	UNP A0A2K3DY10
B	69	HIS	-	expression tag	UNP A0A2K3DY10
B	70	HIS	-	expression tag	UNP A0A2K3DY10
B	71	HIS	-	expression tag	UNP A0A2K3DY10

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	72	MET	-	expression tag	UNP A0A2K3DY10
C	64	MET	-	initiating methionine	UNP A0A2K3DY10
C	65	HIS	-	expression tag	UNP A0A2K3DY10
C	66	HIS	-	expression tag	UNP A0A2K3DY10
C	67	HIS	-	expression tag	UNP A0A2K3DY10
C	68	HIS	-	expression tag	UNP A0A2K3DY10
C	69	HIS	-	expression tag	UNP A0A2K3DY10
C	70	HIS	-	expression tag	UNP A0A2K3DY10
C	71	HIS	-	expression tag	UNP A0A2K3DY10
C	72	MET	-	expression tag	UNP A0A2K3DY10
D	64	MET	-	initiating methionine	UNP A0A2K3DY10
D	65	HIS	-	expression tag	UNP A0A2K3DY10
D	66	HIS	-	expression tag	UNP A0A2K3DY10
D	67	HIS	-	expression tag	UNP A0A2K3DY10
D	68	HIS	-	expression tag	UNP A0A2K3DY10
D	69	HIS	-	expression tag	UNP A0A2K3DY10
D	70	HIS	-	expression tag	UNP A0A2K3DY10
D	71	HIS	-	expression tag	UNP A0A2K3DY10
D	72	MET	-	expression tag	UNP A0A2K3DY10
E	64	MET	-	initiating methionine	UNP A0A2K3DY10
E	65	HIS	-	expression tag	UNP A0A2K3DY10
E	66	HIS	-	expression tag	UNP A0A2K3DY10
E	67	HIS	-	expression tag	UNP A0A2K3DY10
E	68	HIS	-	expression tag	UNP A0A2K3DY10
E	69	HIS	-	expression tag	UNP A0A2K3DY10
E	70	HIS	-	expression tag	UNP A0A2K3DY10
E	71	HIS	-	expression tag	UNP A0A2K3DY10
E	72	MET	-	expression tag	UNP A0A2K3DY10
F	64	MET	-	initiating methionine	UNP A0A2K3DY10
F	65	HIS	-	expression tag	UNP A0A2K3DY10
F	66	HIS	-	expression tag	UNP A0A2K3DY10
F	67	HIS	-	expression tag	UNP A0A2K3DY10
F	68	HIS	-	expression tag	UNP A0A2K3DY10
F	69	HIS	-	expression tag	UNP A0A2K3DY10
F	70	HIS	-	expression tag	UNP A0A2K3DY10
F	71	HIS	-	expression tag	UNP A0A2K3DY10
F	72	MET	-	expression tag	UNP A0A2K3DY10

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O 1 1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FBPase domain-containing protein




- Molecule 1: FBPase domain-containing protein




- Molecule 1: FBPase domain-containing protein




- Molecule 1: FBPase domain-containing protein






- Molecule 1: FBPase domain-containing protein

Chain E: 80% 13% 6%



- Molecule 1: FBPase domain-containing protein

Chain F: 2% 78% 15% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	178.22Å 183.65Å 75.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.47 – 3.09 47.47 – 3.10	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.47-3.09) 99.3 (47.47-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.69 (at 3.12Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.193 , 0.238 0.194 , 0.236	Depositor DCC
R_{free} test set	1989 reflections (4.33%)	wwPDB-VP
Wilson B-factor (Å ²)	64.7	Xtrriage
Anisotropy	0.826	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 37.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.010 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14194	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.24% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2421	0.45	0/3275
1	B	0.24	0/2412	0.47	0/3264
1	C	0.21	0/2408	0.44	0/3259
1	D	0.21	0/2411	0.43	0/3263
1	E	0.21	0/2420	0.46	0/3275
1	F	0.20	0/2412	0.47	1/3263 (0.0%)
All	All	0.22	0/14484	0.45	1/19599 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	122	GLY	N-CA-C	6.19	117.65	111.21

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2372	0	2353	19	0
1	B	2363	0	2339	21	0
1	C	2359	0	2337	19	0
1	D	2362	0	2345	31	0
1	E	2371	0	2351	33	0
1	F	2363	0	2346	31	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	1	0	0	0	0
2	B	3	0	0	0	0
All	All	14194	0	14071	140	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 140 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:329:VAL:HG13	1:E:366:ILE:HD12	1.53	0.89
1:E:325:ILE:HA	1:E:329:VAL:HB	1.56	0.85
1:F:325:ILE:HA	1:F:329:VAL:HG22	1.64	0.80
1:D:366:ILE:HD12	1:D:366:ILE:O	1.83	0.77
1:F:325:ILE:HD12	1:F:326:LEU:N	1.98	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	309/331 (93%)	299 (97%)	10 (3%)	0	100	100
1	B	308/331 (93%)	292 (95%)	16 (5%)	0	100	100
1	C	307/331 (93%)	288 (94%)	19 (6%)	0	100	100
1	D	308/331 (93%)	292 (95%)	16 (5%)	0	100	100
1	E	309/331 (93%)	283 (92%)	23 (7%)	3 (1%)	12	41
1	F	308/331 (93%)	288 (94%)	19 (6%)	1 (0%)	36	67
All	All	1849/1986 (93%)	1742 (94%)	103 (6%)	4 (0%)	43	73

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	325	ILE
1	E	167	ASP
1	E	236	CYS
1	F	255	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	254/270 (94%)	254 (100%)	0	100	100
1	B	253/270 (94%)	251 (99%)	2 (1%)	73	81
1	C	253/270 (94%)	253 (100%)	0	100	100
1	D	253/270 (94%)	253 (100%)	0	100	100
1	E	254/270 (94%)	254 (100%)	0	100	100
1	F	253/270 (94%)	253 (100%)	0	100	100
All	All	1520/1620 (94%)	1518 (100%)	2 (0%)	88	90

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	258[A]	GLU
1	B	258[B]	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	113	HIS
1	E	212	GLN
1	D	266	ASN
1	D	113	HIS
1	E	99	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	311/331 (93%)	-0.46	0 100 100	33, 52, 76, 97	0
1	B	309/331 (93%)	-0.39	1 (0%) 90 80	36, 52, 79, 106	1 (0%)
1	C	309/331 (93%)	-0.20	1 (0%) 90 80	44, 63, 88, 102	0
1	D	310/331 (93%)	-0.20	1 (0%) 90 80	44, 65, 92, 109	0
1	E	311/331 (93%)	0.05	1 (0%) 90 80	47, 72, 106, 120	0
1	F	309/331 (93%)	0.18	5 (1%) 70 49	40, 84, 114, 123	2 (0%)
All	All	1859/1986 (93%)	-0.17	9 (0%) 87 73	33, 64, 102, 123	3 (0%)

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	260[A]	LYS	8.0
1	B	258[A]	GLU	4.0
1	F	318	THR	3.0
1	D	271	PHE	2.8
1	C	271	PHE	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.