



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:45 AM UTC

PDB ID : 1B3B / pdb_00001b3b
Title : THERMOTOGA MARITIMA GLUTAMATE DEHYDROGENASE MUTANT N97D, G376K
Authors : Knapp, S.; Lebbink, J.H.G.; Van Der Oost, J.; Devos, W.M.; Rice, D.; Ladenstein, R.
Deposited on : 1998-12-07
Resolution : 3.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	415	
1	B	415	
1	C	415	
1	D	415	
1	E	415	
1	F	415	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 19008 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (GLUTAMATE DEHYDROGENASE).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	B	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	C	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	D	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	E	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			
1	F	409	Total	C	N	O	S	0	0	0
			3168	2011	552	592	13			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	97	ASP	ASN	engineered mutation	UNP P96110
A	376	LYS	GLY	engineered mutation	UNP P96110
B	97	ASP	ASN	engineered mutation	UNP P96110
B	376	LYS	GLY	engineered mutation	UNP P96110
C	97	ASP	ASN	engineered mutation	UNP P96110
C	376	LYS	GLY	engineered mutation	UNP P96110
D	97	ASP	ASN	engineered mutation	UNP P96110
D	376	LYS	GLY	engineered mutation	UNP P96110
E	97	ASP	ASN	engineered mutation	UNP P96110
E	376	LYS	GLY	engineered mutation	UNP P96110
F	97	ASP	ASN	engineered mutation	UNP P96110
F	376	LYS	GLY	engineered mutation	UNP P96110

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)




• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)






• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)

Chain C: 53% 42% . .




• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)

Chain D: 56% 39% . .




• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)

Chain E: 50% 43% 6% .



• Molecule 1: PROTEIN (GLUTAMATE DEHYDROGENASE)

Chain F: 51% 42% 5% •



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	145.10 Å 145.10 Å 272.50 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 3.10	Depositor
% Data completeness (in resolution range)	94.0 (8.00-3.10)	Depositor
R_{merge}	0.10	Depositor
R_{sym}	0.05	Depositor
Refinement program	X-PLOR 3.8	Depositor
R, R_{free}	0.225 , 0.298	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	19008	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.40	0/3227	0.82	2/4367 (0.0%)
1	B	0.39	0/3227	0.82	3/4367 (0.1%)
1	C	0.38	0/3227	0.82	1/4367 (0.0%)
1	D	0.37	0/3227	0.81	2/4367 (0.0%)
1	E	0.41	0/3227	0.83	3/4367 (0.1%)
1	F	0.40	0/3227	0.80	3/4367 (0.1%)
All	All	0.39	0/19362	0.82	14/26202 (0.1%)

There are no bond length outliers.

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	340	ALA	N-CA-C	6.18	117.70	110.97
1	B	335	VAL	CA-C-N	5.82	126.11	119.83
1	B	335	VAL	C-N-CA	5.82	126.11	119.83
1	F	199	ALA	N-CA-C	-5.65	105.02	111.07
1	F	340	ALA	N-CA-C	5.61	117.08	110.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3168	0	3192	124	18
1	B	3168	0	3192	121	20

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3168	0	3192	118	2
1	D	3168	0	3192	126	2
1	E	3168	0	3192	143	18
1	F	3168	0	3192	128	20
All	All	19008	0	19152	747	40

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 747 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:248:ASN:HD21	1:E:272:LYS:HE2	1.22	1.03
1:F:248:ASN:HD21	1:F:272:LYS:HE2	1.32	0.95
1:A:248:ASN:HD21	1:A:272:LYS:HE2	1.38	0.88
1:E:224:ALA:O	1:E:228:ILE:HG13	1.74	0.88
1:C:354:GLN:HG2	1:C:360:PHE:HA	1.55	0.87

The worst 5 of 40 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:299:HIS:CG	1:E:304:GLU:OE1[6_655]	0.79	1.41
1:B:279:GLU:CG	1:F:330:ARG:O[6_665]	1.00	1.20
1:A:279:GLU:OE1	1:E:330:ARG:NE[6_655]	1.11	1.09
1:B:296:GLY:O	1:F:329:ARG:NH2[6_665]	1.20	1.00
1:A:299:HIS:ND1	1:E:304:GLU:OE1[6_655]	1.28	0.92

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	407/415 (98%)	365 (90%)	39 (10%)	3 (1%)	18	49
1	B	407/415 (98%)	366 (90%)	37 (9%)	4 (1%)	12	41
1	C	407/415 (98%)	369 (91%)	35 (9%)	3 (1%)	18	49
1	D	407/415 (98%)	370 (91%)	33 (8%)	4 (1%)	12	41
1	E	407/415 (98%)	373 (92%)	32 (8%)	2 (0%)	24	57
1	F	407/415 (98%)	366 (90%)	37 (9%)	4 (1%)	12	41
All	All	2442/2490 (98%)	2209 (90%)	213 (9%)	20 (1%)	16	47

5 of 20 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	178	LEU
1	A	178	LEU
1	A	303	ALA
1	B	178	LEU
1	B	303	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/339 (98%)	298 (90%)	35 (10%)	6	26
1	B	333/339 (98%)	301 (90%)	32 (10%)	8	30
1	C	333/339 (98%)	298 (90%)	35 (10%)	6	26
1	D	333/339 (98%)	297 (89%)	36 (11%)	6	25
1	E	333/339 (98%)	298 (90%)	35 (10%)	6	26
1	F	333/339 (98%)	290 (87%)	43 (13%)	4	18
All	All	1998/2034 (98%)	1782 (89%)	216 (11%)	6	25

5 of 216 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	217	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	152	ILE
1	F	267	VAL
1	D	261	LYS
1	D	337	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	61	ASN
1	F	357	GLN
1	D	365	GLN
1	E	389	ASN
1	D	278	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.