



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 12:12 PM UTC

PDB ID : 2B3X / pdb_00002b3x
Title : Structure of an orthorhombic crystal form of human cytosolic aconitase (IRP1)
Authors : Dupuy, J.; Fontecilla-Camps, J.C.; Volbeda, A.
Deposited on : 2005-09-22
Resolution : 2.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1091 (2.54-2.54)
Clashscore	190562	1120 (2.54-2.54)
Ramachandran outliers	187476	1106 (2.54-2.54)
Sidechain outliers	187428	1106 (2.54-2.54)
RSRZ outliers	180081	1091 (2.54-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	888	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7109 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Iron-responsive element binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	888	Total	C	N	O	S	47	0	0
			6928	4431	1183	1285	29			

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	168	Total	O	0	0
			168	168		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Iron-responsive element binding protein 1



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	75.35Å 103.31Å 225.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	113.23 – 2.54 113.23 – 2.54	Depositor EDS
% Data completeness (in resolution range)	96.9 (113.23-2.54) 74.4 (113.23-2.54)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.32 (at 2.54Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.180 , 0.225 (Not available) , 0.220	Depositor DCC
R_{free} test set	1435 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	34.9	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7109	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, EDO, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.76	10/7089 (0.1%)	0.94	15/9626 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	536	ARG	NE-CZ	-24.07	1.06	1.33
1	A	579	ALA	C-N	-19.03	1.07	1.33
1	A	541	ARG	CG-CD	12.48	1.89	1.52
1	A	825	ASP	CB-CG	-10.62	1.25	1.52
1	A	579	ALA	N-CA	-7.86	1.36	1.46
1	A	326	ASP	CB-CG	-7.84	1.32	1.52
1	A	757	GLU	CB-CG	-7.38	1.30	1.52
1	A	541	ARG	CD-NE	6.17	1.54	1.46
1	A	541	ARG	NE-CZ	5.49	1.39	1.33
1	A	527	VAL	CA-CB	5.42	1.59	1.53

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	536	ARG	NE-CZ-NH2	-12.01	108.39	119.20
1	A	536	ARG	NE-CZ-NH1	9.46	130.96	121.50
1	A	225	GLU	CB-CG-CD	8.44	126.95	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	541	ARG	N-CA-C	6.80	121.32	112.89
1	A	801	GLU	CB-CG-CD	6.60	123.82	112.60
1	A	536	ARG	CD-NE-CZ	-6.39	115.45	124.40
1	A	140	GLN	CB-CG-CD	-6.31	101.87	112.60
1	A	582	GLN	N-CA-C	6.05	116.95	108.00
1	A	841	GLU	CB-CG-CD	5.74	122.36	112.60
1	A	541	ARG	CG-CD-NE	5.47	124.03	112.00
1	A	571	GLU	CB-CG-CD	-5.42	103.39	112.60
1	A	326	ASP	CA-CB-CG	5.12	117.72	112.60
1	A	825	ASP	CA-CB-CG	5.05	117.65	112.60
1	A	201	LEU	N-CA-C	5.04	116.51	108.79
1	A	757	GLU	CA-CB-CG	5.02	124.14	114.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	579	ALA	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6928	0	6919	95	0
2	A	1	0	0	0	0
3	A	8	0	0	0	0
4	A	4	0	6	1	0
5	A	168	0	0	15	0
All	All	7109	0	6925	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:345:ASP:H	1:A:351:GLN:HE22	1.13	0.96
1:A:396:LYS:HE3	1:A:396:LYS:H	1.27	0.96
1:A:519:GLU:O	1:A:523:GLN:HG2	1.69	0.93
1:A:82:ARG:HH22	1:A:181:ASN:HD21	0.99	0.91
1:A:82:ARG:HH22	1:A:181:ASN:ND2	1.68	0.90
1:A:287:GLN:HB3	5:A:1148:HOH:O	1.80	0.82
1:A:743:HIS:HE1	5:A:1151:HOH:O	1.61	0.82
1:A:420:THR:CG2	5:A:1170:HOH:O	2.28	0.80
1:A:420:THR:HG22	5:A:1170:HOH:O	1.83	0.79
1:A:222:GLY:HA3	1:A:807:ASN:HD21	1.53	0.72
1:A:396:LYS:HE3	1:A:396:LYS:N	2.06	0.69
1:A:580:LYS:HD2	1:A:581:GLY:N	2.07	0.69
1:A:699:ARG:CG	1:A:699:ARG:HH21	2.06	0.69
1:A:201:LEU:HD21	1:A:211:ILE:HG21	1.74	0.68
1:A:396:LYS:H	1:A:396:LYS:CE	2.04	0.65
1:A:420:THR:HG21	5:A:1170:HOH:O	1.96	0.64
1:A:345:ASP:H	1:A:351:GLN:NE2	1.91	0.63
1:A:743:HIS:CE1	5:A:1151:HOH:O	2.43	0.62
1:A:210:MET:HE1	1:A:437:CYS:O	2.01	0.60
1:A:93:ALA:HA	1:A:227:GLU:OE2	2.00	0.60
1:A:82:ARG:HD3	1:A:199:ASP:OD1	2.02	0.60
1:A:345:ASP:N	1:A:351:GLN:HE22	1.93	0.59
1:A:115:ASN:ND2	1:A:165:HIS:H	2.01	0.59
1:A:699:ARG:HH21	1:A:699:ARG:HG3	1.68	0.58
1:A:248:MET:HE1	1:A:360:GLU:HB3	1.85	0.57
1:A:179:GLN:OE1	1:A:474:GLY:HA3	2.03	0.56
1:A:104:VAL:HG21	1:A:114:ILE:CD1	2.37	0.54
1:A:201:LEU:O	1:A:218:GLY:HA2	2.06	0.54
1:A:253:PRO:HA	1:A:593:ARG:HH21	1.72	0.54
1:A:130:VAL:HG22	1:A:504:MET:HE2	1.89	0.54
1:A:253:PRO:HA	1:A:593:ARG:HE	1.74	0.53
1:A:689:ASN:HD22	1:A:689:ASN:H	1.56	0.53
1:A:324:GLY:O	1:A:610:LYS:HE3	2.10	0.52
1:A:385:MET:HE1	1:A:550:TYR:CE2	2.45	0.51
1:A:743:HIS:HD2	1:A:746:SER:OG	1.93	0.50
1:A:446:MET:HE1	1:A:471:LEU:HD21	1.93	0.50
1:A:435:THR:O	1:A:436:SER:HB3	2.11	0.50
1:A:82:ARG:NH2	1:A:181:ASN:ND2	2.50	0.49
1:A:593:ARG:O	1:A:597:GLN:HB2	2.12	0.49
1:A:227:GLU:OE1	1:A:880:LEU:HD11	2.13	0.49
1:A:255:VAL:O	1:A:593:ARG:HD3	2.13	0.48
1:A:468:LYS:HE3	5:A:1144:HOH:O	2.11	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:435:THR:HG22	5:A:1005:HOH:O	2.13	0.48
1:A:284:GLY:O	1:A:287:GLN:HG2	2.13	0.48
1:A:616:ILE:HD12	5:A:1142:HOH:O	2.13	0.48
1:A:776:ALA:HA	1:A:801:GLU:HG3	1.96	0.48
1:A:838:ILE:HD11	1:A:852:LYS:HD3	1.96	0.48
1:A:519:GLU:O	1:A:523:GLN:CG	2.53	0.48
1:A:8:LEU:HD11	1:A:27:LEU:HD13	1.96	0.47
1:A:344:ARG:HH11	1:A:351:GLN:HE21	1.61	0.47
1:A:247:LEU:HD12	1:A:285:VAL:HG22	1.97	0.46
1:A:133:ASN:HD21	4:A:1002:EDO:H12	1.81	0.46
1:A:137:ASP:HB3	5:A:1103:HOH:O	2.16	0.45
1:A:838:ILE:HD13	1:A:850:GLN:HG3	1.98	0.45
1:A:300:CYS:N	1:A:301:PRO:HD2	2.31	0.45
1:A:671:LEU:HB3	1:A:675:VAL:HG21	1.97	0.45
1:A:318:THR:HA	1:A:321:VAL:HG13	1.98	0.45
1:A:62:ASN:HB3	1:A:74:ILE:HD11	1.97	0.45
1:A:20:LYS:HE2	5:A:1168:HOH:O	2.17	0.45
1:A:389:PHE:CG	1:A:566:ILE:HD11	2.52	0.45
1:A:435:THR:O	1:A:436:SER:CB	2.64	0.45
1:A:699:ARG:CG	1:A:699:ARG:NH2	2.69	0.45
1:A:699:ARG:NH2	1:A:699:ARG:HG2	2.31	0.44
1:A:759:TYR:CG	1:A:766:LEU:HD21	2.52	0.44
1:A:566:ILE:HD12	1:A:566:ILE:C	2.42	0.44
1:A:627:ALA:N	5:A:1157:HOH:O	2.39	0.44
1:A:16:GLN:HE21	1:A:16:GLN:HB3	1.65	0.44
1:A:523:GLN:HA	1:A:523:GLN:OE1	2.17	0.44
1:A:173:GLY:HA2	5:A:1016:HOH:O	2.16	0.43
1:A:280:PHE:HE1	1:A:308:ALA:HB1	1.84	0.43
1:A:374:LYS:HE2	5:A:1167:HOH:O	2.18	0.43
1:A:410:ASN:O	1:A:412:HIS:HD2	2.01	0.43
1:A:616:ILE:HG23	1:A:617:GLU:HG3	2.00	0.43
1:A:689:ASN:HD22	1:A:689:ASN:N	2.15	0.43
1:A:329:LYS:HG3	1:A:332:TYR:OH	2.18	0.43
1:A:176:ILE:HG13	1:A:474:GLY:HA2	2.00	0.42
1:A:329:LYS:HA	1:A:332:TYR:CE2	2.54	0.42
1:A:471:LEU:HD13	1:A:498:VAL:HG22	2.02	0.42
1:A:661:SER:HA	1:A:839:ILE:HD12	2.02	0.42
1:A:110:ASP:HA	1:A:111:PRO:HD3	1.90	0.42
1:A:82:ARG:NH2	1:A:181:ASN:HD21	1.85	0.42
1:A:733:PHE:HB3	1:A:742:ILE:HD12	2.02	0.42
1:A:79:LYS:HG3	1:A:234:PRO:HB3	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:19:LYS:HB3	1:A:196:TYR:CE2	2.54	0.41
1:A:242:VAL:HG22	1:A:277:PHE:HB2	2.01	0.41
1:A:178:HIS:CE1	1:A:208:THR:HA	2.55	0.41
1:A:199:ASP:HB3	1:A:216:ILE:HG12	2.02	0.41
1:A:287:GLN:CB	5:A:1148:HOH:O	2.54	0.41
1:A:452:LEU:C	1:A:452:LEU:HD23	2.45	0.41
1:A:450:GLY:HA3	1:A:487:VAL:HG11	2.03	0.41
1:A:846:GLN:HA	1:A:863:MET:O	2.20	0.41
1:A:312:VAL:HG11	1:A:337:LEU:HD13	2.03	0.41
1:A:115:ASN:ND2	1:A:164:PHE:HA	2.36	0.40
1:A:331:LYS:HD3	1:A:331:LYS:HA	1.94	0.40
1:A:689:ASN:CG	1:A:689:ASN:O	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	886/888 (100%)	857 (97%)	25 (3%)	4 (0%)	24	34

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	580	LYS
1	A	403	GLN
1	A	436	SER
1	A	272	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	751/751 (100%)	704 (94%)	47 (6%)	16	23

All (47) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	SER
1	A	16	GLN
1	A	26	LYS
1	A	27	LEU
1	A	60	ILE
1	A	67	ASN
1	A	74	ILE
1	A	86	GLN
1	A	133	ASN
1	A	140	GLN
1	A	145	LEU
1	A	167	MET
1	A	250	LYS
1	A	262	LEU
1	A	288	LEU
1	A	321	VAL
1	A	329	LYS
1	A	330	LEU
1	A	371	SER
1	A	379	LYS
1	A	380	VAL
1	A	385	MET
1	A	396	LYS
1	A	413	LYS
1	A	435	THR
1	A	471	LEU
1	A	478	VAL
1	A	484	GLU
1	A	487	VAL
1	A	507	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	527	VAL
1	A	532	LEU
1	A	541	ARG
1	A	571	GLU
1	A	572	LYS
1	A	578	ASN
1	A	580	LYS
1	A	597	GLN
1	A	610	LYS
1	A	657	GLN
1	A	678	ASP
1	A	689	ASN
1	A	699	ARG
1	A	730	LEU
1	A	844	LYS
1	A	858	THR
1	A	869	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	16	GLN
1	A	25	ASN
1	A	62	ASN
1	A	65	HIS
1	A	73	ASN
1	A	115	ASN
1	A	133	ASN
1	A	143	GLN
1	A	166	ASN
1	A	181	ASN
1	A	192	GLN
1	A	233	GLN
1	A	322	GLN
1	A	351	GLN
1	A	403	GLN
1	A	412	HIS
1	A	439	ASN
1	A	493	GLN
1	A	509	ASN
1	A	545	ASN
1	A	549	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	602	GLN
1	A	689	ASN
1	A	698	ASN
1	A	740	GLN
1	A	743	HIS
1	A	807	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SF4	A	1000	5,1	0,12,12	-	-	-		
4	EDO	A	1002	-	3,3,3	0.57	0	2,2,2	0.10	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	EDO	A	1002	-	-	1/1/1/1	-
3	SF4	A	1000	5,1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1002	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1002	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	579:ALA	C	580:LYS	N	1.07

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	888/888 (100%)	-0.13	6 (0%) 84 86	11, 30, 46, 64	59 (6%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	593	ARG	3.7
1	A	579	ALA	3.0
1	A	13	ASP	2.2
1	A	580	LYS	2.2
1	A	655	ASP	2.1
1	A	689	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	A	1002	4/4	0.88	0.09	30,31,31,32	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SF4	A	1000	8/8	0.89	0.11	20,22,23,24	1
2	ZN	A	1001	1/1	0.99	0.03	50,50,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.