



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:22 AM UTC

PDB ID : 5B4X / pdb_00005b4x
Title : Crystal structure of the ApoER2 ectodomain in complex with the Reelin R56 fragment
Authors : Yasui, N.; Hirai, H.; Yamashita, K.; Takagi, J.; Nogi, T.
Deposited on : 2016-04-20
Resolution : 3.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	725	% 81% 16% ..
1	C	725	2% 77% 20% ..
2	B	570	% 66% 17% • 16%
2	D	570	2% 69% 15% • 16%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 18892 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Reelin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	710	Total	C	N	O	S	Se	0	0	0
			5609	3559	955	1059	26	10			
1	C	709	Total	C	N	O	S	Se	0	0	0
			5601	3555	953	1057	26	10			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1946	GLY	-	expression tag	UNP Q60841
A	1947	ARG	-	expression tag	UNP Q60841
A	2101	ALA	CYS	engineered mutation	UNP Q60841
A	2663	ARG	-	expression tag	UNP Q60841
A	2664	LEU	-	expression tag	UNP Q60841
A	2665	GLU	-	expression tag	UNP Q60841
A	2666	ASN	-	expression tag	UNP Q60841
A	2667	LEU	-	expression tag	UNP Q60841
A	2668	TYR	-	expression tag	UNP Q60841
A	2669	PHE	-	expression tag	UNP Q60841
A	2670	GLN	-	expression tag	UNP Q60841
C	1946	GLY	-	expression tag	UNP Q60841
C	1947	ARG	-	expression tag	UNP Q60841
C	2101	ALA	CYS	engineered mutation	UNP Q60841
C	2663	ARG	-	expression tag	UNP Q60841
C	2664	LEU	-	expression tag	UNP Q60841
C	2665	GLU	-	expression tag	UNP Q60841
C	2666	ASN	-	expression tag	UNP Q60841
C	2667	LEU	-	expression tag	UNP Q60841
C	2668	TYR	-	expression tag	UNP Q60841
C	2669	PHE	-	expression tag	UNP Q60841
C	2670	GLN	-	expression tag	UNP Q60841

- Molecule 2 is a protein called Low density lipoprotein receptor-related protein 8, apolipoprotein e receptor, isoform CRA_e.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	479	Total 3798	C 2359	N 656	O 746	S 37	0	0	0
2	D	479	Total 3798	C 2359	N 656	O 746	S 37	0	0	0


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	40	SER	-	expression tag	UNP D3DQ39
B	607	ALA	-	expression tag	UNP D3DQ39
B	608	SER	-	expression tag	UNP D3DQ39
B	609	SER	-	expression tag	UNP D3DQ39
D	40	SER	-	expression tag	UNP D3DQ39
D	607	ALA	-	expression tag	UNP D3DQ39
D	608	SER	-	expression tag	UNP D3DQ39
D	609	SER	-	expression tag	UNP D3DQ39

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total 4	Ca 4	0	0
3	B	4	Total 4	Ca 4	0	0
3	C	4	Total 4	Ca 4	0	0
3	D	4	Total 4	Ca 4	0	0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	B	1	Total	C	N	O	0	0
			14	8	1	5		
4	C	1	Total	C	N	O	0	0
			14	8	1	5		
4	D	1	Total	C	N	O	0	0
			14	8	1	5		


3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Reelin




• Molecule 1: Reelin






• Molecule 2: Low density lipoprotein receptor-related protein 8, apolipoprotein e receptor, isoform CRA_e



• Molecule 2: Low density lipoprotein receptor-related protein 8, apolipoprotein e receptor, isoform CRA_e



4 Data and refinement statistics

Property	Value	Source
Space group	P 65	Depositor
Cell constants a, b, c, α , β , γ	205.95Å 205.95Å 169.84Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.52 – 3.20 49.52 – 3.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.52-3.20) 100.0 (49.52-3.20)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 3.19Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.200 , 0.255 0.199 , 0.255	Depositor DCC
R_{free} test set	3467 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å ²)	82.6	Xtriage
Anisotropy	0.120	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 64.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	18892	wwPDB-VP
Average B, all atoms (Å ²)	99.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.74% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/5755	0.81	3/7815 (0.0%)
1	C	0.49	0/5747	0.83	5/7803 (0.1%)
2	B	0.51	0/3885	0.85	8/5265 (0.2%)
2	D	0.48	0/3885	0.84	2/5265 (0.0%)
All	All	0.49	0/19272	0.83	18/26148 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	2643	GLN	CA-C-N	6.03	125.75	119.05
1	C	2643	GLN	C-N-CA	6.03	125.75	119.05
2	B	252	ASP	CA-C-N	6.00	125.48	119.24
2	B	252	ASP	C-N-CA	6.00	125.48	119.24
2	B	283	ASN	N-CA-C	5.96	119.77	110.17

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	434	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5609	0	5322	70	0
1	C	5601	0	5318	95	0
2	B	3798	0	3572	68	0
2	D	3798	0	3572	60	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0
3	C	4	0	0	0	0
3	D	4	0	0	0	0
4	A	28	0	26	0	0
4	B	14	0	13	0	0
4	C	14	0	13	1	0
4	D	14	0	13	0	0
All	All	18892	0	17849	284	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 284 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1975:PRO:HB2	1:C:2006:GLU:O	1.53	1.09
1:C:2023:GLN:NE2	1:C:2142:SER:HB2	1.79	0.97
2:D:367:HIS:ND1	2:D:387:GLY:HA3	1.81	0.95
1:C:2060:TYR:HE1	1:C:2062:SER:HG	0.93	0.89
1:C:2183:PHE:HB2	1:C:2186:MSE:HE2	1.56	0.87

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	706/725 (97%)	628 (89%)	76 (11%)	2 (0%)	36	68
1	C	705/725 (97%)	641 (91%)	64 (9%)	0	100	100
2	B	475/570 (83%)	431 (91%)	41 (9%)	3 (1%)	21	56
2	D	475/570 (83%)	421 (89%)	53 (11%)	1 (0%)	43	73
All	All	2361/2590 (91%)	2121 (90%)	234 (10%)	6 (0%)	36	68

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2174	PHE
2	B	318	MET
2	B	550	GLU
2	B	592	PRO
2	D	592	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	621/624 (100%)	610 (98%)	11 (2%)	51	74
1	C	620/624 (99%)	606 (98%)	14 (2%)	44	70
2	B	427/497 (86%)	418 (98%)	9 (2%)	47	71
2	D	427/497 (86%)	421 (99%)	6 (1%)	59	77
All	All	2095/2242 (93%)	2055 (98%)	40 (2%)	50	73

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	2366	VAL
2	D	221	CYS
1	C	2371	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	2533	LEU
2	D	460	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 20 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	2522	ASN
2	D	298	ASN
2	D	466	GLN
2	D	300	HIS
1	A	2658	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 16 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	A	4006	1	14,14,15	0.57	0	17,19,21	1.24	1 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	C	4005	1	14,14,15	0.52	0	17,19,21	1.48	1 (5%)
4	NAG	D	5005	2	14,14,15	0.60	0	17,19,21	1.64	2 (11%)
4	NAG	B	5005	2	14,14,15	0.47	0	17,19,21	1.87	1 (5%)
4	NAG	A	4005	1	14,14,15	0.48	0	17,19,21	1.41	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	4006	1	-	0/6/23/26	0/1/1/1
4	NAG	C	4005	1	-	0/6/23/26	0/1/1/1
4	NAG	D	5005	2	-	0/6/23/26	0/1/1/1
4	NAG	B	5005	2	-	0/6/23/26	0/1/1/1
4	NAG	A	4005	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	5005	NAG	C1-O5-C5	6.72	121.19	112.19
4	D	5005	NAG	C1-O5-C5	5.25	119.23	112.19
4	A	4005	NAG	C1-O5-C5	4.81	118.64	112.19
4	C	4005	NAG	C1-O5-C5	4.55	118.29	112.19
4	A	4006	NAG	C1-O5-C5	3.44	116.80	112.19

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	C	4005	NAG	1	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	700/725 (96%)	-0.07	10 (1%) 73 54	50, 89, 145, 176	0
1	C	699/725 (96%)	0.00	13 (1%) 66 46	44, 93, 158, 196	0
2	B	479/570 (84%)	-0.08	7 (1%) 72 52	45, 79, 223, 273	0
2	D	479/570 (84%)	0.01	10 (2%) 63 43	51, 90, 232, 316	0
All	All	2357/2590 (91%)	-0.04	40 (1%) 69 49	44, 88, 190, 316	0

The worst 5 of 40 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1992	GLU	4.2
2	D	607	ALA	3.9
1	A	2178	LEU	3.8
1	C	2200	SER	3.4
1	C	1955	PRO	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	D	5002	1/1	0.71	0.17	306,306,306,306	0
4	NAG	A	4005	14/15	0.78	0.10	105,112,121,123	0
4	NAG	B	5005	14/15	0.84	0.13	74,83,92,93	0
4	NAG	D	5005	14/15	0.84	0.12	79,85,89,101	0
4	NAG	A	4006	14/15	0.86	0.12	80,93,104,108	0
3	CA	B	5002	1/1	0.88	0.15	276,276,276,276	0
4	NAG	C	4005	14/15	0.89	0.11	77,82,93,101	0
3	CA	B	5003	1/1	0.90	0.06	196,196,196,196	0
3	CA	D	5003	1/1	0.92	0.05	213,213,213,213	0
3	CA	A	4002	1/1	0.92	0.08	118,118,118,118	0
3	CA	C	4002	1/1	0.94	0.07	132,132,132,132	0
3	CA	C	4003	1/1	0.95	0.05	68,68,68,68	0
3	CA	C	4001	1/1	0.95	0.05	120,120,120,120	0
3	CA	A	4001	1/1	0.96	0.05	108,108,108,108	0
3	CA	C	4004	1/1	0.96	0.07	56,56,56,56	0
3	CA	A	4003	1/1	0.96	0.05	66,66,66,66	0
3	CA	D	5004	1/1	0.97	0.04	129,129,129,129	0
3	CA	A	4004	1/1	0.97	0.04	62,62,62,62	0
3	CA	B	5004	1/1	0.98	0.03	113,113,113,113	0
3	CA	B	5001	1/1	0.99	0.05	86,86,86,86	0
3	CA	D	5001	1/1	0.99	0.08	97,97,97,97	0

6.5 Other polymers

There are no such residues in this entry.