



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 05:57 PM UTC

PDB ID : 2B5D / pdb_00002b5d
Title : Crystal structure of the novel alpha-amylase AmyC from *Thermotoga maritima*
Authors : Dickmanns, A.; Ballschmiter, M.; Liebl, W.; Ficner, R.
Deposited on : 2005-09-28
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	X	528	7% 69% 23% 5%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha-Amylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	X	518	Total	C	N	O	S	0	4	0
			4343	2802	728	789	24			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	X	317	Total	O	0	0
			317	317		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: alpha-Amylase



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	112.16Å 112.16Å 335.54Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 30.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	90.8 (30.00-2.20) 90.7 (30.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0013	Depositor
R, R_{free}	0.219 , 0.257 0.220 , 0.211	Depositor DCC
R_{free} test set	2488 reflections (4.55%)	wwPDB-VP
Wilson B-factor (Å ²)	50.4	Xtriage
Anisotropy	0.462	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4660	wwPDB-VP
Average B, all atoms (Å ²)	60.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	X	0.99	8/4486 (0.2%)	1.11	11/6059 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	X	0	2

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	X	5	ILE	CA-CB	8.36	1.64	1.54
1	X	158	VAL	CA-CB	5.97	1.62	1.54
1	X	316	VAL	CA-CB	5.43	1.61	1.54
1	X	232	VAL	CA-CB	5.37	1.60	1.54
1	X	141	THR	CA-CB	5.24	1.60	1.53
1	X	514	ILE	CA-C	5.22	1.59	1.52
1	X	427	ILE	CA-CB	5.18	1.60	1.54
1	X	343	VAL	CA-CB	5.04	1.60	1.54

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	403	GLY	N-CA-C	6.66	120.91	110.71
1	X	514	ILE	N-CA-C	6.17	122.18	109.34
1	X	343	VAL	CB-CA-C	6.10	119.43	110.77
1	X	507	TYR	N-CA-C	-5.56	104.85	111.03
1	X	5	ILE	N-CA-CB	5.56	117.72	111.21
1	X	525	ARG	CB-CG-CD	-5.46	98.73	111.30
1	X	385	VAL	N-CA-C	5.36	116.13	110.72
1	X	228	THR	CA-C-N	-5.17	113.47	119.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	X	228	THR	C-N-CA	-5.17	113.47	119.47
1	X	242	SER	N-CA-C	5.07	116.65	110.41
1	X	86	ASN	N-CA-C	5.04	116.47	111.07

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	X	402	TRP	Peptide
1	X	527	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	X	4343	0	4177	120	0
2	X	317	0	0	17	0
All	All	4660	0	4177	120	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (120) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:179:ARG:NH2	1:X:202:GLU:OE2	1.83	1.10
1:X:63:GLU:OE2	1:X:514:ILE:O	1.74	1.06
1:X:219:ARG:H	1:X:393:GLN:HE22	1.04	1.02
1:X:5:ILE:HD11	1:X:379:LEU:CD2	1.93	0.98
1:X:526:ASP:HB2	2:X:565:HOH:O	1.62	0.97
1:X:352:LEU:HB3	2:X:650:HOH:O	1.65	0.97
1:X:145:THR:HG21	1:X:457:PHE:O	1.67	0.95
1:X:207:ASP:OD1	1:X:237:ARG:NH2	2.03	0.91
1:X:2:ARG:HB3	1:X:2:ARG:HH11	1.39	0.86
1:X:179:ARG:NE	2:X:709:HOH:O	2.08	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:351:GLU:HB3	2:X:666:HOH:O	1.78	0.84
1:X:179:ARG:CZ	2:X:709:HOH:O	2.24	0.83
1:X:179:ARG:NH1	2:X:709:HOH:O	2.12	0.82
1:X:5:ILE:HD11	1:X:379:LEU:HD23	1.63	0.81
1:X:179:ARG:NH2	1:X:202:GLU:CD	2.41	0.79
1:X:454[B]:ARG:HH21	1:X:514:ILE:N	1.82	0.77
1:X:219:ARG:N	1:X:393:GLN:HE22	1.82	0.75
1:X:2:ARG:HH11	1:X:2:ARG:CB	2.00	0.73
1:X:377:LEU:O	2:X:529:HOH:O	2.09	0.70
1:X:269:ARG:HG2	1:X:273:TYR:CD1	2.26	0.69
1:X:96:HIS:HD2	1:X:98:LEU:H	1.39	0.69
1:X:179:ARG:HH22	1:X:202:GLU:CD	2.00	0.69
1:X:526:ASP:CB	2:X:565:HOH:O	2.28	0.68
1:X:2:ARG:HH11	1:X:2:ARG:CG	2.06	0.67
1:X:219:ARG:H	1:X:393:GLN:NE2	1.86	0.67
1:X:454[B]:ARG:NH1	1:X:515:PHE:O	2.26	0.67
1:X:223:TYR:HB2	1:X:338:MET:HE1	1.78	0.64
1:X:5:ILE:CD1	1:X:379:LEU:HD23	2.26	0.64
1:X:526:ASP:CG	2:X:565:HOH:O	2.39	0.63
1:X:12:HIS:HD2	1:X:58:THR:OG1	1.80	0.63
1:X:224:ARG:NH2	1:X:389:LEU:HG	2.13	0.63
1:X:185:GLU:O	1:X:401:SER:HB2	1.99	0.62
1:X:258:TYR:OH	1:X:319:HIS:HD2	1.83	0.62
1:X:100:HIS:CE1	1:X:104:LYS:HD3	2.35	0.61
1:X:438:TYR:O	1:X:521:ARG:HD3	2.00	0.60
1:X:454[B]:ARG:NH1	1:X:454[B]:ARG:HG2	2.15	0.60
1:X:266:GLY:O	1:X:284:ARG:HD2	2.01	0.59
1:X:155:TYR:OH	1:X:424:HIS:HD2	1.85	0.59
1:X:454[B]:ARG:HH12	1:X:515:PHE:HB2	1.66	0.58
1:X:416:ASN:HB3	2:X:663:HOH:O	2.02	0.58
1:X:309:ILE:O	1:X:313:MET:HG2	2.03	0.58
1:X:132:GLN:NE2	1:X:139:ILE:H	2.02	0.58
1:X:317:GLU:HG3	1:X:321:ARG:NH1	2.19	0.58
1:X:527:VAL:O	1:X:528:ILE:HB	2.06	0.56
1:X:88:GLU:OE2	1:X:269:ARG:HD3	2.06	0.55
1:X:5:ILE:CD1	1:X:379:LEU:CD2	2.76	0.55
1:X:8:PHE:O	1:X:347:PRO:HA	2.07	0.54
1:X:473:SER:O	1:X:475:GLN:N	2.40	0.54
1:X:78:MET:HB3	1:X:114:LEU:HG	1.90	0.54
1:X:265:ILE:HD13	1:X:289:ILE:CD1	2.39	0.53
1:X:63:GLU:OE1	1:X:173:HIS:CE1	2.61	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:86:ASN:HA	1:X:89:VAL:HG12	1.91	0.53
1:X:494:GLN:NE2	1:X:501:ASP:H	2.07	0.53
1:X:450:ASN:ND2	1:X:520:PHE:H	2.06	0.53
1:X:2:ARG:HH22	1:X:375:LYS:HA	1.75	0.52
1:X:78:MET:CB	1:X:114:LEU:HG	2.39	0.52
1:X:429:ARG:HG2	1:X:492:TYR:CE1	2.46	0.51
1:X:2:ARG:HH12	1:X:375:LYS:C	2.18	0.51
1:X:203:TYR:HA	1:X:233:PHE:O	2.09	0.51
1:X:253:PRO:HG3	1:X:357:TRP:CD2	2.46	0.51
1:X:455:GLU:OE1	1:X:512:ASP:OD2	2.28	0.51
1:X:450:ASN:O	1:X:454[A]:ARG:HG3	2.10	0.51
1:X:522:VAL:O	1:X:525:ARG:HD2	2.11	0.51
1:X:8:PHE:CZ	1:X:182[B]:TRP:CD1	2.99	0.50
1:X:494:GLN:HB3	1:X:500:ILE:HA	1.94	0.50
1:X:5:ILE:HD11	1:X:379:LEU:HD22	1.91	0.50
1:X:17:HIS:O	2:X:665:HOH:O	2.19	0.50
1:X:30:LEU:O	1:X:34:ILE:HG13	2.11	0.50
1:X:480:ARG:CZ	1:X:484:HIS:HE1	2.24	0.49
1:X:2:ARG:NH1	1:X:2:ARG:HG2	2.26	0.49
1:X:237:ARG:HG2	2:X:562:HOH:O	2.13	0.49
1:X:89:VAL:HG21	1:X:107:ARG:HH21	1.77	0.49
1:X:93:LYS:HA	1:X:100:HIS:CD2	2.48	0.49
1:X:258:TYR:OH	1:X:319:HIS:CD2	2.65	0.48
1:X:2:ARG:CG	1:X:2:ARG:NH1	2.68	0.48
1:X:455:GLU:OE2	1:X:484:HIS:HD2	1.97	0.48
1:X:361:VAL:HG22	2:X:532:HOH:O	2.14	0.48
1:X:450:ASN:HD21	1:X:520:PHE:H	1.60	0.48
1:X:327:LYS:HA	1:X:327:LYS:HE2	1.97	0.47
1:X:102:MET:HE1	1:X:309:ILE:HB	1.96	0.47
1:X:229:PRO:CD	1:X:396:THR:HG22	2.44	0.47
1:X:480:ARG:HD2	1:X:480:ARG:HA	1.70	0.47
1:X:242:SER:O	1:X:245:VAL:HG13	2.14	0.47
1:X:317:GLU:HG3	1:X:321:ARG:HH11	1.78	0.47
1:X:88:GLU:CD	1:X:269:ARG:HH11	2.22	0.47
1:X:132:GLN:HE21	1:X:139:ILE:H	1.62	0.46
1:X:182[B]:TRP:HD1	1:X:205:PHE:CD1	2.34	0.46
1:X:55:MET:O	1:X:139:ILE:HA	2.15	0.45
1:X:289:ILE:HD11	1:X:291:TYR:CZ	2.52	0.45
1:X:218:PRO:HA	1:X:393:GLN:NE2	2.31	0.45
1:X:237:ARG:HH21	1:X:237:ARG:HD3	1.58	0.45
1:X:54:THR:HG23	1:X:138:GLU:HG3	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:X:179:ARG:NH2	1:X:202:GLU:OE1	2.47	0.44
1:X:255:ASP:HB3	1:X:258:TYR:CD1	2.52	0.44
1:X:450:ASN:O	1:X:454[B]:ARG:HG3	2.18	0.44
1:X:2:ARG:HH11	1:X:2:ARG:HG2	1.82	0.44
1:X:252:TYR:HB2	1:X:253:PRO:HD3	1.99	0.44
1:X:195:TYR:OH	2:X:555:HOH:O	2.21	0.44
1:X:63:GLU:OE1	1:X:173:HIS:HE1	2.00	0.43
1:X:528:ILE:O	2:X:843:HOH:O	2.21	0.43
1:X:265:ILE:HD13	1:X:289:ILE:HD12	2.00	0.43
1:X:204:PHE:CD2	1:X:204:PHE:N	2.86	0.43
1:X:194:LEU:O	1:X:198:GLN:HG3	2.18	0.43
1:X:237:ARG:HD2	1:X:402:TRP:CE2	2.54	0.43
1:X:54:THR:HA	1:X:138:GLU:O	2.19	0.43
1:X:149:LEU:N	1:X:150:PRO:CD	2.83	0.42
1:X:145:THR:CG2	1:X:162:GLN:HE22	2.32	0.42
1:X:182[B]:TRP:CD1	1:X:205:PHE:CD1	3.08	0.42
1:X:364:LEU:HD12	1:X:368:PHE:CZ	2.55	0.42
1:X:44:MET:HE2	1:X:364:LEU:HB3	2.02	0.41
1:X:425:GLU:O	1:X:429:ARG:HD3	2.20	0.41
1:X:289:ILE:HD11	1:X:291:TYR:CE1	2.56	0.41
1:X:454[B]:ARG:HH12	1:X:515:PHE:C	2.26	0.41
1:X:182[B]:TRP:HZ3	2:X:673:HOH:O	2.04	0.41
1:X:454[B]:ARG:HH21	1:X:513:ALA:C	2.28	0.41
1:X:228:THR:O	1:X:229:PRO:C	2.64	0.40
1:X:220:TYR:CD1	1:X:224:ARG:HD3	2.57	0.40
1:X:511:THR:O	2:X:665:HOH:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	X	518/528 (98%)	484 (93%)	30 (6%)	4 (1%)	16 16

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	X	474	VAL
1	X	470	THR
1	X	514	ILE
1	X	473	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	X	462/474 (98%)	411 (89%)	51 (11%)	6 6

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	X	2	ARG
1	X	4	LYS
1	X	6	LEU
1	X	23	HIS
1	X	26	GLU
1	X	30	LEU
1	X	42	LEU
1	X	70	LEU
1	X	76	ARG
1	X	84	LEU
1	X	90[A]	GLU
1	X	90[B]	GLU
1	X	102	MET
1	X	104	LYS
1	X	112	LYS
1	X	114	LEU
1	X	115	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	145	THR
1	X	179	ARG
1	X	192	LEU
1	X	194	LEU
1	X	196	LEU
1	X	204	PHE
1	X	230	SER
1	X	243	GLU
1	X	245	VAL
1	X	259	ARG
1	X	269	ARG
1	X	283	VAL
1	X	289	ILE
1	X	299	LEU
1	X	313	MET
1	X	316	VAL
1	X	334	LEU
1	X	343	VAL
1	X	349	ASP
1	X	361	VAL
1	X	364	LEU
1	X	370	LEU
1	X	385	VAL
1	X	392	VAL
1	X	396	THR
1	X	421	ARG
1	X	423	LEU
1	X	429	ARG
1	X	444	LEU
1	X	458	LEU
1	X	468	MET
1	X	522	VAL
1	X	525	ARG
1	X	528	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (22) such sidechains are listed below:

Mol	Chain	Res	Type
1	X	12	HIS
1	X	71	GLN
1	X	86	ASN
1	X	96	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	X	100	HIS
1	X	109	HIS
1	X	123	ASN
1	X	132	GLN
1	X	173	HIS
1	X	190	GLN
1	X	198	GLN
1	X	217	GLN
1	X	286	ASN
1	X	292	HIS
1	X	319	HIS
1	X	372	ASN
1	X	393	GLN
1	X	424	HIS
1	X	450	ASN
1	X	484	HIS
1	X	490	ASN
1	X	494	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	X	518/528 (98%)	0.58	37 (7%) 22 19	25, 58, 91, 108	4 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	X	23	HIS	4.5
1	X	468	MET	4.4
1	X	29	TRP	4.4
1	X	514	ILE	4.2
1	X	404	ALA	4.1
1	X	415	THR	4.0
1	X	474	VAL	3.9
1	X	476	TYR	3.8
1	X	467	ILE	3.7
1	X	469	THR	3.6
1	X	283	VAL	3.5
1	X	464	TRP	3.4
1	X	352	LEU	3.1
1	X	403	GLY	3.1
1	X	356	TRP	3.1
1	X	472	THR	3.0
1	X	465	ALA	3.0
1	X	402	TRP	3.0
1	X	282	GLY	2.9
1	X	354	GLY	2.9
1	X	477	ALA	2.8
1	X	94	LYS	2.8
1	X	470	THR	2.8
1	X	21	TYR	2.7
1	X	420	TYR	2.7
1	X	22	ASP	2.6
1	X	463	ASP	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	X	5	ILE	2.4
1	X	475	GLN	2.4
1	X	91	ARG	2.3
1	X	466	PHE	2.2
1	X	416	ASN	2.2
1	X	289	ILE	2.2
1	X	528	ILE	2.1
1	X	70	LEU	2.1
1	X	280	PRO	2.1
1	X	25	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.