



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:57 AM UTC

PDB ID : 2B5L / pdb_00002b5l
Title : Crystal Structure of DDB1 In Complex with Simian Virus 5 V Protein
Authors : Li, T.; Chen, X.; Garbutt, K.C.; Zhou, P.; Zheng, N.
Deposited on : 2005-09-28
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1140	2% 50% 40% 9% ..
1	B	1140	4% 38% 47% 12% ..
2	C	222	9% 27% 36% 13% . 22%
2	D	222	14% 23% 36% 16% . 21%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 20399 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called damage-specific DNA binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1132	Total	C	N	O	S	0	0	0
			8860	5610	1493	1708	49			
1	B	1134	Total	C	N	O	S	0	0	0
			8876	5619	1495	1713	49			

- Molecule 2 is a protein called Nonstructural protein V.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	174	Total	C	N	O	S	0	0	0
			1324	826	229	261	8			
2	D	175	Total	C	N	O	S	0	0	0
			1335	834	233	260	8			


- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	2	Total	Zn	0	0
			2	2		
3	D	2	Total	Zn	0	0
			2	2		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: damage-specific DNA binding protein 1







• Molecule 1: damage-specific DNA binding protein 1







• Molecule 2: Nonstructural protein V



• Molecule 2: Nonstructural protein V





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.76Å 240.79Å 117.18Å 90.00° 101.79° 90.00°	Depositor
Resolution (Å)	47.80 – 2.85 47.80 – 2.85	Depositor EDS
% Data completeness (in resolution range)	(Not available) (47.80-2.85) 95.7 (47.80-2.85)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.65Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.229 , 0.299 0.226 , 0.293	Depositor DCC
R_{free} test set	3805 reflections (4.13%)	wwPDB-VP
Wilson B-factor (Å ²)	46.9	Xtriage
Anisotropy	0.604	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 64.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	20399	wwPDB-VP
Average B, all atoms (Å ²)	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	5/9022 (0.1%)	1.13	81/12219 (0.7%)
1	B	0.60	14/9039 (0.2%)	1.32	118/12244 (1.0%)
2	C	0.79	8/1351 (0.6%)	1.42	27/1832 (1.5%)
2	D	1.08	4/1361 (0.3%)	1.63	32/1843 (1.7%)
All	All	0.64	31/20773 (0.1%)	1.27	258/28138 (0.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	1
2	D	0	1
All	All	1	2

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	189	TRP	C-N	-25.44	0.99	1.33
2	D	190	CYS	C-N	17.91	1.59	1.33
1	B	431	GLY	N-CA	-12.68	1.35	1.45
1	B	430	VAL	C-N	-9.31	1.25	1.33
1	B	455	GLN	N-CA	-8.16	1.35	1.46

The worst 5 of 258 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	431	GLY	N-CA-C	-42.26	71.75	110.21
1	B	749	THR	N-CA-C	-31.14	58.10	108.55

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	35	LYS	CB-CA-C	18.23	146.70	110.42
1	B	980	ASP	N-CA-C	-17.33	78.02	108.69
1	B	884	ILE	CB-CA-C	16.86	138.94	111.29

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	1	MET	CA

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	430	VAL	Peptide
2	D	190	CYS	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	8860	0	8828	495	0
1	B	8876	0	8842	826	0
2	C	1324	0	1306	134	0
2	D	1335	0	1323	179	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
All	All	20399	0	20299	1618	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 40.

The worst 5 of 1618 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:367:LEU:HD21	1:B:374:GLN:CG	1.51	1.38
1:B:367:LEU:HD21	1:B:374:GLN:CB	1.50	1.37
1:B:367:LEU:CD2	1:B:374:GLN:HB2	1.53	1.37

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:742:VAL:CG2	1:B:750:THR:HG21	1.57	1.35
1:B:367:LEU:CD2	1:B:374:GLN:CG	2.06	1.34

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1126/1140 (99%)	960 (85%)	128 (11%)	38 (3%)	3	6
1	B	1130/1140 (99%)	892 (79%)	145 (13%)	93 (8%)	0	1
2	C	168/222 (76%)	123 (73%)	33 (20%)	12 (7%)	1	1
2	D	169/222 (76%)	115 (68%)	32 (19%)	22 (13%)	0	0
All	All	2593/2724 (95%)	2090 (81%)	338 (13%)	165 (6%)	1	1

5 of 165 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	254	LYS
1	A	269	SER
1	A	290	GLN
1	A	341	ASN
1	A	370	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	992/999 (99%)	897 (90%)	95 (10%)	8	17
1	B	994/999 (100%)	916 (92%)	78 (8%)	11	25
2	C	151/191 (79%)	130 (86%)	21 (14%)	3	6
2	D	151/191 (79%)	133 (88%)	18 (12%)	5	10
All	All	2288/2380 (96%)	2076 (91%)	212 (9%)	8	18

5 of 212 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	392	ASN
1	B	780	THR
2	D	101	PRO
1	B	456	GLN
1	B	585	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 87 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	481	GLN
1	B	991	HIS
1	B	507	GLN
1	B	803	HIS
1	B	1070	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	D	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	D	189:TRP	C	190:CYS	N	0.99

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1132/1140 (99%)	-0.05	25 (2%) 62 53	18, 47, 99, 160	0
1	B	1134/1140 (99%)	0.39	47 (4%) 41 32	29, 68, 122, 173	0
2	C	174/222 (78%)	0.66	21 (12%) 8 6	32, 62, 118, 183	0
2	D	175/222 (78%)	1.05	31 (17%) 4 3	39, 81, 138, 168	0
All	All	2615/2724 (95%)	0.26	124 (4%) 36 28	18, 59, 117, 183	0

The worst 5 of 124 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	453	ASP	5.4
1	B	571	LEU	4.7
2	D	20	GLY	4.6
1	A	708	GLN	4.6
2	D	16	LEU	4.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	C	3001	1/1	0.89	0.13	35,35,35,35	0
3	ZN	C	3002	1/1	0.90	0.11	35,35,35,35	0
3	ZN	D	3004	1/1	0.95	0.18	35,35,35,35	0
3	ZN	D	3003	1/1	0.98	0.15	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.