



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 12:44 AM UTC

PDB ID : 6B57 / pdb_00006b57
Title : tudor in complex with ligand
Authors : Zhang, H.; Tempel, W.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.M.; Min, J.; Structural Genomics Consortium (SGC)
Deposited on : 2017-09-28
Resolution : 1.93 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	14% 84% • 13%
1	B	231	9% 86% • 10%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tudor and KH domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	202	Total	C	N	O	S	0	1	1
			1532	978	260	290	4			
1	B	207	Total	C	N	O	S	0	2	0
			1615	1026	284	301	4			

There are 84 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	291	MET	-	expression tag	UNP Q9Y2W6
A	292	HIS	-	expression tag	UNP Q9Y2W6
A	293	HIS	-	expression tag	UNP Q9Y2W6
A	294	HIS	-	expression tag	UNP Q9Y2W6
A	295	HIS	-	expression tag	UNP Q9Y2W6
A	296	HIS	-	expression tag	UNP Q9Y2W6
A	297	HIS	-	expression tag	UNP Q9Y2W6
A	298	SER	-	expression tag	UNP Q9Y2W6
A	299	SER	-	expression tag	UNP Q9Y2W6
A	300	GLY	-	expression tag	UNP Q9Y2W6
A	301	ARG	-	expression tag	UNP Q9Y2W6
A	302	GLU	-	expression tag	UNP Q9Y2W6
A	303	ASN	-	expression tag	UNP Q9Y2W6
A	304	LEU	-	expression tag	UNP Q9Y2W6
A	305	TYR	-	expression tag	UNP Q9Y2W6
A	306	PHE	-	expression tag	UNP Q9Y2W6
A	307	GLN	-	expression tag	UNP Q9Y2W6
A	308	GLY	-	expression tag	UNP Q9Y2W6
A	498	GLY	-	expression tag	UNP Q9Y2W6
A	499	SER	-	expression tag	UNP Q9Y2W6
A	500	ALA	-	expression tag	UNP Q9Y2W6
A	501	GLY	-	expression tag	UNP Q9Y2W6
A	502	SER	-	expression tag	UNP Q9Y2W6
A	503	ALA	-	expression tag	UNP Q9Y2W6
A	504	GLY	-	expression tag	UNP Q9Y2W6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	505	SER	-	expression tag	UNP Q9Y2W6
A	506	ALA	-	expression tag	UNP Q9Y2W6
A	507	GLY	-	expression tag	UNP Q9Y2W6
A	508	SER	-	expression tag	UNP Q9Y2W6
A	509	ALA	-	expression tag	UNP Q9Y2W6
A	510	THR	-	expression tag	UNP Q9Y2W6
A	511	GLY	-	expression tag	UNP Q9Y2W6
A	512	ARG	-	expression tag	UNP Q9Y2W6
A	513	ALA	-	expression tag	UNP Q9Y2W6
A	514	ARG	-	expression tag	UNP Q9Y2W6
A	515	ALA	-	expression tag	UNP Q9Y2W6
A	516	ARG	-	expression tag	UNP Q9Y2W6
A	517	ALA	-	expression tag	UNP Q9Y2W6
A	518	ARG	-	expression tag	UNP Q9Y2W6
A	519	GLY	-	expression tag	UNP Q9Y2W6
A	520	ARG	-	expression tag	UNP Q9Y2W6
A	521	ALA	-	expression tag	UNP Q9Y2W6
B	291	MET	-	expression tag	UNP Q9Y2W6
B	292	HIS	-	expression tag	UNP Q9Y2W6
B	293	HIS	-	expression tag	UNP Q9Y2W6
B	294	HIS	-	expression tag	UNP Q9Y2W6
B	295	HIS	-	expression tag	UNP Q9Y2W6
B	296	HIS	-	expression tag	UNP Q9Y2W6
B	297	HIS	-	expression tag	UNP Q9Y2W6
B	298	SER	-	expression tag	UNP Q9Y2W6
B	299	SER	-	expression tag	UNP Q9Y2W6
B	300	GLY	-	expression tag	UNP Q9Y2W6
B	301	ARG	-	expression tag	UNP Q9Y2W6
B	302	GLU	-	expression tag	UNP Q9Y2W6
B	303	ASN	-	expression tag	UNP Q9Y2W6
B	304	LEU	-	expression tag	UNP Q9Y2W6
B	305	TYR	-	expression tag	UNP Q9Y2W6
B	306	PHE	-	expression tag	UNP Q9Y2W6
B	307	GLN	-	expression tag	UNP Q9Y2W6
B	308	GLY	-	expression tag	UNP Q9Y2W6
B	498	GLY	-	expression tag	UNP Q9Y2W6
B	499	SER	-	expression tag	UNP Q9Y2W6
B	500	ALA	-	expression tag	UNP Q9Y2W6
B	501	GLY	-	expression tag	UNP Q9Y2W6
B	502	SER	-	expression tag	UNP Q9Y2W6
B	503	ALA	-	expression tag	UNP Q9Y2W6
B	504	GLY	-	expression tag	UNP Q9Y2W6

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	505	SER	-	expression tag	UNP Q9Y2W6
B	506	ALA	-	expression tag	UNP Q9Y2W6
B	507	GLY	-	expression tag	UNP Q9Y2W6
B	508	SER	-	expression tag	UNP Q9Y2W6
B	509	ALA	-	expression tag	UNP Q9Y2W6
B	510	THR	-	expression tag	UNP Q9Y2W6
B	511	GLY	-	expression tag	UNP Q9Y2W6
B	512	ARG	-	expression tag	UNP Q9Y2W6
B	513	ALA	-	expression tag	UNP Q9Y2W6
B	514	ARG	-	expression tag	UNP Q9Y2W6
B	515	ALA	-	expression tag	UNP Q9Y2W6
B	516	ARG	-	expression tag	UNP Q9Y2W6
B	517	ALA	-	expression tag	UNP Q9Y2W6
B	518	ARG	-	expression tag	UNP Q9Y2W6
B	519	GLY	-	expression tag	UNP Q9Y2W6
B	520	ARG	-	expression tag	UNP Q9Y2W6
B	521	ALA	-	expression tag	UNP Q9Y2W6

- Molecule 2 is UNKNOWN ATOM OR ION (CCD ID: UNX) (formula: X).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	15	Total X 15 15	0	0
2	B	19	Total X 19 19	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	54	Total O 54 54	0	0
3	B	52	Total O 52 52	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tudor and KH domain-containing protein



- Molecule 1: Tudor and KH domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.07Å 72.66Å 130.70Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.35 – 1.93 65.35 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.0 (65.35-1.93) 99.3 (65.35-1.93)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.85 (at 1.94Å)	Xtriage
Refinement program	BUSTER 2.10.2	Depositor
R, R_{free}	0.206 , 0.234 (Not available) , 0.265	Depositor DCC
R_{free} test set	1683 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	33.6	Xtriage
Anisotropy	0.313	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 42.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3287	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.64% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: UNX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	0/1572	1.20	2/2146 (0.1%)
1	B	0.85	0/1662	1.22	6/2262 (0.3%)
All	All	0.85	0/3234	1.21	8/4408 (0.2%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	428	GLY	CA-C-N	7.65	130.39	120.44
1	B	428	GLY	C-N-CA	7.65	130.39	120.44
1	B	316	GLU	N-CA-C	-7.18	101.83	111.96
1	A	316	GLU	N-CA-C	-6.88	102.25	111.96
1	B	481	ASP	CA-CB-CG	6.10	118.70	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1532	0	1407	3	0
1	B	1615	0	1519	1	0
2	A	15	0	0	0	0
2	B	19	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	54	0	0	0	0
3	B	52	0	0	0	0
All	All	3287	0	2926	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:481:ASP:HB3	1:A:484:LEU:HB2	1.91	0.51
1:B:386:LEU:HD11	1:B:397:CYS:HB2	1.93	0.49
1:A:326:VAL:HG11	1:A:450:PRO:HG3	1.98	0.46
1:A:386:LEU:HD11	1:A:397:CYS:HB2	2.00	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/231 (86%)	193 (97%)	6 (3%)	0	100	100
1	B	205/231 (89%)	202 (98%)	3 (2%)	0	100	100
All	All	404/462 (87%)	395 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	150/191 (78%)	149 (99%)	1 (1%)	76	74
1	B	163/191 (85%)	160 (98%)	3 (2%)	51	43
All	All	313/382 (82%)	309 (99%)	4 (1%)	61	54

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	323	ILE
1	B	350	GLU
1	B	352	LEU
1	B	386	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	381	ASN
1	B	383	ASN
1	B	488	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 34 ligands modelled in this entry, 34 are unknown - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	202/231 (87%)	0.78	33 (16%) 4 4	24, 45, 85, 96	1 (0%)
1	B	207/231 (89%)	0.52	21 (10%) 12 14	22, 41, 75, 87	2 (0%)
All	All	409/462 (88%)	0.65	54 (13%) 7 8	22, 43, 80, 96	3 (0%)

The worst 5 of 54 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	495	LEU	6.0
1	A	496	PRO	5.7
1	A	435	ALA	4.2
1	B	353	THR	3.9
1	A	495	LEU	3.9

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	UNX	A	615	1/1	0.45	0.23	60,60,60,60	0
2	UNX	B	604	1/1	0.63	0.19	58,58,58,58	0
2	UNX	B	618	1/1	0.68	0.20	52,52,52,52	0
2	UNX	B	606	1/1	0.80	0.16	39,39,39,39	0
2	UNX	A	602	1/1	0.81	0.14	41,41,41,41	0
2	UNX	A	607	1/1	0.83	0.15	50,50,50,50	0
2	UNX	B	608	1/1	0.84	0.15	33,33,33,33	0
2	UNX	B	611	1/1	0.85	0.12	29,29,29,29	0
2	UNX	B	601	1/1	0.87	0.15	31,31,31,31	0
2	UNX	A	608	1/1	0.87	0.11	48,48,48,48	0
2	UNX	B	614	1/1	0.88	0.14	30,30,30,30	0
2	UNX	A	606	1/1	0.88	0.11	39,39,39,39	0
2	UNX	A	603	1/1	0.89	0.15	46,46,46,46	0
2	UNX	B	616	1/1	0.89	0.14	31,31,31,31	0
2	UNX	B	603	1/1	0.89	0.16	40,40,40,40	0
2	UNX	A	613	1/1	0.90	0.10	40,40,40,40	0
2	UNX	A	614	1/1	0.90	0.20	50,50,50,50	0
2	UNX	B	617	1/1	0.90	0.18	45,45,45,45	0
2	UNX	A	610	1/1	0.90	0.09	38,38,38,38	0
2	UNX	B	619	1/1	0.91	0.15	33,33,33,33	0
2	UNX	B	613	1/1	0.92	0.10	30,30,30,30	0
2	UNX	A	604	1/1	0.92	0.37	46,46,46,46	0
2	UNX	A	605	1/1	0.93	0.13	38,38,38,38	0
2	UNX	B	615	1/1	0.94	0.11	33,33,33,33	0
2	UNX	A	601	1/1	0.95	0.06	26,26,26,26	0
2	UNX	B	607	1/1	0.96	0.13	33,33,33,33	0
2	UNX	A	611	1/1	0.96	0.17	30,30,30,30	0
2	UNX	A	612	1/1	0.96	0.08	29,29,29,29	0
2	UNX	B	605	1/1	0.96	0.18	34,34,34,34	0
2	UNX	A	609	1/1	0.96	0.11	33,33,33,33	0
2	UNX	B	602	1/1	0.97	0.06	30,30,30,30	0
2	UNX	B	610	1/1	0.97	0.12	33,33,33,33	0
2	UNX	B	609	1/1	0.98	0.08	30,30,30,30	0
2	UNX	B	612	1/1	0.99	0.11	28,28,28,28	0

6.5 Other polymers ⓘ

There are no such residues in this entry.