



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 10:28 AM UTC

PDB ID : 6B5J / pdb_00006b5j
Title : TNNI3K complexed with a 4,6-diaminopyrimidine
Authors : Shewchuk, L.M.; Philp, J.
Deposited on : 2017-09-29
Resolution : 2.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3580 (3.00-2.96)
Clashscore	190562	3904 (3.00-2.96)
Ramachandran outliers	187476	3761 (3.00-2.96)
Sidechain outliers	187428	3764 (3.00-2.96)
RSRZ outliers	180081	3579 (3.00-2.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	311	2% 80% 15%
1	B	311	4% 77% 6% 15%
1	C	311	3% 77% 7% 16%
1	D	311	3% 80% 17%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8360 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serine/threonine-protein kinase TNNI3K.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	263	Total	C	N	O	S	0	5	0
			2087	1345	354	372	16			
1	B	264	Total	C	N	O	S	0	5	0
			2089	1346	354	371	18			
1	C	260	Total	C	N	O	S	0	0	0
			2028	1302	351	360	15			
1	D	257	Total	C	N	O	S	0	1	0
			2007	1289	343	359	16			

- Molecule 2 is N-methyl-3-[(6-{[4-(trifluoromethyl)phenyl]amino}pyrimidin-4-yl)amino]benzene-1-sulfonamide (CCD ID: CV4) (formula: $C_{18}H_{16}F_3N_5O_2S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	F	N	O	S	0
			29	18	3	5	2	1	
2	B	1	Total	C	F	N	O	S	0
			29	18	3	5	2	1	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	F	N	O	S	
			29	18	3	5	2	1	
2	D	1	Total	C	F	N	O	S	
			29	18	3	5	2	1	


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	O		
			7	7	0	0
3	B	11	Total	O		
			11	11	0	0
3	C	12	Total	O		
			12	12	0	0
3	D	3	Total	O		
			3	3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serine/threonine-protein kinase TNNI3K




- Molecule 1: Serine/threonine-protein kinase TNNI3K




- Molecule 1: Serine/threonine-protein kinase TNNI3K



- Molecule 1: Serine/threonine-protein kinase TNNI3K





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	85.61Å 108.19Å 92.65Å 90.00° 94.71° 90.00°	Depositor
Resolution (Å)	92.45 – 2.97 92.34 – 2.97	Depositor EDS
% Data completeness (in resolution range)	98.1 (92.45-2.97) 98.3 (92.34-2.97)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.53 (at 2.95Å)	Xtriage
Refinement program	REFMAC 5.8.0135	Depositor
R, R_{free}	0.201 , 0.258 0.205 , 0.258	Depositor DCC
R_{free} test set	1717 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	59.8	Xtriage
Anisotropy	0.214	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 28.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.52$, $\langle L^2 \rangle = 0.36$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8360	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 39.89 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.9665e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CV4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.93	0/2139	1.06	3/2899 (0.1%)
1	B	0.94	1/2142 (0.0%)	1.10	5/2905 (0.2%)
1	C	0.92	0/2076	1.00	1/2814 (0.0%)
1	D	0.95	0/2055	1.05	0/2788
All	All	0.94	1/8412 (0.0%)	1.05	9/11406 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	725	ASN	N-CA	5.61	1.53	1.46

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	491	ARG	CD-NE-CZ	11.53	140.54	124.40
1	B	491	ARG	NE-CZ-NH1	-10.72	110.78	121.50
1	A	491	ARG	NE-CZ-NH2	-10.63	109.63	119.20
1	B	491	ARG	NE-CZ-NH2	8.64	126.97	119.20
1	B	491	ARG	CD-NE-CZ	7.64	135.09	124.40
1	B	491	ARG	CG-CD-NE	-7.03	96.54	112.00
1	B	663	ILE	N-CA-C	-6.86	102.64	109.02
1	A	663	ILE	N-CA-C	-6.63	102.85	109.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	681	HIS	N-CA-C	5.50	119.29	112.47

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	491	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2087	0	2033	4	0
1	B	2089	0	2040	11	0
1	C	2028	0	1989	11	0
1	D	2007	0	1963	4	0
2	A	29	0	0	2	0
2	B	29	0	0	3	0
2	C	29	0	0	1	0
2	D	29	0	0	1	0
3	A	7	0	0	0	0
3	B	11	0	0	0	0
3	C	12	0	0	0	0
3	D	3	0	0	0	0
All	All	8360	0	8025	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:492:TYR:OH	1:A:509:GLU:OE1	2.21	0.59
1:B:545:GLY:HA2	2:B:801:CV4:C21	2.33	0.57
1:B:638:GLN:HG3	1:C:637:THR:HG22	1.93	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:801:CV4:N28	2:B:801:CV4:C29	2.77	0.47
1:D:512:ILE:HG23	1:D:584:ILE:HD13	1.97	0.46
1:C:512:ILE:HG23	1:C:584:ILE:HD13	1.96	0.46
1:C:680:HIS:CE1	1:D:680:HIS:CE1	3.04	0.45
1:A:529:LEU:O	1:A:530:ASN:C	2.58	0.45
2:A:801:CV4:N28	2:A:801:CV4:C29	2.80	0.45
1:A:578:HIS:CE1	1:A:644:ILE:HB	2.52	0.45
1:B:578:HIS:CE1	1:B:644:ILE:HB	2.52	0.44
1:C:587:ARG:NH1	1:C:642:TYR:CG	2.85	0.44
1:B:640:THR:O	1:B:641:ARG:CB	2.65	0.44
2:C:801:CV4:N28	2:C:801:CV4:C29	2.80	0.44
1:D:578:HIS:CE1	1:D:644:ILE:HB	2.53	0.43
1:A:542:ILE:O	2:A:801:CV4:N15	2.51	0.43
1:B:567:ALA:O	1:B:651:TYR:OH	2.34	0.43
1:D:487:VAL:HG23	1:D:489:ILE:HD11	2.01	0.43
1:B:492:TYR:OH	1:B:509:GLU:OE1	2.36	0.43
1:C:452:LEU:O	1:C:453:PRO:C	2.62	0.43
2:D:801:CV4:N28	2:D:801:CV4:C29	2.80	0.43
1:C:590:ASN:HA	1:C:654:CYS:SG	2.59	0.42
1:B:545:GLY:HA2	2:B:801:CV4:C20	2.49	0.42
1:B:585:ILE:HG12	1:B:644:ILE:HA	2.00	0.42
1:C:701:GLY:HA2	1:C:710:PRO:HD2	2.03	0.41
1:B:674:ALA:HB1	1:C:636:PHE:CD1	2.56	0.41
1:C:487:VAL:HG23	1:C:489:ILE:HD11	2.02	0.41
1:C:578:HIS:CE1	1:C:644:ILE:HB	2.57	0.40
1:C:691:PRO:HB2	1:C:693:PRO:HD2	2.03	0.40
1:B:659:LEU:CD1	1:B:694:ILE:HG21	2.51	0.40
1:B:700:ARG:HH22	1:B:714:GLU:CD	2.29	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	261/311 (84%)	252 (97%)	7 (3%)	2 (1%)	16	47
1	B	263/311 (85%)	255 (97%)	6 (2%)	2 (1%)	16	47
1	C	252/311 (81%)	241 (96%)	11 (4%)	0	100	100
1	D	252/311 (81%)	241 (96%)	11 (4%)	0	100	100
All	All	1028/1244 (83%)	989 (96%)	35 (3%)	4 (0%)	30	62

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	530	ASN
1	B	641	ARG
1	A	641	ARG
1	B	530	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/270 (82%)	217 (98%)	4 (2%)	51	76
1	B	221/270 (82%)	216 (98%)	5 (2%)	44	72
1	C	217/270 (80%)	215 (99%)	2 (1%)	70	85
1	D	216/270 (80%)	215 (100%)	1 (0%)	81	89
All	All	875/1080 (81%)	863 (99%)	12 (1%)	59	80

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	531	ASP
1	A	542	ILE
1	A	611	ARG
1	A	713	SER
1	B	531	ASP
1	B	542	ILE
1	B	611	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	640	THR
1	B	725	ASN
1	C	611	ARG
1	C	725	ASN
1	D	611	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	530	ASN
1	A	592	HIS
1	A	627	ASN
1	B	515	GLN
1	B	530	ASN
1	B	592	HIS
1	B	627	ASN
1	C	455	HIS
1	C	540	GLN
1	C	592	HIS
1	C	681	HIS
1	D	455	HIS
1	D	466	HIS
1	D	592	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	CV4	B	801	-	31,31,31	1.60	1 (3%)	45,45,45	1.52	7 (15%)
2	CV4	C	801	-	31,31,31	1.52	1 (3%)	45,45,45	1.44	4 (8%)
2	CV4	A	801	-	31,31,31	1.44	2 (6%)	45,45,45	1.38	5 (11%)
2	CV4	D	801	-	31,31,31	1.33	1 (3%)	45,45,45	1.47	5 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CV4	B	801	-	-	2/23/23/23	0/3/3/3
2	CV4	C	801	-	-	2/23/23/23	0/3/3/3
2	CV4	A	801	-	-	2/23/23/23	0/3/3/3
2	CV4	D	801	-	-	2/23/23/23	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	801	CV4	S03-N02	7.82	1.73	1.61
2	C	801	CV4	S03-N02	7.09	1.72	1.61
2	A	801	CV4	S03-N02	6.68	1.71	1.61
2	D	801	CV4	S03-N02	5.91	1.70	1.61
2	A	801	CV4	O04-S03	2.11	1.46	1.43

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	801	CV4	C06-S03-N02	-5.20	100.38	107.55
2	B	801	CV4	C06-S03-N02	-4.83	100.89	107.55
2	A	801	CV4	C06-S03-N02	-4.52	101.32	107.55
2	B	801	CV4	C14-C13-C12	4.25	120.20	116.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	801	CV4	C06-S03-N02	-4.03	102.00	107.55
2	D	801	CV4	O05-S03-O04	4.01	124.39	119.52
2	A	801	CV4	O05-S03-O04	3.86	124.21	119.52
2	C	801	CV4	O05-S03-O04	3.59	123.88	119.52
2	D	801	CV4	C01-N02-S03	-2.83	115.42	119.16
2	C	801	CV4	C14-C13-C12	2.74	118.77	116.16
2	D	801	CV4	C14-C13-C12	2.69	118.72	116.16
2	B	801	CV4	O05-S03-C06	-2.65	104.64	107.98
2	B	801	CV4	O05-S03-O04	2.59	122.67	119.52
2	A	801	CV4	O05-S03-C06	-2.55	104.76	107.98
2	B	801	CV4	C13-C12-N28	-2.38	119.80	122.92
2	C	801	CV4	O04-S03-N02	2.35	111.39	107.06
2	D	801	CV4	C10-C29-C06	2.34	120.64	118.84
2	B	801	CV4	N11-C12-N28	2.27	123.89	116.78
2	B	801	CV4	O04-S03-N02	2.21	111.13	107.06
2	A	801	CV4	C18-C19-C22	2.07	123.28	119.96
2	A	801	CV4	C01-N02-S03	-2.03	116.47	119.16

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	801	CV4	C29-C06-S03-O05
2	A	801	CV4	C29-C06-S03-O05
2	B	801	CV4	C29-C06-S03-O05
2	D	801	CV4	C07-C06-S03-O05
2	A	801	CV4	C07-C06-S03-O05
2	B	801	CV4	C07-C06-S03-O05
2	C	801	CV4	C29-C06-S03-O05
2	C	801	CV4	C07-C06-S03-O05


There are no ring outliers.


4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	801	CV4	3	0
2	C	801	CV4	1	0
2	A	801	CV4	2	0
2	D	801	CV4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,

bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	263/311 (84%)	0.03	7 (2%) 56 37	17, 53, 95, 120	5 (1%)
1	B	264/311 (84%)	-0.08	12 (4%) 38 23	28, 50, 92, 148	5 (1%)
1	C	260/311 (83%)	0.16	10 (3%) 44 27	41, 63, 99, 116	0
1	D	257/311 (82%)	-0.01	8 (3%) 51 34	33, 56, 88, 105	1 (0%)
All	All	1044/1244 (83%)	0.03	37 (3%) 47 30	17, 55, 94, 148	11 (1%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	492	TYR	5.5
1	A	474	PHE	5.3
1	C	501	SER	5.1
1	D	492	TYR	5.0
1	A	467	GLU	4.9
1	A	471	SER	4.8
1	C	451	GLY	4.6
1	C	625	PRO	3.6
1	C	624	GLN	3.5
1	A	501	SER	3.5
1	B	445	ILE	3.4
1	B	471	SER	3.3
1	B	474	PHE	3.3
1	C	626	GLY	3.2
1	D	501	SER	2.9
1	A	640	THR	2.8
1	C	639	CYS	2.7
1	A	445	ILE	2.6
1	D	474	PHE	2.6
1	D	626	GLY	2.5
1	C	471	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	475	GLY	2.5
1	B	469	ILE	2.5
1	B	446	LEU	2.5
1	B	492	TYR	2.4
1	A	449	ARG	2.3
1	B	472	GLY	2.3
1	D	642	TYR	2.2
1	B	473	SER	2.2
1	C	474	PHE	2.2
1	B	449	ARG	2.1
1	B	726	ILE	2.1
1	B	476	LYS	2.1
1	D	680	HIS	2.1
1	D	681	HIS	2.0
1	C	688	TYR	2.0
1	D	615	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CV4	B	801	29/29	0.92	0.13	46,50,84,107	0
2	CV4	A	801	29/29	0.95	0.09	37,43,51,52	0
2	CV4	C	801	29/29	0.95	0.10	55,66,75,76	0
2	CV4	D	801	29/29	0.97	0.07	41,49,54,58	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers

as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around CV4 B 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CV4 A 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around CV4 C 801:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around CV4 D 801:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.