



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:36 PM UTC

PDB ID : 3B6P / pdb_00003b6p
Title : Structure of TREX1 in complex with a nucleotide and inhibitor ions (sodium and zinc)
Authors : Brucet, M.; Querol-Audi, J.; Fita, I.; Celada, A.
Deposited on : 2007-10-29
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	247	
1	B	247	
1	C	247	
1	D	247	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7299 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Three prime repair exonuclease 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	217	Total	C	N	O	S	0	1	0
			1684	1066	295	313	10			
1	B	217	Total	C	N	O	S	0	1	0
			1684	1066	295	313	10			
1	C	217	Total	C	N	O	S	0	1	0
			1684	1066	295	313	10			
1	D	217	Total	C	N	O	S	0	1	0
			1684	1066	295	313	10			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q91XB0
A	0	LYS	-	expression tag	UNP Q91XB0
A	1	HIS	-	expression tag	UNP Q91XB0
A	2	HIS	-	expression tag	UNP Q91XB0
A	3	HIS	-	expression tag	UNP Q91XB0
A	4	HIS	-	expression tag	UNP Q91XB0
A	5	HIS	-	expression tag	UNP Q91XB0
A	6	HIS	-	expression tag	UNP Q91XB0
A	7	PRO	-	expression tag	UNP Q91XB0
A	8	MET	-	expression tag	UNP Q91XB0
B	-1	MET	-	expression tag	UNP Q91XB0
B	0	LYS	-	expression tag	UNP Q91XB0
B	1	HIS	-	expression tag	UNP Q91XB0
B	2	HIS	-	expression tag	UNP Q91XB0
B	3	HIS	-	expression tag	UNP Q91XB0
B	4	HIS	-	expression tag	UNP Q91XB0
B	5	HIS	-	expression tag	UNP Q91XB0
B	6	HIS	-	expression tag	UNP Q91XB0
B	7	PRO	-	expression tag	UNP Q91XB0
B	8	MET	-	expression tag	UNP Q91XB0
C	-1	MET	-	expression tag	UNP Q91XB0

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	0	LYS	-	expression tag	UNP Q91XB0
C	1	HIS	-	expression tag	UNP Q91XB0
C	2	HIS	-	expression tag	UNP Q91XB0
C	3	HIS	-	expression tag	UNP Q91XB0
C	4	HIS	-	expression tag	UNP Q91XB0
C	5	HIS	-	expression tag	UNP Q91XB0
C	6	HIS	-	expression tag	UNP Q91XB0
C	7	PRO	-	expression tag	UNP Q91XB0
C	8	MET	-	expression tag	UNP Q91XB0
D	-1	MET	-	expression tag	UNP Q91XB0
D	0	LYS	-	expression tag	UNP Q91XB0
D	1	HIS	-	expression tag	UNP Q91XB0
D	2	HIS	-	expression tag	UNP Q91XB0
D	3	HIS	-	expression tag	UNP Q91XB0
D	4	HIS	-	expression tag	UNP Q91XB0
D	5	HIS	-	expression tag	UNP Q91XB0
D	6	HIS	-	expression tag	UNP Q91XB0
D	7	PRO	-	expression tag	UNP Q91XB0
D	8	MET	-	expression tag	UNP Q91XB0

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	D	1	Total Zn 1 1	0	0

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Na 1 1	0	0
3	B	1	Total Na 1 1	0	0
3	C	1	Total Na 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Na	0	0
			1	1		


- Molecule 4 is THYMIDINE-5'-PHOSPHATE (CCD ID: TMP) (formula: $C_{10}H_{15}N_2O_8P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			21	10	2	8	1		
4	B	1	Total	C	N	O	P	0	0
			21	10	2	8	1		
4	C	1	Total	C	N	O	P	0	0
			21	10	2	8	1		
4	D	1	Total	C	N	O	P	0	0
			21	10	2	8	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	125	Total	O	0	0
			125	125		
5	B	103	Total	O	0	0
			103	103		
5	C	105	Total	O	0	0
			105	105		
5	D	138	Total	O	0	0
			138	138		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.77Å 81.50Å 92.54Å 90.00° 103.11° 90.00°	Depositor
Resolution (Å)	24.18 – 2.30 24.18 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.8 (24.18-2.30) 99.7 (24.18-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.62 (at 2.19Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.246 , 0.285 0.264 , 0.303	Depositor DCC
R_{free} test set	2424 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.082	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	7299	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.06 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.1256e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, ZN, TMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.41	0/1724	0.78	0/2350
1	B	0.41	0/1724	0.77	0/2350
1	C	0.40	0/1724	0.75	0/2350
1	D	0.41	0/1724	0.78	0/2350
All	All	0.41	0/6896	0.77	0/9400

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1684	0	1686	6	0
1	B	1684	0	1686	4	0
1	C	1684	0	1686	1	0
1	D	1684	0	1686	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	21	0	13	0	0
4	B	21	0	13	0	0
4	C	21	0	13	0	0
4	D	21	0	13	0	0
5	A	125	0	0	0	0
5	B	103	0	0	1	0
5	C	105	0	0	0	0
5	D	138	0	0	0	0
All	All	7299	0	6796	17	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (17) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:38:ALA:HB3	1:A:65:ASP:HB2	1.91	0.51
1:B:37:LEU:HD21	1:B:205:LEU:HD22	1.94	0.49
1:D:38:ALA:HB3	1:D:65:ASP:HB2	1.96	0.48
1:B:121:LEU:HD11	1:B:147:LEU:HD22	1.95	0.47
1:D:121:LEU:HD11	1:D:147:LEU:HD22	1.97	0.46
1:A:121:LEU:HD11	1:A:147:LEU:HD22	1.99	0.45
1:C:121:LEU:HD11	1:C:147:LEU:HD22	1.98	0.45
1:A:43:ALA:HB1	1:A:61:PRO:HB3	2.00	0.43
1:B:73:PRO:HD2	5:B:1081:HOH:O	2.18	0.43
1:B:14:LEU:HD21	1:B:219:VAL:HG11	2.00	0.43
1:A:93:GLU:HB3	1:D:49:ILE:HD12	2.01	0.42
1:A:53:HIS:HA	1:A:54:PRO:HA	1.90	0.42
1:A:24:LEU:HB3	1:A:25:PRO:HD2	2.01	0.42
1:D:21:ALA:HB3	1:D:129:TYR:OH	2.20	0.41
1:D:37:LEU:HD21	1:D:205:LEU:HD22	2.03	0.41
1:D:103:ASN:HA	1:D:106:ILE:HD12	2.01	0.41
1:D:43:ALA:HB1	1:D:61:PRO:HB3	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	214/247 (87%)	210 (98%)	4 (2%)	0	100	100
1	B	214/247 (87%)	207 (97%)	7 (3%)	0	100	100
1	C	214/247 (87%)	207 (97%)	6 (3%)	1 (0%)	24	31
1	D	214/247 (87%)	210 (98%)	4 (2%)	0	100	100
All	All	856/988 (87%)	834 (97%)	21 (2%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	50	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	187/212 (88%)	183 (98%)	4 (2%)	47	66
1	B	187/212 (88%)	183 (98%)	4 (2%)	47	66
1	C	187/212 (88%)	183 (98%)	4 (2%)	47	66
1	D	187/212 (88%)	184 (98%)	3 (2%)	55	73
All	All	748/848 (88%)	733 (98%)	15 (2%)	48	67

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	133	LEU
1	A	148	ASP
1	A	205	LEU
1	A	215	LEU
1	B	64	VAL
1	B	133	LEU
1	B	153	VAL
1	B	201	VAL
1	C	64	VAL
1	C	133	LEU
1	C	153	VAL
1	C	205	LEU
1	D	133	LEU
1	D	153	VAL
1	D	215	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	117	GLN
1	A	217	GLN
1	A	222	HIS
1	B	51	GLN
1	B	53	HIS
1	B	95	GLN
1	B	117	GLN
1	B	125	ASN
1	B	217	GLN
1	C	53	HIS
1	C	124	HIS
1	C	213	GLN
1	C	217	GLN
1	D	117	GLN
1	D	217	GLN
1	D	222	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TMP	B	1000	3,2	22,22,22	1.42	4 (18%)	32,33,33	1.96	8 (25%)
4	TMP	C	1000	3,2	22,22,22	1.43	4 (18%)	32,33,33	1.93	9 (28%)
4	TMP	D	1000	3,2	22,22,22	1.44	4 (18%)	32,33,33	1.95	7 (21%)
4	TMP	A	1000	3,2	22,22,22	1.44	4 (18%)	32,33,33	1.94	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TMP	B	1000	3,2	-	1/10/22/22	0/2/2/2
4	TMP	C	1000	3,2	-	0/10/22/22	0/2/2/2
4	TMP	D	1000	3,2	-	0/10/22/22	0/2/2/2
4	TMP	A	1000	3,2	-	0/10/22/22	0/2/2/2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1000	TMP	C6-C5	3.05	1.39	1.34
4	D	1000	TMP	C6-C5	3.03	1.39	1.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	1000	TMP	C6-C5	2.96	1.39	1.34
4	C	1000	TMP	C6-C5	2.95	1.39	1.34
4	A	1000	TMP	C4-C5	2.83	1.49	1.44
4	D	1000	TMP	C2-N1	2.83	1.42	1.38
4	C	1000	TMP	C4-C5	2.80	1.49	1.44
4	C	1000	TMP	C2-N1	2.79	1.42	1.38
4	A	1000	TMP	C2-N1	2.77	1.42	1.38
4	B	1000	TMP	C2-N1	2.77	1.42	1.38
4	D	1000	TMP	C4-C5	2.75	1.49	1.44
4	B	1000	TMP	C4-C5	2.69	1.49	1.44
4	A	1000	TMP	C4-N3	-2.37	1.34	1.38
4	C	1000	TMP	C4-N3	-2.36	1.34	1.38
4	D	1000	TMP	C4-N3	-2.34	1.34	1.38
4	B	1000	TMP	C4-N3	-2.31	1.34	1.38

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1000	TMP	N3-C2-N1	4.93	121.31	114.89
4	D	1000	TMP	N3-C2-N1	4.89	121.26	114.89
4	C	1000	TMP	N3-C2-N1	4.88	121.24	114.89
4	B	1000	TMP	N3-C2-N1	4.84	121.20	114.89
4	D	1000	TMP	C4-N3-C2	-4.76	121.11	127.34
4	B	1000	TMP	C4-N3-C2	-4.75	121.11	127.34
4	A	1000	TMP	C4-N3-C2	-4.75	121.12	127.34
4	C	1000	TMP	C4-N3-C2	-4.67	121.21	127.34
4	D	1000	TMP	C5-C4-N3	4.34	119.10	115.32
4	A	1000	TMP	C5-C4-N3	4.27	119.04	115.32
4	B	1000	TMP	C5-C4-N3	4.27	119.03	115.32
4	C	1000	TMP	C5-C4-N3	4.20	118.97	115.32
4	B	1000	TMP	O4-C4-C5	-3.53	120.88	124.92
4	C	1000	TMP	O4-C4-C5	-3.43	120.99	124.92
4	A	1000	TMP	O4-C4-C5	-3.39	121.03	124.92
4	D	1000	TMP	O4-C4-C5	-3.38	121.05	124.92
4	D	1000	TMP	C5-C6-N1	-2.88	120.18	123.31
4	A	1000	TMP	C5-C6-N1	-2.87	120.19	123.31
4	B	1000	TMP	C5-C6-N1	-2.81	120.26	123.31
4	C	1000	TMP	C5-C6-N1	-2.72	120.36	123.31
4	C	1000	TMP	C5M-C5-C4	2.59	121.55	118.78
4	B	1000	TMP	C5M-C5-C4	2.57	121.52	118.78
4	D	1000	TMP	C5M-C5-C4	2.55	121.51	118.78
4	A	1000	TMP	C5M-C5-C4	2.53	121.49	118.78

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1000	TMP	O4'-C1'-N1	2.49	112.28	107.86
4	D	1000	TMP	O4'-C1'-N1	2.43	112.17	107.86
4	B	1000	TMP	O4'-C1'-N1	2.36	112.05	107.86
4	C	1000	TMP	O4'-C1'-N1	2.35	112.04	107.86
4	B	1000	TMP	C2'-C1'-N1	-2.10	108.56	113.81
4	C	1000	TMP	C2'-C1'-N1	-2.03	108.74	113.81
4	C	1000	TMP	O2-C2-N1	-2.02	120.17	122.80

There are no chirality outliers.

All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	B	1000	TMP	C5'-O5'-P-O1P

There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand TMP B 1000



Ligand TMP C 1000





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/247 (87%)	1.28	38 (17%) 4 4	14, 28, 33, 39	1 (0%)
1	B	217/247 (87%)	1.12	30 (13%) 6 7	14, 28, 34, 37	1 (0%)
1	C	217/247 (87%)	1.08	29 (13%) 7 8	14, 28, 36, 39	1 (0%)
1	D	217/247 (87%)	1.19	28 (12%) 7 8	14, 28, 34, 39	1 (0%)
All	All	868/988 (87%)	1.17	125 (14%) 6 7	14, 28, 34, 39	4 (0%)

All (125) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	49	ILE	6.9
1	B	50	SER	6.0
1	B	165	ALA	5.9
1	A	50	SER	5.9
1	A	49	ILE	5.7
1	C	47	THR	5.6
1	A	165	ALA	5.6
1	B	49	ILE	5.2
1	D	49	ILE	5.1
1	A	163	GLU	4.9
1	B	46	ASN	4.9
1	D	164	GLN	4.8
1	A	164	GLN	4.8
1	C	50	SER	4.7
1	A	10	HIS	4.6
1	B	48	SER	4.6
1	B	47	THR	4.5
1	D	51	GLN	4.5
1	D	10	HIS	4.4
1	B	148	ASP	4.4
1	A	59	ARG	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	48	SER	4.2
1	C	46	ASN	4.2
1	A	51	GLN	4.1
1	B	161	ALA	4.0
1	B	9	GLY	4.0
1	A	80	GLY	4.0
1	D	165	ALA	3.9
1	D	59	ARG	3.9
1	B	163	GLU	3.7
1	D	52	GLY	3.7
1	B	10	HIS	3.7
1	A	184	THR	3.6
1	D	80	GLY	3.6
1	A	148	ASP	3.5
1	B	234	GLY	3.5
1	D	76	ALA	3.4
1	C	9	GLY	3.4
1	B	162	LEU	3.4
1	B	164	GLN	3.4
1	C	10	HIS	3.4
1	D	184	THR	3.3
1	C	53	HIS	3.3
1	C	44	LEU	3.2
1	D	42	ARG	3.1
1	A	42	ARG	3.0
1	C	42	ARG	3.0
1	D	163	GLU	3.0
1	D	34	LEU	3.0
1	D	142	SER	2.9
1	C	164	GLN	2.9
1	C	163	GLU	2.9
1	B	184	THR	2.9
1	D	148	ASP	2.8
1	A	157	ALA	2.8
1	B	59	ARG	2.8
1	C	79	PRO	2.8
1	B	44	LEU	2.8
1	C	165	ALA	2.7
1	C	80	GLY	2.7
1	D	50	SER	2.7
1	C	159	LEU	2.7
1	D	157	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	9	GLY	2.6
1	A	52	GLY	2.6
1	C	177	TYR	2.5
1	A	175	LYS	2.5
1	A	53	HIS	2.5
1	A	199	GLY	2.5
1	D	156	ILE	2.4
1	A	234	GLY	2.4
1	B	83	GLU	2.4
1	C	143	THR	2.4
1	D	37	LEU	2.4
1	B	213	GLN	2.4
1	C	59	ARG	2.4
1	D	81	ALA	2.4
1	A	162	LEU	2.4
1	C	157	ALA	2.4
1	A	47	THR	2.4
1	D	64	VAL	2.4
1	B	79	PRO	2.3
1	A	202	LEU	2.3
1	D	196	THR	2.3
1	B	53	HIS	2.3
1	C	148	ASP	2.3
1	B	210	TRP	2.3
1	C	184	THR	2.3
1	A	64	VAL	2.3
1	A	34	LEU	2.3
1	A	143	THR	2.3
1	A	26	SER	2.3
1	C	26	SER	2.2
1	C	34	LEU	2.2
1	A	140	ARG	2.2
1	B	222	HIS	2.2
1	A	217	GLN	2.2
1	B	51	GLN	2.2
1	B	217	GLN	2.2
1	A	188	TRP	2.2
1	B	119[A]	CYS	2.2
1	D	77	CYS	2.2
1	A	48	SER	2.2
1	A	57	VAL	2.1
1	C	45	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	188	TRP	2.1
1	A	76	ALA	2.1
1	A	142	SER	2.1
1	D	143	THR	2.1
1	A	156	ILE	2.1
1	B	220	ASP	2.1
1	D	46	ASN	2.1
1	B	175	LYS	2.1
1	A	177	TYR	2.1
1	C	188	TRP	2.1
1	A	194	SER	2.0
1	A	93	GLU	2.0
1	A	179	LEU	2.0
1	C	220	ASP	2.0
1	A	39	VAL	2.0
1	B	128	ARG	2.0
1	D	180	GLY	2.0
1	C	210	TRP	2.0
1	D	217	GLN	2.0
1	C	99	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NA	C	801	1/1	0.81	0.10	31,31,31,31	0
3	NA	A	801	1/1	0.85	0.13	22,22,22,22	0
3	NA	B	801	1/1	0.87	0.09	25,25,25,25	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	TMP	A	1000	21/21	0.88	0.11	22,23,24,24	0
3	NA	D	801	1/1	0.89	0.16	25,25,25,25	0
4	TMP	B	1000	21/21	0.90	0.10	23,24,24,24	0
4	TMP	D	1000	21/21	0.90	0.11	25,25,26,27	0
4	TMP	C	1000	21/21	0.91	0.10	26,27,27,27	0
2	ZN	A	800	1/1	0.99	0.02	28,28,28,28	0
2	ZN	B	800	1/1	0.99	0.02	25,25,25,25	0
2	ZN	C	800	1/1	0.99	0.04	27,27,27,27	0
2	ZN	D	800	1/1	0.99	0.03	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around TMP A 1000:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TMP B 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TMP D 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around TMP C 1000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.