



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 11:33 AM UTC

PDB ID : 6B67 / pdb_00006b67
Title : Human PP2Calpha (PPM1A) complexed with cyclic peptide c(MpSIpYVA)
Authors : Dyda, F.; Kosek, D.
Deposited on : 2017-10-01
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	296	86% 14%
1	B	296	89% 10%
1	C	296	83% 16%
2	D	7	71% 29%
2	E	7	71% 29%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	7	29% 71% 29%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7726 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein phosphatase 1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	296	Total	C	N	O	S	0	0	0
			2297	1424	410	445	18			
1	B	296	Total	C	N	O	S	0	0	0
			2297	1424	410	445	18			
1	C	296	Total	C	N	O	S	0	0	0
			2297	1424	410	445	18			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	146	GLU	ASP	engineered mutation	UNP P35813
B	146	GLU	ASP	engineered mutation	UNP P35813
C	146	GLU	ASP	engineered mutation	UNP P35813

- Molecule 2 is a protein called cyclic peptide c(MpSIpYVA).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	7	Total	C	N	O	P	S	0	0
			64	36	8	16	2	2		
2	E	7	Total	C	N	O	P	S	0	0
			64	36	8	16	2	2		
2	F	7	Total	C	N	O	P	S	0	0
			64	36	8	16	2	2		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Ca	0	0
			4	4		
3	B	3	Total	Ca	0	0
			3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	3	Total 3	Ca 3	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	220	Total 220	O 220	0	0
4	B	202	Total 202	O 202	0	0
4	C	189	Total 189	O 189	0	0
4	D	7	Total 7	O 7	0	0
4	E	13	Total 13	O 13	0	0
4	F	2	Total 2	O 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Protein phosphatase 1A




- Molecule 1: Protein phosphatase 1A



- Molecule 1: Protein phosphatase 1A



- Molecule 2: cyclic peptide c(MpSIpYVA)




- Molecule 2: cyclic peptide c(MpSIpYVA)

Chain E:  71% 29%



- Molecule 2: cyclic peptide c(MpSIpYVA)

Chain F:  29% 71% 29%



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	159.03Å 89.57Å 70.05Å 90.00° 113.86° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 30.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.9 (30.00-2.20) 95.9 (30.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.03	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.19Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.169 , 0.229 0.164 , 0.221	Depositor DCC
R_{free} test set	1096 reflections (2.50%)	wwPDB-VP
Wilson B-factor (Å ²)	22.1	Xtriage
Anisotropy	0.335	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.024 for -h-2*k,l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7726	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SEP, PTR, CA, 48V

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2343	1.00	6/3160 (0.2%)
1	B	0.66	1/2343 (0.0%)	1.01	3/3160 (0.1%)
1	C	0.61	0/2343	0.99	11/3160 (0.3%)
2	D	0.93	0/25	0.75	0/29
2	E	0.58	0/25	0.88	0/29
2	F	0.35	0/25	0.54	0/29
All	All	0.64	1/7104 (0.0%)	1.00	20/9567 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	29	MET	SD-CE	-10.00	1.54	1.79

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	147	SER	N-CA-C	-8.84	98.14	110.35
1	A	147	SER	N-CA-C	-8.22	99.00	110.35
1	B	147	SER	N-CA-C	-7.37	101.02	110.53
1	A	283	ASN	N-CA-C	-7.34	100.84	110.53
1	C	283	ASN	N-CA-C	-6.66	101.29	110.35
1	C	157	VAL	N-CA-C	-6.45	100.32	108.84
1	A	197	LEU	N-CA-C	-6.45	100.07	110.32
1	B	283	ASN	N-CA-C	-6.44	101.59	110.35
1	C	13	HIS	N-CA-C	-6.10	97.20	107.99
1	C	240	GLY	N-CA-C	-6.02	106.87	114.16
1	C	66	GLN	N-CA-C	6.01	118.32	111.11
1	C	7	LYS	CA-C-N	5.86	125.80	119.76
1	C	7	LYS	C-N-CA	5.86	125.80	119.76
1	A	66	GLN	N-CA-C	5.74	117.21	111.07

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	51	GLU	N-CA-C	5.48	118.01	111.71
1	A	48	SER	N-CA-C	5.40	118.65	111.30
1	A	259	GLU	N-CA-C	-5.38	106.88	113.50
1	C	133	LEU	N-CA-C	-5.32	100.06	108.73
1	C	123	ASP	N-CA-C	-5.31	103.38	110.55
1	B	197	LEU	N-CA-C	-5.05	103.05	110.52

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2297	0	2211	25	0
1	B	2297	0	2211	23	0
1	C	2297	0	2211	32	0
2	D	64	0	53	0	0
2	E	64	0	53	0	0
2	F	64	0	53	0	0
3	A	4	0	0	0	0
3	B	3	0	0	0	0
3	C	3	0	0	0	0
4	A	220	0	0	8	0
4	B	202	0	0	1	0
4	C	189	0	0	7	0
4	D	7	0	0	0	0
4	E	13	0	0	0	0
4	F	2	0	0	0	0
All	All	7726	0	6792	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:252:ASP:HB3	4:A:604:HOH:O	1.80	0.81
1:A:148:ARG:HH11	1:A:247:ASN:HD21	1.31	0.77
1:B:10:MET:HA	1:B:29:MET:HE2	1.72	0.71
1:A:155:ARG:H	1:A:232:GLN:NE2	1.89	0.71
1:C:186:ARG:NH1	1:C:192:ALA:HB2	2.09	0.67
1:B:8:PRO:HG3	1:B:32:TRP:CE2	2.33	0.63
1:A:111:HIS:O	1:A:114:VAL:HG22	1.99	0.62
1:A:148:ARG:HH11	1:A:247:ASN:ND2	1.97	0.62
1:C:296:LYS:HA	4:C:405:HOH:O	1.99	0.62
1:A:229:GLU:HG2	4:A:560:HOH:O	2.00	0.60
1:B:10:MET:CA	1:B:29:MET:HE2	2.32	0.59
1:C:290:CYS:HB3	1:C:296:LYS:HE3	1.84	0.59
1:C:111:HIS:O	1:C:114:VAL:HG22	2.05	0.57
1:C:174:ARG:HD2	1:C:213:GLN:O	2.04	0.56
1:C:296:LYS:HB2	4:C:439:HOH:O	2.04	0.56
1:C:172:LYS:HZ2	1:C:176:GLN:HE22	1.54	0.56
1:B:113:ARG:O	1:B:117:GLU:HG2	2.06	0.55
1:C:119:LYS:HB2	1:C:122:ALA:HB2	1.89	0.55
1:A:155:ARG:H	1:A:232:GLN:HE21	1.53	0.55
1:B:174:ARG:HD3	1:B:213:GLN:O	2.06	0.55
1:A:46:LEU:HB2	1:A:50:LEU:HB2	1.89	0.54
1:C:124:ARG:HA	1:C:124:ARG:HE	1.72	0.54
1:B:10:MET:O	1:B:29:MET:HE2	2.07	0.53
1:C:163:ASP:HB2	1:C:165:LYS:HE2	1.91	0.53
1:A:146:GLU:OE1	1:A:165:LYS:HE2	2.10	0.52
1:B:10:MET:HE3	1:B:29:MET:HE1	1.91	0.51
1:C:290:CYS:CB	1:C:296:LYS:HE3	2.40	0.51
1:B:171:GLU:OE2	1:B:174:ARG:NH2	2.43	0.51
1:B:46:LEU:HB2	1:B:50:LEU:HB2	1.92	0.51
1:A:130:VAL:HG23	1:A:143:ASN:HD22	1.76	0.51
1:C:171:GLU:OE1	1:C:174:ARG:NH2	2.43	0.50
1:C:172:LYS:NZ	1:C:176:GLN:HE22	2.09	0.50
1:C:146:GLU:CD	1:C:165:LYS:HD3	2.36	0.50
1:B:160:PHE:CZ	1:C:169:PRO:HD3	2.47	0.49
1:C:23:ARG:NH2	1:C:262:ASP:O	2.46	0.49
1:C:297:VAL:HG11	4:C:445:HOH:O	2.13	0.49
1:A:148:ARG:NH1	1:A:247:ASN:HD21	2.06	0.49
1:B:264:LEU:HD22	1:B:288:LEU:HD13	1.94	0.48
1:C:46:LEU:HD22	1:C:76:LEU:HD21	1.95	0.48
1:C:262:ASP:HB3	4:C:425:HOH:O	2.14	0.47
1:C:184:ILE:HG22	4:C:464:HOH:O	2.14	0.46
1:C:255:ARG:HD2	4:C:551:HOH:O	2.14	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:262:ASP:HB3	4:B:560:HOH:O	2.14	0.46
1:C:183:MET:HE3	1:C:188:ASN:C	2.41	0.46
1:B:10:MET:SD	1:B:276:LEU:CD2	3.04	0.46
1:A:171:GLU:OE1	1:A:174:ARG:NH2	2.48	0.46
1:B:148:ARG:HB3	1:B:242:TRP:NE1	2.32	0.45
1:A:12:LYS:HE2	4:A:598:HOH:O	2.17	0.44
1:C:295:PRO:O	1:C:296:LYS:C	2.59	0.44
1:A:241:ILE:HG13	1:A:286:VAL:CG2	2.47	0.44
1:A:2:GLY:HA3	4:A:511:HOH:O	2.17	0.44
1:A:206:HIS:HB3	4:A:579:HOH:O	2.16	0.44
1:C:23:ARG:HE	1:C:296:LYS:HE2	1.83	0.44
1:A:71:CYS:SG	1:A:197:LEU:HD13	2.57	0.44
1:A:74:HIS:HE1	4:A:408:HOH:O	2.01	0.43
1:B:12:LYS:NZ	1:B:29:MET:HE3	2.33	0.43
1:C:59:TYR:CE2	1:C:72:CYS:HA	2.52	0.43
1:C:191:LEU:HG	1:C:193:VAL:HG12	2.00	0.43
1:B:10:MET:C	1:B:29:MET:HE2	2.43	0.43
1:C:295:PRO:O	1:C:297:VAL:HG23	2.17	0.43
1:C:23:ARG:HD3	1:C:296:LYS:HE2	2.00	0.43
1:A:12:LYS:HE3	4:A:495:HOH:O	2.18	0.43
1:A:46:LEU:HD22	1:A:76:LEU:HD21	2.01	0.43
1:A:33:ARG:NH1	4:A:402:HOH:O	2.51	0.42
1:A:174:ARG:HD2	1:A:213:GLN:O	2.19	0.42
1:B:10:MET:O	1:B:29:MET:CE	2.67	0.42
1:B:47:PRO:O	1:B:49:GLY:N	2.53	0.41
1:B:148:ARG:HB3	1:B:242:TRP:CE2	2.55	0.41
1:B:10:MET:HE3	1:B:29:MET:CE	2.49	0.41
1:B:66:GLN:HE21	1:B:119:LYS:NZ	2.18	0.41
1:A:26:LEU:C	1:A:26:LEU:HD12	2.46	0.41
1:A:71:CYS:SG	1:A:197:LEU:CD1	3.09	0.41
1:B:12:LYS:HZ2	1:B:29:MET:HE3	1.86	0.41
1:C:237:ALA:HA	1:C:285:SER:O	2.21	0.41
1:A:124:ARG:HD3	1:A:124:ARG:HA	1.88	0.40
1:C:150:LEU:C	1:C:150:LEU:HD23	2.47	0.40
1:C:172:LYS:HE2	4:C:415:HOH:O	2.21	0.40
1:C:181:SER:HB2	1:C:183:MET:HE2	2.04	0.40
1:B:12:LYS:CE	1:B:29:MET:HE3	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	294/296 (99%)	278 (95%)	14 (5%)	2 (1%)	18	19
1	B	294/296 (99%)	283 (96%)	9 (3%)	2 (1%)	18	19
1	C	294/296 (99%)	282 (96%)	10 (3%)	2 (1%)	18	19
2	D	2/7 (29%)	1 (50%)	1 (50%)	0	100	100
2	E	2/7 (29%)	2 (100%)	0	0	100	100
2	F	2/7 (29%)	2 (100%)	0	0	100	100
All	All	888/909 (98%)	848 (96%)	34 (4%)	6 (1%)	18	19

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	48	SER
1	A	88	SER
1	A	89	ALA
1	B	47	PRO
1	C	120	HIS
1	C	296	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	252/252 (100%)	245 (97%)	7 (3%)	38	52
1	B	252/252 (100%)	245 (97%)	7 (3%)	38	52

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	252/252 (100%)	248 (98%)	4 (2%)	55	71
2	D	3/3 (100%)	3 (100%)	0	100	100
2	E	3/3 (100%)	3 (100%)	0	100	100
2	F	3/3 (100%)	3 (100%)	0	100	100
All	All	765/765 (100%)	747 (98%)	18 (2%)	43	58

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	28	SER
1	A	43	VAL
1	A	72	CYS
1	A	170	LEU
1	A	190	SER
1	A	259	GLU
1	A	288	LEU
1	B	72	CYS
1	B	76	LEU
1	B	114	VAL
1	B	146	GLU
1	B	151	LEU
1	B	170	LEU
1	B	288	LEU
1	C	16	GLN
1	C	124	ARG
1	C	151	LEU
1	C	193	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	74	HIS
1	A	81	ASN
1	A	111	HIS
1	A	143	ASN
1	A	176	GLN
1	A	177	ASN
1	A	222	HIS
1	A	232	GLN
1	A	247	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	66	GLN
1	B	78	HIS
1	B	96	ASN
1	B	99	ASN
1	B	111	HIS
1	B	120	HIS
1	B	138	HIS
1	B	154	ASN
1	B	176	GLN
1	B	177	ASN
1	C	14	ASN
1	C	18	GLN
1	C	40	HIS
1	C	66	GLN
1	C	74	HIS
1	C	81	ASN
1	C	99	ASN
1	C	176	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

9 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PTR	E	4	2	15,16,17	1.38	1 (6%)	17,22,24	1.68	5 (29%)
2	PTR	D	4	2	15,16,17	0.95	0	17,22,24	1.23	2 (11%)
2	PTR	F	4	2	15,16,17	0.89	0	17,22,24	1.27	2 (11%)
2	48V	F	7	2	9,9,10	0.47	0	8,10,12	0.46	0
2	48V	D	7	2	9,9,10	0.48	0	8,10,12	0.52	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	48V	E	7	2	9,9,10	0.59	0	8,10,12	0.53	0
2	SEP	E	2	2,3	8,9,10	1.45	2 (25%)	7,12,14	2.62	6 (85%)
2	SEP	D	2	2,3	8,9,10	1.29	1 (12%)	7,12,14	2.81	4 (57%)
2	SEP	F	2	2	8,9,10	0.82	0	7,12,14	2.98	4 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PTR	E	4	2	-	3/10/11/13	0/1/1/1
2	PTR	D	4	2	-	2/10/11/13	0/1/1/1
2	PTR	F	4	2	-	1/10/11/13	0/1/1/1
2	48V	F	7	2	-	2/8/9/10	-
2	48V	D	7	2	-	2/8/9/10	-
2	48V	E	7	2	-	1/8/9/10	-
2	SEP	E	2	2,3	-	1/6/8/10	-
2	SEP	D	2	2,3	-	1/6/8/10	-
2	SEP	F	2	2	-	3/6/8/10	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	4	PTR	P-O1P	3.98	1.62	1.50
2	E	2	SEP	P-O1P	2.45	1.58	1.50
2	E	2	SEP	P-O3P	-2.24	1.46	1.54
2	D	2	SEP	P-O3P	2.12	1.62	1.54

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	2	SEP	OG-CB-CA	4.66	112.68	108.14
2	D	2	SEP	OG-CB-CA	4.64	112.66	108.14
2	E	2	SEP	O2P-P-OG	3.98	117.05	106.67
2	F	2	SEP	O2P-P-OG	3.82	116.63	106.67
2	D	2	SEP	OG-P-O1P	3.72	116.48	106.44
2	E	4	PTR	O3P-P-OH	3.63	116.06	105.32
2	F	4	PTR	O2P-P-OH	3.57	115.88	105.32
2	F	2	SEP	OG-P-O1P	3.55	116.04	106.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	4	PTR	O2P-P-OH	3.38	115.33	105.32
2	D	4	PTR	O2P-P-OH	3.29	115.05	105.32
2	D	2	SEP	O2P-P-OG	3.10	114.76	106.67
2	E	2	SEP	O3P-P-OG	2.92	114.30	106.67
2	E	2	SEP	O2P-P-O1P	-2.66	100.45	110.83
2	E	4	PTR	O3P-P-O1P	-2.61	100.65	110.83
2	E	2	SEP	OG-CB-CA	2.48	110.56	108.14
2	E	4	PTR	OH-P-O1P	-2.43	101.39	109.48
2	E	2	SEP	O3P-P-O1P	-2.36	101.63	110.83
2	D	4	PTR	O3P-P-O1P	-2.36	101.66	110.83
2	D	2	SEP	O3P-P-O1P	-2.26	102.02	110.83
2	F	4	PTR	O3P-P-O1P	-2.24	102.11	110.83
2	E	4	PTR	O3P-P-O2P	2.23	116.18	107.80
2	F	2	SEP	O3P-P-O1P	-2.17	102.39	110.83
2	E	2	SEP	O3P-P-O2P	2.09	115.65	107.80

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	2	SEP	CA-CB-OG-P
2	F	2	SEP	CB-OG-P-O2P
2	D	7	48V	N01-C02-C03-N05
2	E	7	48V	N01-C02-C03-N05
2	F	7	48V	N01-C02-C03-N05
2	D	4	PTR	N-CA-CB-CG
2	E	4	PTR	N-CA-CB-CG
2	E	2	SEP	CA-CB-OG-P
2	F	2	SEP	CA-CB-OG-P
2	D	4	PTR	C-CA-CB-CG
2	D	7	48V	N01-C02-C03-O04
2	F	2	SEP	CB-OG-P-O3P
2	E	4	PTR	CZ-OH-P-O1P
2	F	4	PTR	CZ-OH-P-O2P
2	F	7	48V	C02-C06-S07-C08
2	E	4	PTR	C-CA-CB-CG

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	296/296 (100%)	-0.74	4 (1%) 73 71	8, 17, 53, 108	0
1	B	296/296 (100%)	-0.82	1 (0%) 90 88	8, 18, 49, 78	0
1	C	296/296 (100%)	-0.64	5 (1%) 69 66	10, 22, 52, 97	0
2	D	4/7 (57%)	-0.53	0 100 100	21, 22, 24, 24	0
2	E	4/7 (57%)	-0.67	0 100 100	17, 21, 24, 27	0
2	F	4/7 (57%)	2.07	2 (50%) 0 0	72, 76, 96, 97	0
All	All	900/909 (99%)	-0.72	12 (1%) 75 73	8, 19, 53, 108	0

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	122	ALA	3.8
1	C	121	GLY	3.4
1	C	120	HIS	3.4
1	C	3	ALA	3.1
1	C	297	VAL	3.1
2	F	5	VAL	2.9
1	A	122	ALA	2.7
2	F	6	ALA	2.5
1	A	120	HIS	2.5
1	A	121	GLY	2.5
1	B	297	VAL	2.1
1	A	124	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column

labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PTR	F	4	16/17	0.59	0.15	88,106,116,116	0
2	48V	F	7	10/11	0.78	0.12	86,96,97,97	0
2	SEP	F	2	10/11	0.92	0.12	54,60,71,73	0
2	PTR	D	4	16/17	0.93	0.07	18,29,42,44	0
2	PTR	E	4	16/17	0.94	0.06	17,29,48,48	0
2	48V	E	7	10/11	0.95	0.06	25,27,29,31	0
2	SEP	E	2	10/11	0.96	0.07	22,23,26,26	0
2	SEP	D	2	10/11	0.98	0.05	16,18,20,20	0
2	48V	D	7	10/11	0.98	0.04	25,27,30,30	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	304	1/1	0.81	0.09	54,54,54,54	0
3	CA	C	303	1/1	0.93	0.05	36,36,36,36	0
3	CA	C	301	1/1	0.95	0.03	26,26,26,26	0
3	CA	B	301	1/1	0.95	0.04	18,18,18,18	0
3	CA	A	302	1/1	0.98	0.03	21,21,21,21	0
3	CA	B	302	1/1	0.98	0.04	26,26,26,26	0
3	CA	A	303	1/1	0.98	0.07	36,36,36,36	0
3	CA	C	302	1/1	0.98	0.03	33,33,33,33	0
3	CA	A	301	1/1	0.98	0.02	12,12,12,12	0
3	CA	B	303	1/1	0.99	0.02	35,35,35,35	0

6.5 Other polymers [i](#)

There are no such residues in this entry.