



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 09:58 AM UTC

PDB ID : 7B69 / pdb\_00007b69  
Title : BK Polyomavirus VP1 pentamer core(residues 26-299) mutant C104S  
Authors : Osipov, E.M.; Beelen, S.; Strelkov, S.V.  
Deposited on : 2020-12-07  
Resolution : 1.47 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	6779 (1.50-1.46)
Clashscore	190562	7025 (1.50-1.46)
Ramachandran outliers	187476	6917 (1.50-1.46)
Sidechain outliers	187428	6914 (1.50-1.46)
RSRZ outliers	180081	6781 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	275	 7% 92% 5%
1	BBB	275	 8% 90% 6%
1	CCC	275	 4% 90% 7%
1	DDD	275	 8% 88% 9%
1	EEE	275	 15% 85% 7% 8%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	FFF	275	 9% 89% 6% .
1	GGG	275	 8% 93% 7% .
1	HHH	275	 8% 87% 7% 5%
1	III	275	 10% 91% 7% ..
1	JJJ	275	 4% 91% 8% ..

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 22750 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Major capsid protein VP1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	AAA	262	Total 2045	C 1287	N 356	O 390	S 12	0	4	0
1	BBB	265	Total 2059	C 1298	N 357	O 391	S 13	0	4	0
1	CCC	268	Total 2077	C 1309	N 359	O 396	S 13	0	6	0
1	DDD	270	Total 2105	C 1323	N 367	O 401	S 14	0	6	0
1	EEE	253	Total 1960	C 1233	N 340	O 373	S 14	0	4	0
1	FFF	264	Total 2068	C 1299	N 361	O 395	S 13	0	5	0
1	GGG	273	Total 2109	C 1329	N 360	O 406	S 14	0	5	0
1	HHH	260	Total 2039	C 1284	N 354	O 387	S 14	0	7	0
1	III	273	Total 2105	C 1324	N 361	O 407	S 13	0	3	0
1	JJJ	272	Total 2113	C 1329	N 367	O 404	S 13	0	6	0

There are 20 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
AAA	25	GLY	-	expression tag	UNP P03088
AAA	104	SER	CYS	engineered mutation	UNP P03088
BBB	25	GLY	-	expression tag	UNP P03088
BBB	104	SER	CYS	engineered mutation	UNP P03088
CCC	25	GLY	-	expression tag	UNP P03088
CCC	104	SER	CYS	engineered mutation	UNP P03088
DDD	25	GLY	-	expression tag	UNP P03088
DDD	104	SER	CYS	engineered mutation	UNP P03088
EEE	25	GLY	-	expression tag	UNP P03088

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
EEE	104	SER	CYS	engineered mutation	UNP P03088
FFF	25	GLY	-	expression tag	UNP P03088
FFF	104	SER	CYS	engineered mutation	UNP P03088
GGG	25	GLY	-	expression tag	UNP P03088
GGG	104	SER	CYS	engineered mutation	UNP P03088
HHH	25	GLY	-	expression tag	UNP P03088
HHH	104	SER	CYS	engineered mutation	UNP P03088
III	25	GLY	-	expression tag	UNP P03088
III	104	SER	CYS	engineered mutation	UNP P03088
JJJ	25	GLY	-	expression tag	UNP P03088
JJJ	104	SER	CYS	engineered mutation	UNP P03088

- Molecule 2 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C<sub>2</sub>H<sub>6</sub>OS).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	AAA	1	Total	C	O	S	0	0
			4	2	1	1		
2	AAA	1	Total	C	O	S	0	0
			4	2	1	1		
2	BBB	1	Total	C	O	S	0	0
			4	2	1	1		
2	BBB	1	Total	C	O	S	0	0
			4	2	1	1		
2	CCC	1	Total	C	O	S	0	0
			4	2	1	1		
2	CCC	1	Total	C	O	S	0	0
			4	2	1	1		

Continued on next page...

*Continued from previous page...*

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	CCC	1	Total 4	C 2	O 1	S 1	0	0
2	DDD	1	Total 4	C 2	O 1	S 1	0	0
2	DDD	1	Total 4	C 2	O 1	S 1	0	0
2	EEE	1	Total 4	C 2	O 1	S 1	0	0
2	EEE	1	Total 4	C 2	O 1	S 1	0	0
2	EEE	1	Total 4	C 2	O 1	S 1	0	0
2	EEE	1	Total 4	C 2	O 1	S 1	0	0
2	FFF	1	Total 4	C 2	O 1	S 1	0	0
2	FFF	1	Total 4	C 2	O 1	S 1	0	0
2	GGG	1	Total 4	C 2	O 1	S 1	0	0
2	GGG	1	Total 4	C 2	O 1	S 1	0	0
2	GGG	1	Total 4	C 2	O 1	S 1	0	0
2	HHH	1	Total 4	C 2	O 1	S 1	0	0
2	HHH	1	Total 4	C 2	O 1	S 1	0	0
2	III	1	Total 4	C 2	O 1	S 1	0	0
2	III	1	Total 4	C 2	O 1	S 1	0	0
2	JJJ	1	Total 4	C 2	O 1	S 1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	AAA	183	Total 183	O 183	0	0
3	BBB	201	Total 201	O 201	0	0

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	CCC	244	Total 244	O 244	0	0
3	DDD	204	Total 204	O 204	0	0
3	EEE	179	Total 179	O 179	0	0
3	FFF	194	Total 194	O 194	0	0
3	GGG	184	Total 184	O 184	0	0
3	HHH	179	Total 179	O 179	0	0
3	III	205	Total 205	O 205	0	0
3	JJJ	205	Total 205	O 205	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Major capsid protein VP1

Chain AAA: 



- Molecule 1: Major capsid protein VP1

Chain BBB: 





- Molecule 1: Major capsid protein VP1

Chain CCC: 




- Molecule 1: Major capsid protein VP1

Chain DDD: 



- Molecule 1: Major capsid protein VP1





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.07Å 135.72Å 156.43Å 90.00° 94.98° 90.00°	Depositor
Resolution (Å)	102.35 – 1.47 102.35 – 1.47	Depositor EDS
% Data completeness (in resolution range)	96.6 (102.35-1.47) 96.6 (102.35-1.47)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.26 (at 1.47Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.181 , 0.197 (Not available) , 0.216	Depositor DCC
$R_{free}$ test set	20576 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	18.9	Xtrriage
Anisotropy	0.029	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 28.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	22750	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.10% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	AAA	0.88	0/2099	1.06	1/2848 (0.0%)
1	BBB	0.93	1/2117 (0.0%)	1.08	1/2875 (0.0%)
1	CCC	0.91	0/2141	1.05	1/2910 (0.0%)
1	DDD	0.94	1/2167 (0.0%)	1.09	3/2942 (0.1%)
1	EEE	0.89	0/2017	1.01	0/2738
1	FFF	0.91	0/2129	1.06	2/2889 (0.1%)
1	GGG	0.93	0/2171	1.04	0/2955
1	HHH	0.89	0/2104	1.03	0/2855
1	III	0.95	2/2160 (0.1%)	1.10	3/2937 (0.1%)
1	JJJ	0.91	0/2180	1.04	0/2961
All	All	0.92	4/21285 (0.0%)	1.06	11/28910 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	BBB	0	1
1	FFF	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	III	32	VAL	C-O	6.31	1.30	1.23
1	DDD	96	PRO	C-O	-5.81	1.17	1.23
1	III	100	GLU	C-O	5.51	1.30	1.23
1	BBB	49	CYS	C-N	-5.06	1.26	1.33

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	III	161	MET	CG-SD-CE	-10.67	77.43	100.90
1	FFF	161	MET	CG-SD-CE	-10.15	78.58	100.90
1	AAA	161	MET	CG-SD-CE	-10.12	78.63	100.90
1	DDD	100	GLU	CB-CA-C	-6.28	100.77	110.81
1	BBB	73	ASN	OD1-CG-ND2	-5.74	116.86	122.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	BBB	73	ASN	Sidechain
1	FFF	296	LYS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	2045	0	1995	4	0
1	BBB	2059	0	2019	10	0
1	CCC	2077	0	2026	17	0
1	DDD	2105	0	2062	17	0
1	EEE	1960	0	1906	10	0
1	FFF	2068	0	2029	11	0
1	GGG	2109	0	2033	11	0
1	HHH	2039	0	1994	12	0
1	III	2105	0	2030	16	0
1	JJJ	2113	0	2054	19	0
2	AAA	8	0	12	0	0
2	BBB	8	0	12	0	0
2	CCC	12	0	18	1	0
2	DDD	8	0	12	0	0
2	EEE	16	0	24	0	0
2	FFF	8	0	12	0	0
2	GGG	12	0	18	0	0
2	HHH	8	0	12	0	0
2	III	8	0	12	0	0
2	JJJ	4	0	6	0	0
3	AAA	183	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	BBB	201	0	0	2	1
3	CCC	244	0	0	6	0
3	DDD	204	0	0	4	1
3	EEE	179	0	0	1	0
3	FFF	194	0	0	2	0
3	GGG	184	0	0	1	0
3	HHH	179	0	0	3	0
3	III	205	0	0	4	0
3	JJJ	205	0	0	3	0
All	All	22750	0	20286	114	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 114 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:III:109:MET:CE	1:III:255:LYS:HA	2.04	0.87
1:FFF:211:ASP:OD2	1:FFF:214[B]:ARG:HD3	1.73	0.87
1:III:109:MET:HE3	1:III:255:LYS:HA	1.56	0.86
1:JJJ:211:ASP:OD2	1:JJJ:214[A]:ARG:HD3	1.75	0.85
1:DDD:211:ASP:OD2	1:DDD:214[B]:ARG:HD3	1.76	0.84

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:BBB:402:HOH:O	3:DDD:487:HOH:O[2_446]	2.06	0.14

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	260/275 (94%)	249 (96%)	10 (4%)	1 (0%)	30	11
1	BBB	265/275 (96%)	254 (96%)	10 (4%)	1 (0%)	30	11
1	CCC	270/275 (98%)	260 (96%)	9 (3%)	1 (0%)	30	11
1	DDD	274/275 (100%)	263 (96%)	10 (4%)	1 (0%)	30	11
1	EEE	251/275 (91%)	241 (96%)	9 (4%)	1 (0%)	30	11
1	FFF	265/275 (96%)	254 (96%)	10 (4%)	1 (0%)	30	11
1	GGG	276/275 (100%)	267 (97%)	8 (3%)	1 (0%)	30	11
1	HHH	261/275 (95%)	251 (96%)	9 (3%)	1 (0%)	30	11
1	III	274/275 (100%)	262 (96%)	11 (4%)	1 (0%)	30	11
1	JJJ	274/275 (100%)	263 (96%)	10 (4%)	1 (0%)	30	11
All	All	2670/2750 (97%)	2564 (96%)	96 (4%)	10 (0%)	30	11

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	188	VAL
1	BBB	188	VAL
1	CCC	188	VAL
1	DDD	188	VAL
1	EEE	188	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	225/235 (96%)	224 (100%)	1 (0%)	84	70
1	BBB	227/235 (97%)	225 (99%)	2 (1%)	70	48
1	CCC	228/235 (97%)	228 (100%)	0	100	100
1	DDD	232/235 (99%)	231 (100%)	1 (0%)	84	70
1	EEE	216/235 (92%)	215 (100%)	1 (0%)	81	65
1	FFF	230/235 (98%)	229 (100%)	1 (0%)	84	70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	GGG	231/235 (98%)	230 (100%)	1 (0%)	84	70
1	HHH	225/235 (96%)	223 (99%)	2 (1%)	70	48
1	III	229/235 (97%)	226 (99%)	3 (1%)	61	33
1	JJJ	232/235 (99%)	228 (98%)	4 (2%)	53	24
All	All	2275/2350 (97%)	2259 (99%)	16 (1%)	76	56

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	JJJ	98	LEU
1	JJJ	38	LYS
1	HHH	273	SER
1	JJJ	27	LEU
1	HHH	28	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	DMS	GGG	303	-	3,3,3	0.25	0	3,3,3	0.04	0
2	DMS	EEE	304	-	3,3,3	0.33	0	3,3,3	0.23	0
2	DMS	DDD	302	-	3,3,3	0.33	0	3,3,3	0.19	0
2	DMS	III	301	-	3,3,3	0.52	0	3,3,3	0.89	0
2	DMS	BBB	302	-	3,3,3	0.39	0	3,3,3	0.35	0
2	DMS	FFF	302	-	3,3,3	0.28	0	3,3,3	0.12	0
2	DMS	CCC	303	-	3,3,3	0.54	0	3,3,3	0.19	0
2	DMS	BBB	301	-	3,3,3	0.65	0	3,3,3	1.17	0
2	DMS	CCC	302	-	3,3,3	0.38	0	3,3,3	0.40	0
2	DMS	JJJ	301	-	3,3,3	0.45	0	3,3,3	1.02	0
2	DMS	CCC	301	-	3,3,3	0.67	0	3,3,3	1.49	1 (33%)
2	DMS	HHH	302	-	3,3,3	0.40	0	3,3,3	0.16	0
2	DMS	GGG	302	-	3,3,3	0.44	0	3,3,3	0.26	0
2	DMS	HHH	301	-	3,3,3	0.39	0	3,3,3	1.12	0
2	DMS	III	302	-	3,3,3	0.42	0	3,3,3	0.41	0
2	DMS	AAA	302	-	3,3,3	0.32	0	3,3,3	0.19	0
2	DMS	EEE	303	-	3,3,3	0.47	0	3,3,3	0.32	0
2	DMS	EEE	302	-	3,3,3	0.44	0	3,3,3	0.29	0
2	DMS	DDD	301	-	3,3,3	0.43	0	3,3,3	1.01	0
2	DMS	EEE	301	-	3,3,3	0.49	0	3,3,3	1.10	0
2	DMS	FFF	301	-	3,3,3	0.42	0	3,3,3	0.97	0
2	DMS	AAA	301	-	3,3,3	0.60	0	3,3,3	1.19	0
2	DMS	GGG	301	-	3,3,3	0.33	0	3,3,3	0.85	0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	CCC	301	DMS	O-S-C1	-2.57	93.64	106.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	CCC	303	DMS	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	262/275 (95%)	0.37	18 (6%) 23 25	8, 19, 44, 63	4 (1%)
1	BBB	265/275 (96%)	0.24	21 (7%) 18 20	9, 17, 44, 77	4 (1%)
1	CCC	268/275 (97%)	0.00	11 (4%) 41 45	8, 16, 33, 85	6 (2%)
1	DDD	270/275 (98%)	0.35	23 (8%) 16 18	9, 19, 40, 70	6 (2%)
1	EEE	253/275 (92%)	0.69	42 (16%) 4 4	10, 20, 71, 95	4 (1%)
1	FFF	264/275 (96%)	0.55	24 (9%) 15 16	10, 20, 47, 67	5 (1%)
1	GGG	273/275 (99%)	0.36	21 (7%) 19 21	9, 19, 45, 76	5 (1%)
1	HHH	260/275 (94%)	0.37	23 (8%) 15 17	10, 19, 48, 71	7 (2%)
1	III	273/275 (99%)	0.38	28 (10%) 12 12	9, 18, 42, 66	3 (1%)
1	JJJ	272/275 (98%)	0.30	12 (4%) 39 43	9, 20, 36, 79	6 (2%)
All	All	2660/2750 (96%)	0.36	223 (8%) 17 18	8, 19, 45, 95	50 (1%)

The worst 5 of 223 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	EEE	295	VAL	9.3
1	BBB	26	LEU	8.4
1	EEE	37	VAL	8.1
1	EEE	35	LEU	7.9
1	BBB	41	VAL	6.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	DMS	GGG	302	4/4	0.80	0.19	32,42,43,43	0
2	DMS	FFF	302	4/4	0.86	0.15	31,38,38,39	0
2	DMS	DDD	302	4/4	0.86	0.17	34,34,36,37	0
2	DMS	GGG	303	4/4	0.86	0.18	37,40,48,49	0
2	DMS	III	302	4/4	0.86	0.17	31,32,37,38	0
2	DMS	AAA	302	4/4	0.87	0.16	36,36,36,39	0
2	DMS	EEE	304	4/4	0.87	0.16	37,37,42,42	0
2	DMS	CCC	303	4/4	0.88	0.17	30,35,36,36	0
2	DMS	HHH	302	4/4	0.89	0.15	33,35,37,38	0
2	DMS	GGG	301	4/4	0.89	0.15	24,27,29,30	0
2	DMS	AAA	301	4/4	0.90	0.15	24,27,30,30	0
2	DMS	EEE	303	4/4	0.92	0.14	25,32,32,33	0
2	DMS	BBB	302	4/4	0.92	0.13	25,28,30,31	0
2	DMS	DDD	301	4/4	0.92	0.13	25,27,28,28	0
2	DMS	CCC	301	4/4	0.92	0.13	23,23,29,30	0
2	DMS	BBB	301	4/4	0.93	0.15	19,22,24,25	0
2	DMS	HHH	301	4/4	0.93	0.12	24,26,31,32	0
2	DMS	EEE	301	4/4	0.93	0.12	24,25,30,30	0
2	DMS	FFF	301	4/4	0.93	0.12	25,26,29,30	0
2	DMS	III	301	4/4	0.94	0.11	24,27,28,30	0
2	DMS	EEE	302	4/4	0.95	0.12	31,32,32,35	0
2	DMS	CCC	302	4/4	0.95	0.12	24,25,26,27	0
2	DMS	JJJ	301	4/4	0.95	0.11	22,24,27,27	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.