



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 03:26 PM UTC

PDB ID : 8B61 / pdb_00008b61
Title : Crystal structure of BfrC protein from Bacteroides fragilis NCTC 9343
Authors : Antonyuk, S.V.; Barnett, K.; Strange, R.W.; Olczak, T.
Deposited on : 2022-09-25
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.81 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1112 (1.82-1.82)
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)
RSRZ outliers	180081	1112 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	238	
1	B	238	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	301	-	-	X	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6761 atoms, of which 3017 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Conserved hypothetical lipoprotein.


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	197	Total	C	H	N	O	S	48	2	0
			3053	1001	1475	268	303	6			
1	B	200	Total	C	H	N	O	S	50	3	0
			3117	1025	1502	271	313	6			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	2	0
			14	3	8	3		
2	A	1	Total	C	H	O	2	0
			14	3	8	3		
2	B	1	Total	C	H	O	2	0
			14	3	8	3		

- Molecule 3 is pentane-1,3,5-tricarboxylic acid (CCD ID: P7W) (formula: $C_8H_{12}O_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			22	8	8	6		
3	B	1	Total	C	H	O	0	0
			22	8	8	6		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	246	Total	O	0	5
			247	247		
4	B	258	Total	O	0	0
			258	258		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Conserved hypothetical lipoprotein



• Molecule 1: Conserved hypothetical lipoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	52.08Å 89.84Å 99.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	66.55 – 1.81 66.55 – 1.81	Depositor EDS
% Data completeness (in resolution range)	99.2 (66.55-1.81) 99.3 (66.55-1.81)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.48 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.201 , 0.251 0.200 , 0.252	Depositor DCC
R_{free} test set	2219 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	21.6	Xtriage
Anisotropy	0.246	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 44.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6761	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 23.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.3464e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, P7W

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.14	2/1628 (0.1%)	1.32	4/2219 (0.2%)
1	B	1.17	2/1673 (0.1%)	1.33	4/2282 (0.2%)
All	All	1.16	4/3301 (0.1%)	1.33	8/4501 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	195	HIS	CE1-NE2	7.41	1.40	1.32
1	A	46	VAL	C-O	6.60	1.31	1.24
1	A	203	ASP	C-O	5.63	1.31	1.24
1	B	128	VAL	C-O	5.27	1.29	1.24

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	153	ASP	CA-CB-CG	6.34	118.94	112.60
1	B	174	ASN	CA-CB-CG	-5.26	107.34	112.60
1	B	32	ASP	CA-CB-CG	5.20	117.80	112.60
1	A	201	THR	CA-C-N	5.18	127.94	120.38
1	A	201	THR	C-N-CA	5.18	127.94	120.38
1	B	39	LYS	CA-C-N	5.17	130.68	122.62
1	B	39	LYS	C-N-CA	5.17	130.68	122.62
1	A	174	ASN	CA-CB-CG	-5.07	107.53	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1578	1475	1467	20	0
1	B	1615	1502	1500	18	0
2	A	12	16	16	6	0
2	B	6	8	8	1	0
3	A	14	8	0	1	0
3	B	14	8	0	0	0
4	A	247	0	0	6	0
4	B	258	0	0	6	0
All	All	3744	3017	2991	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:230:SER:HB3	4:B:454:HOH:O	1.87	0.75
2:A:301:GOL:O1	4:A:401:HOH:O	2.08	0.71
1:B:163:ASN:C	4:B:471:HOH:O	2.34	0.70
1:B:132[A]:GLN:HG2	1:B:172:ASN:ND2	2.07	0.69
1:A:42:TYR:HD1	2:A:301:GOL:H11	1.58	0.67
1:A:76:MET:O	1:A:99:ILE:HG23	1.95	0.66
1:B:163:ASN:CA	4:B:471:HOH:O	2.46	0.63
1:B:163:ASN:HA	4:B:471:HOH:O	1.98	0.61
1:A:237:GLN:OE1	1:B:68:ARG:NH1	2.27	0.60
1:B:52:TYR:HB2	1:B:61:PHE:HB2	1.88	0.55
1:B:67:ASN:OD1	1:B:71:LYS:HA	2.07	0.54
1:A:205[A]:LYS:NZ	4:A:405:HOH:O	2.42	0.51
1:A:132:GLN:HG2	4:A:487:HOH:O	2.12	0.49
1:B:132[A]:GLN:NE2	4:B:409:HOH:O	2.42	0.48
1:B:56:ARG:HD2	4:B:548:HOH:O	2.14	0.48
1:A:58:GLY:HA3	1:B:35[B]:THR:HG22	1.96	0.46
1:B:132[A]:GLN:HG2	1:B:172:ASN:HD21	1.79	0.46
3:A:303:P7W:O31	3:A:303:P7W:C21	2.63	0.46
1:A:31:PHE:CE2	1:A:230:SER:HB3	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:183:PHE:CD1	1:B:183:PHE:C	2.95	0.44
1:A:51:ILE:O	1:A:84:ALA:HA	2.18	0.44
1:A:94:SER:HB2	1:A:129:GLU:HG2	1.99	0.44
1:A:238:PRO:HG3	4:A:516:HOH:O	2.18	0.43
1:B:134:VAL:HB	1:B:173:ALA:HB3	2.00	0.43
1:A:87:GLY:HA3	2:A:301:GOL:H2	2.01	0.42
1:B:30:PRO:HD3	1:B:213:ILE:HG21	2.00	0.42
2:A:302:GOL:C3	4:A:427:HOH:O	2.67	0.42
1:A:67:ASN:OD1	1:A:71:LYS:HA	2.19	0.42
1:A:54:ASN:HB2	1:A:80:ASP:O	2.20	0.41
1:A:60:ILE:O	1:B:32:ASP:HA	2.20	0.41
1:A:154:PHE:CD1	1:A:154:PHE:C	2.97	0.41
1:A:42:TYR:CD1	2:A:301:GOL:H11	2.46	0.41
1:B:44:THR:HB	1:B:66:VAL:HG21	2.03	0.41
1:B:41:GLY:HA3	2:B:301:GOL:C3	2.51	0.41
1:A:41:GLY:HA3	2:A:301:GOL:H12	2.02	0.41
1:A:35:THR:HA	1:A:229:LEU:O	2.22	0.40
1:A:90:MET:N	4:A:416:HOH:O	2.53	0.40
1:A:111:ASN:HA	1:A:114:ASN:ND2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/238 (82%)	193 (100%)	1 (0%)	0	100	100
1	B	199/238 (84%)	198 (100%)	1 (0%)	0	100	100
All	All	393/476 (83%)	391 (100%)	2 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/201 (84%)	168 (100%)	0	100	100
1	B	173/201 (86%)	171 (99%)	2 (1%)	63	54
All	All	341/402 (85%)	339 (99%)	2 (1%)	78	74

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	163	ASN
1	B	182	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	ASN
1	A	114	ASN
1	A	145	HIS
1	A	172	ASN
1	B	172	ASN
1	B	211	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	B	301	-	5,5,5	0.23	0	5,5,5	0.50	0
3	P7W	B	302	-	13,13,13	1.09	2 (15%)	14,16,16	1.19	1 (7%)
2	GOL	A	301	-	5,5,5	0.21	0	5,5,5	0.84	0
3	P7W	A	303	-	13,13,13	1.23	2 (15%)	14,16,16	1.12	2 (14%)
2	GOL	A	302	-	5,5,5	0.17	0	5,5,5	0.38	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	301	-	-	2/4/4/4	-
3	P7W	B	302	-	-	6/14/14/14	-
2	GOL	A	301	-	-	2/4/4/4	-
3	P7W	A	303	-	-	5/14/14/14	-
2	GOL	A	302	-	-	2/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	303	P7W	O22-C2	-2.40	1.22	1.30
3	A	303	P7W	O42-C4	2.32	1.29	1.22
3	B	302	P7W	O31-C3	2.04	1.28	1.22
3	B	302	P7W	O42-C4	2.04	1.28	1.22

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	303	P7W	O31-C3-C31	-2.14	117.19	122.86
3	B	302	P7W	O21-C2-C21	-2.11	116.40	123.09
3	A	303	P7W	C22-C21-C2	-2.04	107.06	112.49

There are no chirality outliers.

All (17) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	301	GOL	O1-C1-C2-C3
2	B	301	GOL	O1-C1-C2-O2
2	B	301	GOL	O1-C1-C2-C3
3	A	303	P7W	O31-C3-C31-C32
3	A	303	P7W	O32-C3-C31-C32
3	B	302	P7W	O31-C3-C31-C32
3	B	302	P7W	O32-C3-C31-C32
3	B	302	P7W	C21-C22-C31-C3
2	A	301	GOL	O1-C1-C2-O2
3	A	303	P7W	C31-C32-C41-C4
3	B	302	P7W	C31-C32-C41-C4
2	A	302	GOL	C1-C2-C3-O3
3	B	302	P7W	O22-C2-C21-C22
3	A	303	P7W	O22-C2-C21-C22
3	A	303	P7W	O21-C2-C21-C22
3	B	302	P7W	O21-C2-C21-C22
2	A	302	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	301	GOL	1	0
2	A	301	GOL	5	0
3	A	303	P7W	1	0
2	A	302	GOL	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/238 (82%)	-0.08	5 (2%) 58 64	16, 26, 49, 78	5 (2%)
1	B	200/238 (84%)	-0.18	8 (4%) 42 48	12, 24, 44, 76	5 (2%)
All	All	397/476 (83%)	-0.13	13 (3%) 49 56	12, 25, 48, 78	10 (2%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	163	ASN	5.9
1	B	216	TYR	5.4
1	A	227	GLY	4.1
1	B	30	PRO	3.2
1	B	217	ASP	2.8
1	A	238	PRO	2.6
1	B	162	PRO	2.5
1	A	99	ILE	2.4
1	A	162	PRO	2.4
1	A	215	TRP	2.2
1	B	238	PRO	2.2
1	B	31	PHE	2.2
1	B	164	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	P7W	B	302	14/14	0.78	0.17	55,63,71,77	0
3	P7W	A	303	14/14	0.79	0.17	54,59,66,73	0
2	GOL	A	302	6/6	0.82	0.12	20,46,49,49	2
2	GOL	A	301	6/6	0.86	0.13	20,40,44,46	2
2	GOL	B	301	6/6	0.87	0.12	20,37,39,41	2

6.5 Other polymers [i](#)

There are no such residues in this entry.