



wwPDB EM Validation Summary Report ⓘ

Mar 23, 2026 – 05:23 AM UTC

PDB ID : 8B6G / pdb_00008b6g
EMDB ID : EMD-15866
Title : Cryo-EM structure of succinate dehydrogenase complex (complex-II) in respiratory supercomplex of *Tetrahymena thermophila*
Authors : Muhleip, A.; Kock Flygaard, R.; Baradaran, R.; Amunts, A.
Deposited on : 2022-09-27
Resolution : 3.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDb archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14081 (2.50 - 3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	CH	195	5% 50% 6% 44%
2	CM	76	88% 9% .
3	CL	89	10% 88% 11% .
4	CA	636	9% 82% 12% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	CI	114	
6	CB	312	
7	CF	296	
8	CG	198	
9	CK	93	
10	CE	322	
11	CJ	103	
12	CN	62	
13	CC	60	
14	CO	43	
15	CD	44	

2 Entry composition

There are 25 unique types of molecules in this entry. The entry contains 41412 atoms, of which 20810 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Diphthamide synthesis protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
1	CH	110	Total	C	H	N	O	S	0	0
			1700	529	845	147	171	8		

- Molecule 2 is a protein called Transmembrane protein, putative.

Mol	Chain	Residues	Atoms						AltConf	Trace
2	CM	74	Total	C	H	N	O	S	0	0
			1232	403	603	115	109	2		

- Molecule 3 is a protein called Transposase.

Mol	Chain	Residues	Atoms						AltConf	Trace
3	CL	88	Total	C	H	N	O	S	0	0
			1522	499	752	125	144	2		

- Molecule 4 is a protein called Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial.

Mol	Chain	Residues	Atoms						AltConf	Trace
4	CA	599	Total	C	H	N	O	S	0	0
			9198	2907	4574	825	866	26		

- Molecule 5 is a protein called DUF4885 domain-containing protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
5	CI	114	Total	C	H	N	O	S	0	0
			1805	580	890	153	180	2		

- Molecule 6 is a protein called Succinate dehydrogenase (quinone).

Mol	Chain	Residues	Atoms						AltConf	Trace
6	CB	285	Total	C	H	N	O	S	0	0
			4561	1457	2261	392	430	21		

- Molecule 7 is a protein called Transmembrane protein, putative.

Mol	Chain	Residues	Atoms						AltConf	Trace
7	CF	218	Total	C	H	N	O	S	0	0
			3598	1171	1786	306	331	4		

- Molecule 8 is a protein called SDHTT3.

Mol	Chain	Residues	Atoms						AltConf	Trace
8	CG	196	Total	C	H	N	O	S	0	0
			3247	1072	1593	273	305	4		

- Molecule 9 is a protein called Transmembrane protein, putative.

Mol	Chain	Residues	Atoms						AltConf	Trace
9	CK	93	Total	C	H	N	O	S	0	0
			1577	530	782	129	134	2		

- Molecule 10 is a protein called NmrA domain-containing protein.

Mol	Chain	Residues	Atoms						AltConf	Trace
10	CE	321	Total	C	H	N	O	S	0	0
			5115	1623	2554	449	488	1		

- Molecule 11 is a protein called Transmembrane protein, putative.

Mol	Chain	Residues	Atoms						AltConf	Trace
11	CJ	103	Total	C	H	N	O	S	0	0
			1663	554	815	140	151	3		

- Molecule 12 is a protein called Transmembrane protein, putative.

Mol	Chain	Residues	Atoms						AltConf	Trace
12	CN	62	Total	C	H	N	O	S	0	0
			1029	345	515	80	87	2		

- Molecule 13 is a protein called Cytochrome b-c1 complex subunit 8.

Mol	Chain	Residues	Atoms						AltConf	Trace
13	CC	59	Total	C	H	N	O	S	0	0
			976	319	487	86	83	1		


- Molecule 14 is a protein called SDHTT11.

Mol	Chain	Residues	Atoms						AltConf	Trace
14	CO	43	Total	C	H	N	O	S	0	0
			742	245	378	60	57	2		

- Molecule 15 is a protein called SDHD.


Mol	Chain	Residues	Atoms						AltConf	Trace
15	CD	44	Total	C	H	N	O	S	0	0
			807	271	412	61	62	1		

- Molecule 16 is CARDIOLIPIN (CCD ID: CDL) (formula: $C_{81}H_{156}O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
16	CM	1	Total	C	H	O	P	0
			256	81	156	17	2	
16	CG	1	Total	C	H	O	P	0
			256	81	156	17	2	
16	CG	1	Total	C	H	O	P	0
			256	81	156	17	2	
16	CK	1	Total	C	H	O	P	0
			256	81	156	17	2	
16	CJ	1	Total	C	H	O	P	0
			256	81	156	17	2	
16	CO	1	Total	C	H	O	P	0
			256	81	156	17	2	
16	CD	1	Total	C	H	O	P	0
			256	81	156	17	2	

- Molecule 18 is FLAVIN-ADENINE DINUCLEOTIDE (CCD ID: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf	
18	CA	1	Total	C	H	N	O	P	0
			84	27	31	9	15	2	

- Molecule 19 is FE2/S2 (INORGANIC) CLUSTER (CCD ID: FES) (formula: Fe_2S_2) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
19	CB	1	Total	Fe	S	0
			4	2	2	

- Molecule 20 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe_4S_4) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			AltConf
20	CB	1	Total	Fe	S	0
			8	4	4	

- Molecule 21 is FE3-S4 CLUSTER (CCD ID: F3S) (formula: Fe_3S_4) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
21	CB	1	Total	Fe	S	0
			7	3	4	

- Molecule 22 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (CCD ID: 3PE) (formula: $C_{41}H_{82}NO_8P$).




Mol	Chain	Residues	Atoms							AltConf
22	CK	1	Total 133	C 41	H 82	N 1	O 8	P 1	0	
22	CC	1	Total 133	C 41	H 82	N 1	O 8	P 1	0	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf	
22	CD	1	Total	C	H	N	O	P	0
			133	41	82	1	8	1	

- Molecule 23 is HEME C (CCD ID: HEC) (formula: $C_{34}H_{34}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						AltConf
23	CE	1	Total	C	Fe	H	N	O	0
			75	34	1	32	4	4	

- Molecule 24 is 1,2-DIACYL-SN-GLYCERO-3-PHOSPHOCHOLINE (CCD ID: PC1) (formula: $C_{44}H_{88}NO_8P$).



Mol	Chain	Residues	Atoms					AltConf	
24	CC	1	Total	C	H	N	O	P	0
			142	44	88	1	8	1	

- Molecule 25 is Ubiquinone-8 (CCD ID: UQ8) (formula: $C_{49}H_{74}O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
25	CC	1	Total	C	H	O	0
			127	49	74	4	


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Diphthamide synthesis protein




- Molecule 2: Transmembrane protein, putative




- Molecule 3: Transposase





- Molecule 4: Succinate dehydrogenase [ubiquinone] flavoprotein subunit, mitochondrial








• Molecule 5: DUF4885 domain-containing protein



• Molecule 6: Succinate dehydrogenase (quinone)




• Molecule 7: Transmembrane protein, putative




• Molecule 8: SDHTT3







- Molecule 9: Transmembrane protein, putative





- Molecule 10: NmrA domain-containing protein




- Molecule 11: Transmembrane protein, putative





- Molecule 12: Transmembrane protein, putative



- Molecule 13: Cytochrome b-c1 complex subunit 8



- Molecule 14: SDHTT11



● Molecule 15: SDHD



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	138746	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	25.66	Depositor
Minimum defocus (nm)	600	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	Not provided	
Image detector	GATAN K2 QUANTUM (4k x 4k)	Depositor
Maximum map value	6.278	Depositor
Minimum map value	-2.240	Depositor
Average map value	-0.004	Depositor
Map value standard deviation	0.109	Depositor
Recommended contour level	0.8	Depositor
Map size (Å)	600.0, 600.0, 600.0	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.25, 1.25, 1.25	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA, FAD, CDL, FES, PC1, UQ8, HEC, SF4, F3S, 3PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	CH	0.18	0/867	0.25	0/1166
2	CM	0.17	0/648	0.23	0/879
3	CL	0.18	0/788	0.24	0/1058
4	CA	0.16	0/4722	0.26	0/6385
5	CI	0.18	0/930	0.25	0/1244
6	CB	0.17	0/2355	0.25	0/3191
7	CF	0.18	0/1857	0.30	0/2512
8	CG	0.18	0/1699	0.27	0/2296
9	CK	0.20	0/821	0.28	0/1112
10	CE	0.17	0/2615	0.29	0/3553
11	CJ	0.18	0/875	0.24	0/1186
12	CN	0.17	0/530	0.28	0/719
13	CC	0.20	0/501	0.26	0/674
14	CO	0.20	0/375	0.24	0/508
15	CD	0.19	0/406	0.26	0/546
All	All	0.17	0/19989	0.27	0/27029

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	CH	855	845	844	9	0
2	CM	629	603	602	6	0
3	CL	770	752	752	7	0
4	CA	4624	4574	4573	48	0
5	CI	915	890	890	7	0
6	CB	2300	2261	2270	22	0
7	CF	1812	1786	1785	16	0
8	CG	1654	1593	1592	17	0
9	CK	795	782	782	11	0
10	CE	2561	2554	2553	24	0
11	CJ	848	815	815	8	0
12	CN	514	515	515	7	0
13	CC	489	487	487	4	0
14	CO	364	378	375	3	0
15	CD	395	412	412	2	0
16	CD	100	156	156	1	0
16	CG	200	312	312	1	0
16	CJ	100	156	156	2	0
16	CK	100	156	156	1	0
16	CM	100	156	156	1	0
16	CO	100	156	156	0	0
17	CA	1	0	0	0	0
17	CB	1	0	0	0	0
18	CA	53	31	29	7	0
19	CB	4	0	0	0	0
20	CB	8	0	0	1	0
21	CB	7	0	0	1	0
22	CC	51	82	82	1	0
22	CD	51	82	82	2	0
22	CK	51	82	82	0	0
23	CE	43	32	31	2	0
24	CC	54	88	88	1	0
25	CC	53	74	74	0	0
All	All	20602	20810	20807	162	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 162 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:CJ:49:ARG:NH1	16:CJ:201:CDL:OB7	2.08	0.86

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:CA:589:ARG:NH2	4:CA:632:ALA:O	2.12	0.82
16:CG:301:CDL:OB3	11:CJ:38:TYR:OH	1.99	0.80
8:CG:114:ARG:NH1	16:CD:301:CDL:OB4	2.18	0.77
7:CF:277:SER:HB3	9:CK:80:ILE:HD11	1.67	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	CH	108/195 (55%)	104 (96%)	4 (4%)	0	100	100
2	CM	72/76 (95%)	70 (97%)	2 (3%)	0	100	100
3	CL	86/89 (97%)	84 (98%)	2 (2%)	0	100	100
4	CA	597/636 (94%)	581 (97%)	15 (2%)	1 (0%)	43	76
5	CI	112/114 (98%)	111 (99%)	1 (1%)	0	100	100
6	CB	283/312 (91%)	272 (96%)	10 (4%)	1 (0%)	30	65
7	CF	216/296 (73%)	208 (96%)	8 (4%)	0	100	100
8	CG	194/198 (98%)	190 (98%)	4 (2%)	0	100	100
9	CK	91/93 (98%)	88 (97%)	3 (3%)	0	100	100
10	CE	319/322 (99%)	293 (92%)	23 (7%)	3 (1%)	14	48
11	CJ	101/103 (98%)	99 (98%)	2 (2%)	0	100	100
12	CN	60/62 (97%)	56 (93%)	4 (7%)	0	100	100
13	CC	57/60 (95%)	57 (100%)	0	0	100	100
14	CO	41/43 (95%)	41 (100%)	0	0	100	100
15	CD	42/44 (96%)	42 (100%)	0	0	100	100
All	All	2379/2643 (90%)	2296 (96%)	78 (3%)	5 (0%)	44	76

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	CA	405	TRP
10	CE	83	THR
10	CE	153	LEU
6	CB	266	ASP
10	CE	88	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	CH	100/184 (54%)	100 (100%)	0	100	100
2	CM	65/67 (97%)	65 (100%)	0	100	100
3	CL	82/83 (99%)	82 (100%)	0	100	100
4	CA	481/515 (93%)	479 (100%)	2 (0%)	84	90
5	CI	97/97 (100%)	97 (100%)	0	100	100
6	CB	259/283 (92%)	253 (98%)	6 (2%)	44	74
7	CF	194/268 (72%)	194 (100%)	0	100	100
8	CG	179/181 (99%)	177 (99%)	2 (1%)	65	83
9	CK	85/85 (100%)	84 (99%)	1 (1%)	63	82
10	CE	286/287 (100%)	283 (99%)	3 (1%)	68	84
11	CJ	86/86 (100%)	86 (100%)	0	100	100
12	CN	56/56 (100%)	56 (100%)	0	100	100
13	CC	50/51 (98%)	50 (100%)	0	100	100
14	CO	38/38 (100%)	38 (100%)	0	100	100
15	CD	43/43 (100%)	43 (100%)	0	100	100
All	All	2101/2324 (90%)	2087 (99%)	14 (1%)	73	86

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
6	CB	272	CYS
8	CG	99	HIS
10	CE	87	TYR
10	CE	3	LEU
10	CE	47	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
11	CJ	25	ASN
15	CD	25	ASN
15	CD	29	ASN
12	CN	44	ASN
6	CB	205	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 19 ligands modelled in this entry, 2 are monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
21	F3S	CB	1002	-	0,9,9	-	-	-		
16	CDL	CJ	201	-	99,99,99	0.29	0	105,111,111	0.26	0
16	CDL	CM	201	-	99,99,99	0.28	0	105,111,111	0.26	0
19	FES	CB	1000	-	0,4,4	-	-	-		
23	HEC	CE	401	10	46,50,50	1.82	5 (10%)	58,82,82	2.03	4 (6%)
16	CDL	CD	301	-	99,99,99	0.30	0	105,111,111	0.27	0
20	SF4	CB	1001	-	0,12,12	-	-	-		
16	CDL	CK	102	-	99,99,99	0.29	0	105,111,111	0.27	0
22	3PE	CC	301	-	50,50,50	0.26	0	53,55,55	0.21	0
18	FAD	CA	702	4	58,58,58	0.46	0	85,89,89	0.33	0
22	3PE	CD	302	-	50,50,50	0.27	0	53,55,55	0.20	0
24	PC1	CC	302	-	53,53,53	0.29	0	59,61,61	0.29	0
16	CDL	CG	302	-	99,99,99	0.30	0	105,111,111	0.25	0
16	CDL	CO	101	-	99,99,99	0.29	0	105,111,111	0.28	0
25	UQ8	CC	303	-	53,53,53	0.53	0	66,67,67	0.78	5 (7%)
22	3PE	CK	101	-	50,50,50	0.27	0	53,55,55	0.21	0
16	CDL	CG	301	-	99,99,99	0.29	0	105,111,111	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
21	F3S	CB	1002	-	-	-	0/3/3/3
16	CDL	CJ	201	-	-	17/110/110/110	-
16	CDL	CM	201	-	-	15/110/110/110	-
19	FES	CB	1000	-	-	-	0/1/1/1
23	HEC	CE	401	10	-	6/14/54/54	-
16	CDL	CD	301	-	-	32/110/110/110	-
20	SF4	CB	1001	-	-	-	0/6/5/5
16	CDL	CK	102	-	-	31/110/110/110	-
22	3PE	CC	301	-	-	5/54/54/54	-
18	FAD	CA	702	4	-	14/34/50/50	0/6/6/6
22	3PE	CD	302	-	-	13/54/54/54	-
24	PC1	CC	302	-	-	15/57/57/57	-
16	CDL	CG	302	-	-	26/110/110/110	-
16	CDL	CO	101	-	-	19/110/110/110	-
25	UQ8	CC	303	-	-	9/51/75/75	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
22	3PE	CK	101	-	-	4/54/54/54	-
16	CDL	CG	301	-	-	23/110/110/110	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
23	CE	401	HEC	CAC-C3C	6.09	1.54	1.35
23	CE	401	HEC	CAB-C3B	6.07	1.54	1.35
23	CE	401	HEC	C3D-C2D	5.53	1.53	1.38
23	CE	401	HEC	C3B-C2B	-2.21	1.33	1.41
23	CE	401	HEC	C3C-C2C	-2.10	1.34	1.41

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
23	CE	401	HEC	CBB-CAB-C3B	-9.28	108.89	127.43
23	CE	401	HEC	CBC-CAC-C3C	-9.23	108.99	127.43
23	CE	401	HEC	C4D-ND-C1D	3.31	111.22	105.82
25	CC	303	UQ8	C7-C8-C9	2.89	131.81	126.83
25	CC	303	UQ8	C11-C9-C8	-2.49	115.58	121.17

There are no chirality outliers.

5 of 229 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
16	CM	201	CDL	OB6-CB4-CB6-OB8
16	CG	301	CDL	CA2-OA2-PA1-OA3
16	CG	301	CDL	CA2-OA2-PA1-OA4
16	CG	301	CDL	CA2-OA2-PA1-OA5
16	CG	301	CDL	CB2-OB2-PB2-OB3

There are no ring outliers.

12 monomers are involved in 21 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
21	CB	1002	F3S	1	0
16	CJ	201	CDL	2	0
16	CM	201	CDL	1	0
23	CE	401	HEC	2	0
16	CD	301	CDL	1	0



Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
20	CB	1001	SF4	1	0
16	CK	102	CDL	1	0
22	CC	301	3PE	1	0
18	CA	702	FAD	7	0
22	CD	302	3PE	2	0
24	CC	302	PC1	1	0
16	CG	301	CDL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









Ligand HEC CE 401




Bond lengths



Bond angles




Torsions




Rings


Ligand CDL CD 301




Bond lengths




Bond angles






Torsions





Rings











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-15866. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 240




Y Index: 240




Z Index: 240

6.2.2 Raw map



X Index: 240



Y Index: 240




Z Index: 240


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)

6.3.1 Primary map



X Index: 193




Y Index: 331




Z Index: 210


6.3.2 Raw map



X Index: 0



Y Index: 0




Z Index: 0


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.8. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_15866_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 114 nm³; this corresponds to an approximate mass of 103 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.333 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.00	-	-
Author-provided FSC curve	2.96	3.50	3.00
Unmasked-calculated*	8.57	21.41	9.35

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 8.57 differs from the reported value 3.0 by more than 10 %

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-15866 and PDB model 8B6G. Per-residue inclusion information can be found in section [3](#) on page [12](#).

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.8 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.8).

9.4 Atom inclusion ⓘ



At the recommended contour level, 65% of all backbone atoms, 65% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.8) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6500	 0.4020
CA	 0.6540	 0.4150
CB	 0.7150	 0.4140
CC	 0.6250	 0.4050
CD	 0.6390	 0.3790
CE	 0.6420	 0.3920
CF	 0.6350	 0.4080
CG	 0.6770	 0.4080
CH	 0.6430	 0.3620
CI	 0.6070	 0.3840
CJ	 0.6690	 0.3930
CK	 0.6530	 0.3860
CL	 0.6570	 0.4000
CM	 0.6930	 0.3980
CN	 0.6220	 0.3890
CO	 0.6330	 0.4080

