



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 18, 2026 – 04:56 AM UTC

PDB ID : 4B7S / pdb_00004b7s
Title : PikC D50N mutant bound to the 10-DML analog with the 3-(N,N- dimethylamino)propanoate anchoring group
Authors : Podust, L.M.
Deposited on : 2012-08-21
Resolution : 1.84 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.84 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1296 (1.84-1.84)
Clashscore	190562	1329 (1.84-1.84)
Ramachandran outliers	187476	1318 (1.84-1.84)
Sidechain outliers	187428	1318 (1.84-1.84)
RSRZ outliers	180081	1296 (1.84-1.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	% 74% 16% • 9%
1	B	436	% 77% 11% • 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	A	1417	-	-	X	-
5	GOL	A	1420	-	-	X	-

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 7135 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYTOCHROME P450 HYDROXYLASE PIKC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	397	Total	C	N	O	S	0	3	0
			3084	1948	553	570	13			
1	B	393	Total	C	N	O	S	0	5	0
			3100	1959	554	574	13			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	expression tag	UNP O87605
A	-18	GLY	-	expression tag	UNP O87605
A	-17	SER	-	expression tag	UNP O87605
A	-16	SER	-	expression tag	UNP O87605
A	-15	HIS	-	expression tag	UNP O87605
A	-14	HIS	-	expression tag	UNP O87605
A	-13	HIS	-	expression tag	UNP O87605
A	-12	HIS	-	expression tag	UNP O87605
A	-11	HIS	-	expression tag	UNP O87605
A	-10	HIS	-	expression tag	UNP O87605
A	-9	SER	-	expression tag	UNP O87605
A	-8	SER	-	expression tag	UNP O87605
A	-7	GLY	-	expression tag	UNP O87605
A	-6	LEU	-	expression tag	UNP O87605
A	-5	VAL	-	expression tag	UNP O87605
A	-4	PRO	-	expression tag	UNP O87605
A	-3	ARG	-	expression tag	UNP O87605
A	-2	GLY	-	expression tag	UNP O87605
A	-1	SER	-	expression tag	UNP O87605
A	0	HIS	-	expression tag	UNP O87605
A	50	ASN	ASP	engineered mutation	UNP O87605
B	-19	MET	-	expression tag	UNP O87605
B	-18	GLY	-	expression tag	UNP O87605
B	-17	SER	-	expression tag	UNP O87605
B	-16	SER	-	expression tag	UNP O87605

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-15	HIS	-	expression tag	UNP O87605
B	-14	HIS	-	expression tag	UNP O87605
B	-13	HIS	-	expression tag	UNP O87605
B	-12	HIS	-	expression tag	UNP O87605
B	-11	HIS	-	expression tag	UNP O87605
B	-10	HIS	-	expression tag	UNP O87605
B	-9	SER	-	expression tag	UNP O87605
B	-8	SER	-	expression tag	UNP O87605
B	-7	GLY	-	expression tag	UNP O87605
B	-6	LEU	-	expression tag	UNP O87605
B	-5	VAL	-	expression tag	UNP O87605
B	-4	PRO	-	expression tag	UNP O87605
B	-3	ARG	-	expression tag	UNP O87605
B	-2	GLY	-	expression tag	UNP O87605
B	-1	SER	-	expression tag	UNP O87605
B	0	HIS	-	expression tag	UNP O87605
B	50	ASN	ASP	engineered mutation	UNP O87605

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is [(3R,4S,5S,7R,9E,11R,12R)-12-ethyl-3,5,7,11-tetramethyl-2,8-bis(oxidanylidene)-1-oxacyclododec-9-en-4-yl] 3-(dimethylamino)propanoate (CCD ID: QLE) (formula: C₂₂H₃₇NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			28	22	1	5		
4	B	1	Total	C	N	O	0	0
			28	22	1	5		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		
5	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	403	Total	O	0	0
			403	403		
6	B	324	Total	O	0	0
			324	324		

i

- Molecule 1: CYTOCHROME P450 HYDROXYLASE PIKC



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	61.25Å 92.02Å 69.53Å 90.00° 90.48° 90.00°	Depositor
Resolution (Å)	61.24 – 1.84 61.24 – 1.84	Depositor EDS
% Data completeness (in resolution range)	94.4 (61.24-1.84) 94.4 (61.24-1.84)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.74 (at 1.84Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.166 , 0.224 0.167 , 0.225	Depositor DCC
R_{free} test set	3187 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.415	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.076 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7135	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, QLE, GOL, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.27	4/3157 (0.1%)	1.15	9/4312 (0.2%)
1	B	1.20	2/3175 (0.1%)	1.12	10/4336 (0.2%)
All	All	1.23	6/6332 (0.1%)	1.13	19/8648 (0.2%)

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	185	ALA	CA-CB	-5.83	1.44	1.53
1	A	105	LYS	N-CA	5.77	1.53	1.46
1	A	331	ASP	CA-C	5.66	1.57	1.53
1	B	107	VAL	CA-CB	5.65	1.60	1.54
1	B	163	VAL	CA-CB	5.32	1.59	1.53

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	179	VAL	N-CA-C	9.03	119.21	111.90
1	B	308	VAL	CA-C-N	-6.57	117.86	122.59
1	B	308	VAL	C-N-CA	-6.57	117.86	122.59
1	A	178	PHE	N-CA-C	6.30	120.80	113.18
1	B	179	VAL	CB-CA-C	-6.25	105.30	111.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3084	0	3037	72	0
1	B	3100	0	3049	58	0
2	A	43	0	30	1	0
2	B	43	0	30	4	0
3	A	30	0	0	2	0
3	B	10	0	0	0	0
4	A	28	0	37	5	0
4	B	28	0	37	4	0
5	A	36	0	48	22	0
5	B	6	0	8	0	0
6	A	403	0	0	10	0
6	B	324	0	0	6	0
All	All	7135	0	6276	139	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 139 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:287:GLU:CB	5:A:1417:GOL:H31	1.63	1.27
1:B:180[B]:PHE:CD1	1:B:180[B]:PHE:O	2.05	1.09
5:A:1420:GOL:H12	6:A:2048:HOH:O	1.51	1.09
1:A:15:LEU:HD13	5:A:1420:GOL:O3	1.64	0.96
1:A:287:GLU:CB	5:A:1417:GOL:C3	2.44	0.95

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	398/436 (91%)	388 (98%)	8 (2%)	2 (0%)	24	12
1	B	396/436 (91%)	388 (98%)	8 (2%)	0	100	100
All	All	794/872 (91%)	776 (98%)	16 (2%)	2 (0%)	36	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	384	PRO
1	A	11	SER

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/355 (90%)	309 (97%)	10 (3%)	35	18
1	B	323/355 (91%)	310 (96%)	13 (4%)	28	11
All	All	642/710 (90%)	619 (96%)	23 (4%)	32	13

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	150[B]	TRP
1	B	202	ILE
1	B	179	VAL
1	B	287	GLU
1	A	150	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	188	GLN
1	B	208	GLN
1	B	344	HIS
1	B	265	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	344	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	GOL	A	1415	-	5,5,5	0.49	0	5,5,5	0.92	0
3	SO4	A	1410	-	4,4,4	0.31	0	6,6,6	0.71	0
2	HEM	A	1407	1	50,50,50	2.14	13 (26%)	67,82,82	1.86	8 (11%)
3	SO4	A	1412	-	4,4,4	0.29	0	6,6,6	0.19	0
5	GOL	A	1419	-	5,5,5	0.44	0	5,5,5	0.76	0
5	GOL	A	1420	-	5,5,5	0.44	0	5,5,5	1.04	0
3	SO4	A	1409	-	4,4,4	0.24	0	6,6,6	0.59	0
5	GOL	B	1411	-	5,5,5	0.66	0	5,5,5	0.94	0
5	GOL	A	1417	-	5,5,5	0.57	0	5,5,5	1.20	0
5	GOL	A	1418	-	5,5,5	0.62	0	5,5,5	0.43	0
4	QLE	B	1408	-	28,28,28	1.38	2 (7%)	34,38,38	1.80	5 (14%)
5	GOL	A	1416	-	5,5,5	0.51	0	5,5,5	0.70	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	B	1407	1	50,50,50	1.91	10 (20%)	67,82,82	1.56	17 (25%)
4	QLE	A	1408	-	28,28,28	1.37	3 (10%)	34,38,38	1.69	5 (14%)
3	SO4	A	1413	-	4,4,4	0.27	0	6,6,6	0.11	0
3	SO4	B	1409	-	4,4,4	0.41	0	6,6,6	0.45	0
3	SO4	A	1414	-	4,4,4	0.39	0	6,6,6	0.36	0
3	SO4	A	1411	-	4,4,4	0.50	0	6,6,6	1.07	0
3	SO4	B	1410	-	4,4,4	0.22	0	6,6,6	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	1407	1	-	0/14/54/54	-
5	GOL	A	1415	-	-	0/4/4/4	-
4	QLE	A	1408	-	-	3/44/44/44	0/0/1/1
5	GOL	B	1411	-	-	1/4/4/4	-
5	GOL	A	1417	-	-	4/4/4/4	-
2	HEM	A	1407	1	-	2/14/54/54	-
5	GOL	A	1419	-	-	1/4/4/4	-
5	GOL	A	1420	-	-	4/4/4/4	-
5	GOL	A	1418	-	-	0/4/4/4	-
4	QLE	B	1408	-	-	4/44/44/44	0/0/1/1
5	GOL	A	1416	-	-	0/4/4/4	-

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1407	HEM	C3D-C2D	7.25	1.52	1.36
2	B	1407	HEM	C3D-C2D	6.59	1.51	1.36
2	A	1407	HEM	FE-ND	5.89	2.13	1.94
2	B	1407	HEM	FE-NC	5.54	2.13	1.95
4	B	1408	QLE	O1-C8	5.13	1.46	1.34

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1407	HEM	C4D-ND-C1D	7.57	114.17	105.21
4	A	1408	QLE	C13-C14-N	-6.30	104.95	113.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1408	QLE	C13-C14-N	-5.67	105.81	113.50
2	A	1407	HEM	CHD-C1D-ND	5.15	129.96	124.42
4	B	1408	QLE	O1-C5-C6	4.38	113.87	106.90

There are no chirality outliers.

5 of 19 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1408	QLE	O1-C5-C6-C7
4	A	1408	QLE	C3-C5-C6-C7
5	A	1417	GOL	O1-C1-C2-O2
5	A	1417	GOL	O1-C1-C2-C3
5	A	1417	GOL	C1-C2-C3-O3


There are no ring outliers.


10 monomers are involved in 38 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1407	HEM	1	0
5	A	1419	GOL	3	0
5	A	1420	GOL	8	0
5	A	1417	GOL	8	0
4	B	1408	QLE	4	0
5	A	1416	GOL	3	0
2	B	1407	HEM	4	0
4	A	1408	QLE	5	0
3	A	1413	SO4	1	0
3	A	1411	SO4	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand HEM A 1407







Ligand HEM B 1407




Bond lengths




Bond angles



Torsions



Rings



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	397/436 (91%)	-0.51	3 (0%) 82 88	7, 18, 34, 49	3 (0%)
1	B	393/436 (90%)	-0.37	5 (1%) 75 82	12, 21, 35, 49	5 (1%)
All	All	790/872 (90%)	-0.44	8 (1%) 79 86	7, 19, 35, 49	8 (1%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	10	ALA	4.0
1	B	180[A]	PHE	3.0
1	B	150[A]	TRP	2.7
1	A	384	PRO	2.6
1	B	171	PHE	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	1411	5/5	0.70	0.16	51,54,56,59	0
3	SO4	A	1414	5/5	0.72	0.22	80,81,83,83	0
3	SO4	A	1409	5/5	0.78	0.15	62,64,65,67	0
3	SO4	A	1413	5/5	0.78	0.12	89,89,90,90	0
3	SO4	A	1410	5/5	0.78	0.12	63,63,65,66	0
5	GOL	A	1418	6/6	0.78	0.17	55,56,57,58	0
3	SO4	B	1409	5/5	0.79	0.12	63,64,65,66	0
3	SO4	B	1410	5/5	0.81	0.13	70,70,70,71	0
3	SO4	A	1412	5/5	0.83	0.12	82,82,82,82	0
5	GOL	A	1420	6/6	0.88	0.11	32,34,39,43	0
5	GOL	A	1419	6/6	0.89	0.12	32,42,45,45	0
5	GOL	A	1417	6/6	0.89	0.12	34,38,39,41	0
5	GOL	A	1416	6/6	0.91	0.11	32,40,42,43	0
4	QLE	B	1408	28/28	0.94	0.07	15,24,27,29	0
4	QLE	A	1408	28/28	0.94	0.08	17,24,31,32	0
5	GOL	A	1415	6/6	0.96	0.08	17,26,31,37	0
5	GOL	B	1411	6/6	0.96	0.06	19,27,31,39	0
2	HEM	A	1407	43/43	0.98	0.05	7,12,16,24	0
2	HEM	B	1407	43/43	0.99	0.04	9,13,16,20	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around QLE B 1408:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around QLE A 1408:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around HEM A 1407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around HEM B 1407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.