



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 09:14 PM UTC

PDB ID : 4B8C / pdb_00004b8c
Title : nuclease module of the yeast Ccr4-Not complex
Authors : Basquin, J.; Conti, E.
Deposited on : 2012-08-26
Resolution : 3.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1210 (3.48-3.36)
Clashscore	190562	1234 (3.48-3.36)
Ramachandran outliers	187476	1222 (3.48-3.36)
Sidechain outliers	187428	1222 (3.48-3.36)
RSRZ outliers	180081	1210 (3.48-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	2% 57% 28% 6% • 7%
1	C	288	2% 56% 30% 6% • 7%
1	E	288	3% 59% 27% 6% • 7%
1	F	288	2% 60% 26% 6% • 7%
2	B	249	2% 61% 32% • 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	G	249	5% 59% 33% • 6%
2	H	249	4% 59% 33% • 6%
2	I	249	4% 60% 31% • 6%
3	D	727	2% 16% 22% 6% 56%
3	J	727	2% 10% 13% 5% • 71%
3	K	727	2% 17% 21% 6% 56%
3	L	727	2% 10% 13% 6% • 71%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 24298 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POLY(A) RIBONUCLEASE POP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	267	Total	C	N	O	S	0	0	0
			2170	1409	346	404	11			
1	C	267	Total	C	N	O	S	0	0	0
			2174	1412	347	404	11			
1	E	267	Total	C	N	O	S	0	0	0
			2174	1411	347	405	11			
1	F	267	Total	C	N	O	S	0	0	0
			2170	1409	346	404	11			

- Molecule 2 is a protein called GENERAL NEGATIVE REGULATOR OF TRANSCRIPTION SUBUNIT 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	233	Total	C	N	O	S	0	0	0
			1869	1221	309	333	6			
2	G	233	Total	C	N	O	S	0	0	0
			1869	1221	309	333	6			
2	H	233	Total	C	N	O	S	0	0	0
			1869	1221	309	333	6			
2	I	233	Total	C	N	O	S	0	0	0
			1869	1221	309	333	6			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	752	ARG	-	expression tag	UNP P25655
B	753	SER	-	expression tag	UNP P25655
B	754	MET	-	expression tag	UNP P25655
G	752	ARG	-	expression tag	UNP P25655
G	753	SER	-	expression tag	UNP P25655
G	754	MET	-	expression tag	UNP P25655
H	752	ARG	-	expression tag	UNP P25655
H	753	SER	-	expression tag	UNP P25655

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	754	MET	-	expression tag	UNP P25655
I	752	ARG	-	expression tag	UNP P25655
I	753	SER	-	expression tag	UNP P25655
I	754	MET	-	expression tag	UNP P25655


- Molecule 3 is a protein called GLUCOSE-REPRESSIBLE ALCOHOL DEHYDROGENASE TRANSCRIPTIONAL EFFECTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	318	Total	C	N	O	S	0	0	0
			2456	1599	387	459	11			
3	J	210	Total	C	N	O	S	0	0	0
			1604	1034	266	298	6			
3	K	318	Total	C	N	O	S	0	0	0
			2470	1608	392	459	11			
3	L	210	Total	C	N	O	S	0	0	0
			1604	1034	266	298	6			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: POLY(A) RIBONUCLEASE POP2




• Molecule 1: POLY(A) RIBONUCLEASE POP2




• Molecule 1: POLY(A) RIBONUCLEASE POP2







• Molecule 1: POLY(A) RIBONUCLEASE POP2




• Molecule 2: GENERAL NEGATIVE REGULATOR OF TRANSCRIPTION SUBUNIT 1




• Molecule 2: GENERAL NEGATIVE REGULATOR OF TRANSCRIPTION SUBUNIT 1





● Molecule 3: GLUCOSE-REPRESSIBLE ALCOHOL DEHYDROGENASE TRANSCRIPTIONAL EFFECTOR



GLU VAL ASP PRO GLU TYR VAL SER LYS PHE ILE GLY PHE PRO ASN ASP LYS PHE PRO SER ASP HIS ILE PRO LEU LEU ALA ARG PHE GLU PHE MET LYS THR ASN THR GLY SER LYS VAL

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	122.65Å 122.91Å 126.42Å 89.47° 89.74° 64.22°	Depositor
Resolution (Å)	47.95 – 3.41 47.95 – 3.41	Depositor EDS
% Data completeness (in resolution range)	97.1 (47.95-3.41) 98.5 (47.95-3.41)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.29 (at 3.25Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: 1.7.1_743)	Depositor
R, R_{free}	0.233 , 0.270 0.227 , 0.265	Depositor DCC
R_{free} test set	4942 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	77.9	Xtriage
Anisotropy	0.333	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 70.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l 0.018 for k,h,-l 0.022 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	24298	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	0/2231	1.01	11/3028 (0.4%)
1	C	0.64	0/2235	1.01	9/3032 (0.3%)
1	E	0.64	0/2235	1.00	8/3033 (0.3%)
1	F	0.65	1/2231 (0.0%)	0.98	7/3028 (0.2%)
2	B	0.54	0/1908	0.92	2/2588 (0.1%)
2	G	0.53	0/1908	0.91	1/2588 (0.0%)
2	H	0.53	0/1908	0.92	2/2588 (0.1%)
2	I	0.53	0/1908	0.90	1/2588 (0.0%)
3	D	0.69	1/2520 (0.0%)	1.19	30/3446 (0.9%)
3	J	0.81	3/1640 (0.2%)	1.50	24/2240 (1.1%)
3	K	0.70	3/2535 (0.1%)	1.20	30/3463 (0.9%)
3	L	0.83	3/1640 (0.2%)	1.56	30/2240 (1.3%)
All	All	0.65	11/24899 (0.0%)	1.10	155/33862 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	1
1	E	0	1
3	D	0	1
3	J	0	1
3	K	0	1
3	L	0	2
All	All	0	8

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	L	462	ARG	C-N	7.32	1.50	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	J	462	ARG	C-N	7.11	1.50	1.33
3	L	332	ASP	CA-CB	6.55	1.66	1.53
3	L	361	ARG	CA-C	6.45	1.58	1.52
3	K	334	GLN	CA-C	5.79	1.60	1.52

The worst 5 of 155 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	462	ARG	O-C-N	27.71	145.96	121.30
3	J	462	ARG	O-C-N	27.41	145.70	121.30
3	L	465	ILE	CA-C-N	15.11	136.09	119.92
3	L	465	ILE	C-N-CA	15.11	136.09	119.92
3	J	397	ALA	N-CA-C	-12.97	94.10	110.19

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	194	THR	Peptide
1	C	194	THR	Peptide
3	D	344	LEU	Peptide
1	E	194	THR	Peptide
3	J	332	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2170	0	2083	99	0
1	C	2174	0	2094	102	0
1	E	2174	0	2089	94	1
1	F	2170	0	2083	81	1
2	B	1869	0	1936	56	0
2	G	1869	0	1936	55	1
2	H	1869	0	1936	64	0
2	I	1869	0	1936	56	1
3	D	2456	0	2268	232	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	J	1604	0	1505	150	1
3	K	2470	0	2293	211	0
3	L	1604	0	1505	164	1
All	All	24298	0	23664	1318	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 27.

The worst 5 of 1318 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:176:ILE:HD11	3:D:343:ASN:HD22	1.20	1.04
3:J:422:PHE:HD1	3:J:422:PHE:O	1.42	1.00
3:K:176:ILE:HD11	3:K:343:ASN:HD22	1.21	1.00
3:K:567:VAL:HG13	3:K:568:PRO:HD3	1.41	0.99
3:D:424:ASN:HD21	3:D:807:ILE:HG21	1.29	0.98

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:166:SER:OG	3:L:146:LEU:O[1_554]	2.08	0.12
2:G:835:TYR:OH	2:I:800:ASN:CA[1_465]	2.10	0.10
1:E:166:SER:OG	3:J:146:LEU:O[1_455]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	263/288 (91%)	250 (95%)	12 (5%)	1 (0%)	30 59
1	C	263/288 (91%)	250 (95%)	12 (5%)	1 (0%)	30 59

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	E	263/288 (91%)	249 (95%)	13 (5%)	1 (0%)	30	59
1	F	263/288 (91%)	250 (95%)	13 (5%)	0	100	100
2	B	231/249 (93%)	222 (96%)	9 (4%)	0	100	100
2	G	231/249 (93%)	222 (96%)	9 (4%)	0	100	100
2	H	231/249 (93%)	221 (96%)	10 (4%)	0	100	100
2	I	231/249 (93%)	222 (96%)	9 (4%)	0	100	100
3	D	300/727 (41%)	253 (84%)	46 (15%)	1 (0%)	36	65
3	J	204/727 (28%)	168 (82%)	34 (17%)	2 (1%)	12	39
3	K	300/727 (41%)	253 (84%)	46 (15%)	1 (0%)	36	65
3	L	204/727 (28%)	171 (84%)	31 (15%)	2 (1%)	12	39
All	All	2984/5056 (59%)	2731 (92%)	244 (8%)	9 (0%)	36	65

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	D	380	SER
3	K	380	SER
3	J	464	GLU
3	J	465	ILE
3	L	465	ILE

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/264 (91%)	215 (89%)	26 (11%)	6	22
1	C	242/264 (92%)	214 (88%)	28 (12%)	5	20
1	E	242/264 (92%)	215 (89%)	27 (11%)	6	21
1	F	241/264 (91%)	213 (88%)	28 (12%)	5	20
2	B	211/231 (91%)	200 (95%)	11 (5%)	21	47
2	G	211/231 (91%)	198 (94%)	13 (6%)	16	42

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	H	211/231 (91%)	198 (94%)	13 (6%)	16	42
2	I	211/231 (91%)	197 (93%)	14 (7%)	15	40
3	D	258/648 (40%)	206 (80%)	52 (20%)	1	4
3	J	165/648 (26%)	126 (76%)	39 (24%)	1	2
3	K	261/648 (40%)	207 (79%)	54 (21%)	1	4
3	L	165/648 (26%)	128 (78%)	37 (22%)	1	3
All	All	2659/4572 (58%)	2317 (87%)	342 (13%)	4	16

5 of 342 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	J	378	ASN
3	K	512	THR
3	J	398	GLU
3	K	341	LEU
3	K	779	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
3	J	427	ASN
3	L	175	ASN
3	K	142	HIS
3	K	424	ASN
3	L	389	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/288 (92%)	0.03	4 (1%) 72 58	36, 58, 98, 121	0
1	C	267/288 (92%)	-0.02	6 (2%) 62 47	36, 58, 99, 121	0
1	E	267/288 (92%)	0.09	9 (3%) 48 35	35, 58, 100, 121	0
1	F	267/288 (92%)	0.13	7 (2%) 57 42	36, 59, 98, 124	0
2	B	233/249 (93%)	0.19	4 (1%) 69 54	48, 79, 115, 146	0
2	G	233/249 (93%)	0.28	12 (5%) 33 24	49, 81, 112, 146	0
2	H	233/249 (93%)	0.28	10 (4%) 40 29	48, 80, 115, 145	0
2	I	233/249 (93%)	0.28	11 (4%) 36 27	50, 81, 114, 145	0
3	D	318/727 (43%)	0.42	11 (3%) 47 34	50, 87, 136, 160	0
3	J	210/727 (28%)	0.17	14 (6%) 24 19	35, 61, 101, 127	0
3	K	318/727 (43%)	0.44	18 (5%) 29 22	50, 87, 136, 160	0
3	L	210/727 (28%)	0.17	14 (6%) 24 19	35, 62, 102, 131	0
All	All	3056/5056 (60%)	0.21	120 (3%) 43 30	35, 72, 115, 160	0

The worst 5 of 120 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	991	THR	9.3
2	H	991	THR	8.4
3	J	252	ASP	4.8
2	I	773	SER	4.4
3	L	468	PRO	4.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.