



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:48 PM UTC

PDB ID : 3B98 / pdb_00003b98
Title : Crystal structure of zebrafish prostacyclin synthase (cytochrome P450 8A1)
Authors : Li, Y.-C.; Chiang, C.-W.; Yeh, H.-C.; Hsu, P.-Y.; Whitby, F.G.; Wang, L.-H.;
Chan, N.-L.
Deposited on : 2007-11-03
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8172 (2.10-2.06)
Clashscore	190562	8714 (2.10-2.06)
Ramachandran outliers	187476	8641 (2.10-2.06)
Sidechain outliers	187428	8642 (2.10-2.06)
RSRZ outliers	180081	8177 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	475	
1	B	475	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7443 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostaglandin I2 synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	446	Total	C	N	O	S	0	0	0
			3613	2300	641	653	19			
1	B	427	Total	C	N	O	S	0	0	0
			3462	2214	611	618	19			

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	MET	-	initiating methionine	UNP A9LLA5
A	11	ALA	-	expression tag	UNP A9LLA5
A	12	LYS	-	expression tag	UNP A9LLA5
A	13	LYS	-	expression tag	UNP A9LLA5
A	14	THR	-	expression tag	UNP A9LLA5
A	15	SER	-	expression tag	UNP A9LLA5
A	16	SER	-	expression tag	UNP A9LLA5
A	481	HIS	-	expression tag	UNP A9LLA5
A	482	HIS	-	expression tag	UNP A9LLA5
A	483	HIS	-	expression tag	UNP A9LLA5
A	484	HIS	-	expression tag	UNP A9LLA5
B	10	MET	-	initiating methionine	UNP A9LLA5
B	11	ALA	-	expression tag	UNP A9LLA5
B	12	LYS	-	expression tag	UNP A9LLA5
B	13	LYS	-	expression tag	UNP A9LLA5
B	14	THR	-	expression tag	UNP A9LLA5
B	15	SER	-	expression tag	UNP A9LLA5
B	16	SER	-	expression tag	UNP A9LLA5
B	481	HIS	-	expression tag	UNP A9LLA5
B	482	HIS	-	expression tag	UNP A9LLA5
B	483	HIS	-	expression tag	UNP A9LLA5
B	484	HIS	-	expression tag	UNP A9LLA5

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
2	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	198	Total	O	0	0
			198	198		
3	B	84	Total	O	0	0
			84	84		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Prostaglandin I2 synthase



• Molecule 1: Prostaglandin I2 synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	58.67Å 87.90Å 190.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.08 30.00 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.8 (30.00-2.08) 99.8 (30.00-2.08)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.58 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.227 , 0.264 (Not available) , 0.262	Depositor DCC
R_{free} test set	3031 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	27.0	Xtriage
Anisotropy	0.086	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 30.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7443	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/3695	0.78	0/4993
1	B	0.47	0/3539	0.81	0/4777
All	All	0.49	0/7234	0.80	0/9770

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3613	0	3614	32	0
1	B	3462	0	3472	26	0
2	A	43	0	30	2	0
2	B	43	0	30	2	0
3	A	198	0	0	4	0
3	B	84	0	0	0	0
All	All	7443	0	7146	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:MET:HE1	1:A:437:THR:HG21	1.48	0.95
1:A:146:MET:CE	1:A:437:THR:HG21	1.98	0.94
1:B:142:LEU:HG	1:B:146:MET:HE2	1.54	0.89
1:B:85:VAL:HG22	1:B:347:ILE:HG12	1.74	0.69
1:A:83:ASP:OD1	1:A:416:THR:HG23	1.94	0.67
2:B:600:HEM:HBB2	2:B:600:HEM:HHC	1.78	0.66
1:A:201:ASP:O	1:A:460:ARG:NH2	2.24	0.65
1:B:78:ASP:OD1	1:B:80:ASN:ND2	2.32	0.63
1:A:147:THR:HG22	1:A:150:GLU:H	1.64	0.62
1:B:378:HIS:HE1	1:B:410:PRO:O	1.83	0.62
1:A:146:MET:CE	3:A:711:HOH:O	2.49	0.60
1:A:210:THR:HG22	1:A:210:THR:O	2.01	0.60
1:A:149:SER:HA	1:A:154:LYS:HD2	1.84	0.60
1:A:211:THR:HG22	1:A:211:THR:O	2.01	0.59
1:A:146:MET:HE3	1:A:437:THR:HG21	1.83	0.58
1:B:138:MET:HE2	1:B:174:THR:HG21	1.86	0.57
1:A:146:MET:HE1	1:A:437:THR:CG2	2.31	0.55
1:B:300:ARG:NH2	1:B:439:LEU:O	2.40	0.55
1:B:33:LYS:O	1:B:67:ARG:HD2	2.08	0.53
1:A:146:MET:HE1	3:A:711:HOH:O	2.09	0.53
1:A:33:LYS:O	1:A:67:ARG:HD2	2.09	0.53
1:B:146:MET:CE	1:B:437:THR:HG21	2.40	0.52
1:A:378:HIS:HE1	1:A:410:PRO:O	1.93	0.52
2:A:600:HEM:HBB2	2:A:600:HEM:HMB2	1.91	0.51
1:B:448:ASP:HB2	1:B:451:ALA:HB2	1.92	0.51
2:A:600:HEM:HBB2	2:A:600:HEM:CMB	2.41	0.51
1:A:410:PRO:HG2	3:A:705:HOH:O	2.11	0.50
1:A:85:VAL:HG22	1:A:347:ILE:HG12	1.94	0.50
1:A:321:PHE:HD2	1:A:432:LYS:HZ3	1.58	0.49
1:B:284:TRP:CG	1:B:332:THR:HG21	2.49	0.48
1:B:125:HIS:CG	1:B:245:TRP:HB2	2.48	0.48
1:A:221:SER:O	1:A:225:GLU:HG3	2.14	0.47
1:A:142:LEU:HG	1:A:146:MET:HE2	1.95	0.47
1:A:206:LYS:HB3	1:A:212:VAL:HG13	1.95	0.47
1:A:158:TRP:CE3	1:A:475:ARG:HG3	2.49	0.47
1:A:211:THR:O	1:A:211:THR:CG2	2.61	0.47
1:B:324:VAL:HG22	1:B:386:PHE:HB2	1.97	0.47
1:B:296:LEU:O	1:B:300:ARG:HG3	2.14	0.47
1:A:171:LEU:HG	1:A:434:LEU:HD22	1.96	0.46
1:B:171:LEU:HG	1:B:434:LEU:HD22	1.97	0.46
1:B:80:ASN:HD21	1:B:350:SER:HB3	1.81	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:147:THR:HG22	1:B:149:SER:H	1.80	0.46
1:B:177:LEU:HD23	1:B:182:ALA:HB2	1.98	0.45
1:B:80:ASN:ND2	1:B:350:SER:HB3	2.31	0.45
1:A:153:LEU:HD13	1:A:478:ILE:HG13	1.98	0.45
1:A:321:PHE:CE2	1:A:432:LYS:HG2	2.51	0.44
1:A:416:THR:HG22	3:A:794:HOH:O	2.16	0.43
1:A:120:LYS:O	1:A:124:MET:HG2	2.18	0.43
1:B:85:VAL:HG13	1:B:347:ILE:HD11	1.99	0.43
1:A:140:ASN:OD1	1:A:143:ARG:NH2	2.45	0.43
1:A:239:LYS:HD3	1:A:240:PRO:HD2	2.01	0.43
1:A:417:GLU:O	1:A:420:LEU:HD22	2.19	0.42
1:B:146:MET:HE3	1:B:437:THR:HG21	2.01	0.42
1:B:349:LEU:HD22	1:B:355:TYR:CE1	2.53	0.42
1:B:125:HIS:CD2	1:B:134:LEU:HD11	2.55	0.42
1:B:427:ALA:HB2	2:B:600:HEM:HMC2	2.02	0.42
1:A:85:VAL:HG13	1:A:347:ILE:HD11	2.02	0.41
1:A:142:LEU:HD23	1:A:433:GLU:HG3	2.03	0.41
1:B:345:LYS:HE3	1:B:347:ILE:HD11	2.03	0.40
1:B:146:MET:HE1	1:B:437:THR:HG21	2.03	0.40
1:B:83:ASP:OD1	1:B:416:THR:HG23	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/475 (93%)	434 (98%)	8 (2%)	0	100	100
1	B	417/475 (88%)	414 (99%)	3 (1%)	0	100	100
All	All	859/950 (90%)	848 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	389/414 (94%)	374 (96%)	15 (4%)	28	29
1	B	371/414 (90%)	355 (96%)	16 (4%)	26	26
All	All	760/828 (92%)	729 (96%)	31 (4%)	27	27

All (31) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	43	LEU
1	A	67	ARG
1	A	76	LEU
1	A	77	LEU
1	A	120	LYS
1	A	134	LEU
1	A	147	THR
1	A	155	THR
1	A	347	ILE
1	A	358	ARG
1	A	395	THR
1	A	399	ASP
1	A	416	THR
1	A	420	LEU
1	A	472	LEU
1	B	66	VAL
1	B	76	LEU
1	B	80	ASN
1	B	107	ASN
1	B	120	LYS
1	B	246	LEU
1	B	273	VAL
1	B	340	ASP
1	B	347	ILE
1	B	358	ARG
1	B	382	GLU
1	B	416	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	420	LEU
1	B	453	VAL
1	B	460	ARG
1	B	465	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	139	GLN
1	A	184	ASN
1	A	185	ASN
1	A	192	GLN
1	A	376	GLN
1	A	378	HIS
1	A	385	GLN
1	A	403	ASN
1	A	425	HIS
1	B	80	ASN
1	B	317	ASN
1	B	343	GLN
1	B	376	GLN
1	B	378	HIS
1	B	385	GLN
1	B	419	ASN
1	B	425	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HEM	B	600	1	50,50,50	1.94	11 (22%)	67,82,82	1.27	5 (7%)
2	HEM	A	600	3,1	50,50,50	1.92	10 (20%)	67,82,82	1.24	3 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	B	600	1	-	2/14/54/54	-
2	HEM	A	600	3,1	-	3/14/54/54	-

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	600	HEM	C3D-C2D	8.02	1.54	1.36
2	A	600	HEM	C3D-C2D	7.70	1.53	1.36
2	A	600	HEM	FE-NC	4.40	2.09	1.95
2	B	600	HEM	FE-NA	4.36	2.09	1.95
2	A	600	HEM	FE-ND	4.31	2.08	1.94
2	B	600	HEM	FE-NB	4.17	2.07	1.94
2	A	600	HEM	FE-NB	3.86	2.06	1.94
2	B	600	HEM	FE-NC	3.36	2.06	1.95
2	B	600	HEM	CAC-C3C	3.25	1.56	1.47
2	B	600	HEM	FE-ND	3.15	2.04	1.94
2	A	600	HEM	CAC-C3C	3.07	1.55	1.47
2	B	600	HEM	CAB-C3B	2.99	1.55	1.47
2	A	600	HEM	FE-NA	2.84	2.04	1.95
2	A	600	HEM	CAB-C3B	2.75	1.54	1.47
2	A	600	HEM	CMB-C2B	2.51	1.55	1.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	600	HEM	CMD-C2D	2.24	1.55	1.50
2	B	600	HEM	CMB-C2B	2.22	1.55	1.50
2	B	600	HEM	CMA-C3A	2.19	1.55	1.50
2	A	600	HEM	CMC-C2C	2.14	1.55	1.50
2	B	600	HEM	CMC-C2C	2.12	1.55	1.50
2	A	600	HEM	CMD-C2D	2.08	1.55	1.50

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600	HEM	C4D-ND-C1D	5.40	111.61	105.21
2	B	600	HEM	C4D-ND-C1D	5.20	111.37	105.21
2	B	600	HEM	C3B-C4B-NB	-3.02	107.30	109.47
2	B	600	HEM	C3B-C2B-C1B	2.48	108.28	106.41
2	B	600	HEM	CHB-C1B-NB	2.43	127.37	124.37
2	A	600	HEM	C1B-NB-C4B	2.40	108.06	105.21
2	A	600	HEM	C3B-C4B-NB	-2.28	107.83	109.47
2	B	600	HEM	C1B-NB-C4B	2.23	107.85	105.21

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	600	HEM	C3D-CAD-CBD-CGD
2	A	600	HEM	C2A-CAA-CBA-CGA
2	B	600	HEM	C4B-C3B-CAB-CBB
2	A	600	HEM	CAA-CBA-CGA-O2A
2	A	600	HEM	CAA-CBA-CGA-O1A


There are no ring outliers.


2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	600	HEM	2	0
2	A	600	HEM	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	446/475 (93%)	0.03	15 (3%) 48 50	10, 21, 41, 55	0
1	B	427/475 (89%)	0.72	54 (12%) 8 8	15, 32, 57, 73	0
All	All	873/950 (91%)	0.37	69 (7%) 18 20	10, 26, 53, 73	0

All (69) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	248	SER	4.3
1	B	147	THR	4.2
1	B	182	ALA	3.7
1	B	231	LEU	3.7
1	B	154	LYS	3.7
1	B	187	SER	3.4
1	B	219	ILE	3.4
1	B	153	LEU	3.4
1	A	80	ASN	3.3
1	B	212	VAL	3.3
1	B	152	GLY	3.3
1	B	222	ALA	3.1
1	B	181	GLY	3.1
1	A	103	LYS	3.0
1	B	230	TRP	2.9
1	B	223	ALA	2.9
1	B	304	GLN	2.9
1	A	347	ILE	2.8
1	B	191	THR	2.8
1	B	348	CYS	2.8
1	B	220	ALA	2.7
1	B	211	THR	2.6
1	B	214	LYS	2.6
1	B	448	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	151	MET	2.6
1	B	80	ASN	2.6
1	A	90	ALA	2.6
1	B	192	GLN	2.5
1	B	317	ASN	2.5
1	B	441	ARG	2.5
1	A	395	THR	2.5
1	B	158	TRP	2.5
1	B	221	SER	2.5
1	A	392	ALA	2.5
1	B	260	ALA	2.5
1	B	452	THR	2.4
1	A	470	GLY	2.4
1	A	118	GLU	2.4
1	A	399	ASP	2.4
1	B	228	TRP	2.4
1	B	449	LYS	2.4
1	A	393	ASP	2.4
1	B	189	ALA	2.3
1	A	304	GLN	2.3
1	A	317	ASN	2.3
1	A	450	ASN	2.3
1	A	394	ARG	2.3
1	B	218	GLN	2.3
1	B	246	LEU	2.3
1	B	478	ILE	2.3
1	B	215	GLU	2.3
1	B	475	ARG	2.3
1	B	109	ILE	2.3
1	B	438	ILE	2.3
1	B	447	CYS	2.3
1	B	446	LEU	2.3
1	B	267	MET	2.2
1	B	35	MET	2.2
1	A	115	PRO	2.1
1	B	321	PHE	2.1
1	B	190	LEU	2.1
1	B	460	ARG	2.1
1	B	117	SER	2.1
1	B	294	GLU	2.1
1	B	473	GLU	2.1
1	B	247	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	303	ILE	2.1
1	B	474	ILE	2.1
1	B	440	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	HEM	B	600	43/43	0.95	0.11	20,22,37,39	0
2	HEM	A	600	43/43	0.96	0.09	10,15,33,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around HEM B 600:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.