



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 4, 2026 – 10:57 PM UTC

PDB ID : 4B91 / pdb_00004b91
Title : Crystal structure of truncated human CRMP-5
Authors : Ponnusamy, R.; Lohkamp, B.
Deposited on : 2012-08-31
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	484	2% 88% 10% .
1	B	484	4% 89% 8% .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8054 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DIHYDROPYRIMIDINASE-RELATED PROTEIN 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	474	Total	C	N	O	S	0	20	0
			3751	2374	649	701	27			
1	B	475	Total	C	N	O	S	0	14	0
			3715	2352	640	697	26			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9BPU6
B	0	SER	-	expression tag	UNP Q9BPU6


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	338	Total	O	0	2
			340	340		
2	B	244	Total	O	0	4
			248	248		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: DIHYDROPYRIMIDINASE-RELATED PROTEIN 5



• Molecule 1: DIHYDROPYRIMIDINASE-RELATED PROTEIN 5



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	90.09Å 90.09Å 248.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.60 – 1.70 39.60 – 1.70	Depositor EDS
% Data completeness (in resolution range)	97.6 (39.60-1.70) 97.6 (39.60-1.70)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.97 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.170 , 0.202 0.179 , 0.208	Depositor DCC
R_{free} test set	5532 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	22.0	Xtriage
Anisotropy	0.336	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 30.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8054	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.33	7/3871 (0.2%)	1.13	5/5252 (0.1%)
1	B	1.31	12/3831 (0.3%)	1.12	5/5198 (0.1%)
All	All	1.32	19/7702 (0.2%)	1.13	10/10450 (0.1%)

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	80	ASP	CA-C	-7.69	1.43	1.52
1	B	314[A]	LEU	C-O	-6.49	1.16	1.24
1	B	314[B]	LEU	C-O	-6.49	1.16	1.24
1	B	72[A]	THR	C-O	-5.93	1.16	1.24
1	B	72[B]	THR	C-O	-5.93	1.16	1.24
1	A	108	SER	N-CA	-5.70	1.39	1.46
1	A	152	LYS	CA-C	5.42	1.59	1.52
1	A	191	HIS	N-CA	-5.39	1.39	1.46
1	B	471	VAL	N-CA	-5.24	1.39	1.46
1	A	157	PHE	N-CA	5.23	1.52	1.45
1	B	256	VAL	C-O	5.23	1.30	1.24
1	A	12	ILE	CA-CB	5.20	1.60	1.53
1	B	372	ARG	C-O	5.18	1.30	1.24
1	B	12	ILE	CA-CB	5.14	1.59	1.53
1	A	194	ASN	C-O	-5.13	1.18	1.24
1	B	89	ALA	C-O	5.09	1.29	1.24
1	B	93	GLY	C-O	-5.08	1.17	1.24
1	B	169	LEU	C-O	5.07	1.30	1.23
1	A	203	LYS	N-CA	5.04	1.52	1.46

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	476	VAL	N-CA-C	-5.93	104.57	110.62
1	B	40	ARG	N-CA-C	5.92	118.62	111.40
1	B	222	GLU	N-CA-C	5.82	119.47	112.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	472	TYR	N-CA-C	5.68	119.94	112.89
1	B	474	LYS	CD-CE-NZ	5.40	129.18	111.90
1	A	471	VAL	N-CA-CB	-5.38	105.50	111.41
1	B	472	TYR	N-CA-C	5.32	119.48	112.89
1	B	234	THR	N-CA-C	-5.22	105.50	111.14
1	A	61	GLY	CA-C-N	5.13	124.84	119.92
1	A	61	GLY	C-N-CA	5.13	124.84	119.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3751	0	3784	27	0
1	B	3715	0	3749	16	0
2	A	340	0	0	1	0
2	B	248	0	0	0	0
All	All	8054	0	7533	43	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (43) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:430:MET:HE2	1:A:432:CYS:SG	1.95	1.05
1:B:159[B]:MET:HE1	1:B:188:ALA:HB1	1.69	0.74
1:A:9[B]:ARG:NH2	1:A:46:GLY:O	2.27	0.67
1:B:316[B]:ASN:OD1	1:B:318:THR:N	2.26	0.66
1:A:76:ALA:HB2	1:A:336:MET:HE1	1.82	0.60
1:A:297:VAL:HA	1:A:298:PRO:C	2.29	0.57
1:A:409[A]:ALA:O	1:A:433[A]:HIS:C	2.48	0.56
1:A:244:TYR:CE2	1:A:384:LEU:CD2	2.88	0.56
1:A:408[A]:GLU:OE1	1:A:408[A]:GLU:HA	2.05	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251:ILE:HA	1:A:309[A]:TYR:CE2	2.42	0.55
1:B:297:VAL:HA	1:B:298:PRO:C	2.31	0.55
1:A:244:TYR:CE2	1:A:384:LEU:HD22	2.43	0.53
1:B:409:ALA:HB2	1:B:435:VAL:HG22	1.92	0.51
1:A:409[A]:ALA:O	1:A:433[A]:HIS:O	2.27	0.51
1:A:244:TYR:HE2	1:A:384:LEU:HD22	1.76	0.51
1:B:251:ILE:HA	1:B:309[A]:TYR:CE2	2.47	0.50
1:B:244:TYR:CE2	1:B:384:LEU:CD2	2.94	0.49
1:A:170[A]:ARG:HD3	2:A:2157:HOH:O	2.12	0.49
1:A:178:LEU:HD12	1:A:235:ILE:HG22	1.93	0.49
1:B:244:TYR:CE2	1:B:384:LEU:HD22	2.47	0.49
1:B:190[A]:VAL:HG21	1:B:243:ILE:HD11	1.95	0.48
1:A:409[A]:ALA:HB3	1:A:435[A]:VAL:CG2	2.45	0.47
1:A:410:THR:HG21	1:A:431:ARG:HD3	1.97	0.46
1:A:9[B]:ARG:HD2	1:A:29:TYR:OH	2.17	0.44
1:A:428:GLU:O	1:A:429:ASN:HB2	2.17	0.44
1:A:406:ASP:HB2	1:A:437[A]:LEU:HG	1.98	0.44
1:B:178:LEU:HD12	1:B:235:ILE:HG22	2.00	0.44
1:A:410:THR:HG22	1:A:433[A]:HIS:CD2	2.53	0.43
1:A:181[B]:CYS:SG	1:A:186:ALA:HB3	2.57	0.43
1:B:98:ILE:HG21	1:B:131:HIS:CE1	2.54	0.43
1:B:244:TYR:HE2	1:B:384:LEU:HD22	1.83	0.43
1:A:313:LEU:HB3	1:A:318:THR:HB	2.01	0.42
1:A:437[A]:LEU:HD23	1:A:437[A]:LEU:HA	1.76	0.42
1:B:159[B]:MET:HE1	1:B:188:ALA:CB	2.45	0.42
1:A:190[A]:VAL:HG21	1:A:243:ILE:HD11	2.00	0.42
1:B:18:VAL:HB	1:B:59:ILE:HG22	2.01	0.42
1:A:410:THR:CG2	1:A:433[A]:HIS:CD2	3.03	0.41
1:A:309[A]:TYR:CZ	1:A:313:LEU:HD11	2.54	0.41
1:A:313:LEU:HA	1:A:316[A]:ASN:ND2	2.35	0.41
1:B:288:TRP:CZ2	1:B:338:LYS:HA	2.56	0.40
1:B:409:ALA:CB	1:B:435:VAL:HG22	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	492/484 (102%)	483 (98%)	8 (2%)	1 (0%)	43	28
1	B	487/484 (101%)	475 (98%)	11 (2%)	1 (0%)	43	28
All	All	979/968 (101%)	958 (98%)	19 (2%)	2 (0%)	43	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	350	GLY
1	B	350	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	410/400 (102%)	407 (99%)	3 (1%)	76	69
1	B	407/400 (102%)	400 (98%)	7 (2%)	53	38
All	All	817/800 (102%)	807 (99%)	10 (1%)	70	51

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	179[A]	HIS
1	A	179[B]	HIS
1	A	474	LYS
1	B	179[A]	HIS
1	B	179[B]	HIS
1	B	182	LYS
1	B	279	THR
1	B	349[A]	SER
1	B	349[B]	SER
1	B	474	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	474/484 (97%)	-0.11	11 (2%) 61 65	10, 20, 37, 78	20 (4%)
1	B	475/484 (98%)	0.10	17 (3%) 46 50	10, 23, 42, 73	14 (2%)
All	All	949/968 (98%)	-0.01	28 (2%) 52 56	10, 21, 40, 78	34 (3%)

All (28) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	482	LEU	5.0
1	B	43	MET	4.8
1	B	481	THR	3.7
1	A	481	THR	3.7
1	B	8	VAL	3.6
1	A	409[A]	ALA	3.5
1	A	8	VAL	3.5
1	A	43	MET	3.2
1	B	41	GLU	3.1
1	A	410	THR	3.1
1	B	40	ARG	2.9
1	B	48	ALA	2.9
1	A	480	LYS	2.9
1	B	50	VAL	2.7
1	A	408[A]	GLU	2.7
1	B	453	MET	2.6
1	B	42	LEU	2.6
1	A	207	ASP	2.4
1	B	208	LEU	2.3
1	B	286	GLN	2.3
1	A	304	THR	2.2
1	B	409	ALA	2.1
1	A	47	GLY	2.1
1	B	47	GLY	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	453	MET	2.1
1	B	410	THR	2.1
1	B	44	ILE	2.1
1	B	46	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.