



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 10:37 AM UTC

PDB ID : 7B94 / pdb_00007b94
Title : MEK1 in complex with compound 6
Authors : Kack, H.; Oster, L.
Deposited on : 2020-12-14
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	326	
1	B	326	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 5073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dual specificity mitogen-activated protein kinase kinase 1, Dual specificity mitogen-activated protein kinase kinase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	301	Total	C	N	O	S	0	0	0
			2380	1519	406	440	15			
1	B	295	Total	C	N	O	S	0	0	0
			2316	1475	399	427	15			

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	initiating methionine	UNP Q02750
A	21	ALA	-	expression tag	UNP Q02750
A	22	HIS	-	expression tag	UNP Q02750
A	23	HIS	-	expression tag	UNP Q02750
A	24	HIS	-	expression tag	UNP Q02750
A	25	HIS	-	expression tag	UNP Q02750
A	26	HIS	-	expression tag	UNP Q02750
A	27	HIS	-	expression tag	UNP Q02750
A	28	ALA	-	expression tag	UNP Q02750
A	29	ALA	-	expression tag	UNP Q02750
A	30	ALA	-	expression tag	UNP Q02750
A	31	GLU	-	expression tag	UNP Q02750
A	32	ASN	-	expression tag	UNP Q02750
A	33	LEU	-	expression tag	UNP Q02750
A	34	TYR	-	expression tag	UNP Q02750
A	35	PHE	-	expression tag	UNP Q02750
A	36	GLN	-	expression tag	UNP Q02750
A	302	GLY	-	linker	UNP Q02750
A	303	SER	-	linker	UNP Q02750
A	304	GLY	-	linker	UNP Q02750
A	305	SER	-	linker	UNP Q02750
A	306	GLY	-	linker	UNP Q02750
A	307	SER	-	linker	UNP Q02750
B	20	MET	-	initiating methionine	UNP Q02750

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	21	ALA	-	expression tag	UNP Q02750
B	22	HIS	-	expression tag	UNP Q02750
B	23	HIS	-	expression tag	UNP Q02750
B	24	HIS	-	expression tag	UNP Q02750
B	25	HIS	-	expression tag	UNP Q02750
B	26	HIS	-	expression tag	UNP Q02750
B	27	HIS	-	expression tag	UNP Q02750
B	28	ALA	-	expression tag	UNP Q02750
B	29	ALA	-	expression tag	UNP Q02750
B	30	ALA	-	expression tag	UNP Q02750
B	31	GLU	-	expression tag	UNP Q02750
B	32	ASN	-	expression tag	UNP Q02750
B	33	LEU	-	expression tag	UNP Q02750
B	34	TYR	-	expression tag	UNP Q02750
B	35	PHE	-	expression tag	UNP Q02750
B	36	GLN	-	expression tag	UNP Q02750
B	264	GLY	-	linker	UNP Q02750
B	303	SER	-	linker	UNP Q02750
B	304	GLY	-	linker	UNP Q02750
B	305	SER	-	linker	UNP Q02750
B	306	GLY	-	linker	UNP Q02750
B	307	SER	-	linker	UNP Q02750

- Molecule 2 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (CCD ID: ANP) (formula: $C_{10}H_{17}N_6O_{12}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	10	6	12	3		
2	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is 2-(4-iodophenyl)-8 {H}-imidazo[1,2-c]pyrimidin-5-one (CCD ID: T3W) (formula: C₁₂H₈IN₃O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	I	N	O	0	0
			17	12	1	3	1		
4	B	1	Total	C	I	N	O	0	0
			17	12	1	3	1		

- Molecule 5 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	135	Total	O	0	0
			135	135		
6	B	134	Total	O	0	0
			134	134		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dual specificity mitogen-activated protein kinase kinase 1, Dual specificity mitogen-activated protein kinase kinase 1



- Molecule 1: Dual specificity mitogen-activated protein kinase kinase 1, Dual specificity mitogen-activated protein kinase kinase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.28Å 71.75Å 157.80Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	78.90 – 2.00 78.90 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.7 (78.90-2.00) 96.7 (78.90-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.27 (at 2.00Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.216 , 0.265 0.213 , 0.263	Depositor DCC
R_{free} test set	2115 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	37.8	Xtriage
Anisotropy	0.400	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 65.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5073	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ANP, MG, T3W

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.88	1/2423 (0.0%)	1.25	4/3257 (0.1%)
1	B	0.87	1/2357 (0.0%)	1.29	10/3168 (0.3%)
All	All	0.88	2/4780 (0.0%)	1.27	14/6425 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	94	MET	SD-CE	-8.70	1.57	1.79
1	B	308	MET	SD-CE	-7.91	1.59	1.79

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	139	ILE	N-CA-CB	8.57	120.06	110.72
1	B	382	ASN	CA-CB-CG	7.67	120.27	112.60
1	A	84	LYS	N-CA-C	-5.92	99.55	108.96
1	B	147	ASP	CA-CB-CG	5.75	118.34	112.60
1	A	345	ASN	CA-CB-CG	5.64	118.24	112.60
1	A	43	ASP	CA-CB-CG	5.53	118.13	112.60
1	B	136	ASP	CA-CB-CG	5.44	118.04	112.60
1	B	91	GLY	CA-C-N	5.34	128.44	120.90
1	B	91	GLY	C-N-CA	5.34	128.44	120.90
1	B	117	VAL	CA-C-N	5.19	127.75	120.28
1	B	117	VAL	C-N-CA	5.19	127.75	120.28
1	A	154	VAL	N-CA-CB	5.12	120.04	110.77
1	B	138	GLU	CA-C-N	-5.01	115.90	122.37
1	B	138	GLU	C-N-CA	-5.01	115.90	122.37

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2380	0	2406	22	0
1	B	2316	0	2350	25	0
2	A	31	0	13	0	0
2	B	31	0	13	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	17	0	0	1	0
4	B	17	0	0	1	0
5	A	5	0	0	0	0
5	B	5	0	0	0	0
6	A	135	0	0	1	0
6	B	134	0	0	0	0
All	All	5073	0	4782	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:94:MET:CE	1:A:142:CYS:HB3	1.93	0.98
1:A:188:HIS:HD2	1:A:190:ASP:H	1.23	0.85
1:B:343:ILE:O	1:B:349:ARG:HD3	1.83	0.78
1:A:94:MET:HE1	1:A:142:CYS:HB3	1.70	0.73
1:A:343:ILE:O	1:A:349:ARG:HD3	1.90	0.70
1:B:188:HIS:HD2	1:B:190:ASP:H	1.40	0.70
1:A:227:ARG:HE	1:A:261:TYR:HB2	1.58	0.69
1:A:112:ILE:HD11	1:A:134:TYR:HB2	1.77	0.66
1:A:43:ASP:H	1:A:46:GLN:HE21	1.45	0.65
1:B:234:ARG:HD3	1:B:240:TYR:HD1	1.65	0.61
1:B:234:ARG:HD3	1:B:240:TYR:CD1	2.37	0.60
1:A:343:ILE:HG21	1:A:348:GLU:HB3	1.83	0.59
1:A:94:MET:HE2	1:A:142:CYS:HB3	1.82	0.59
1:B:115:LEU:O	1:B:118:LEU:HB2	2.05	0.57
1:B:118:LEU:HD13	1:B:211:VAL:HG21	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:57:LYS:NZ	1:B:119:HIS:HD2	2.06	0.54
1:A:261:TYR:HE2	1:A:313:LEU:HD11	1.73	0.53
1:B:260:ARG:HH11	1:B:260:ARG:HG2	1.74	0.53
1:B:343:ILE:O	1:B:349:ARG:CD	2.56	0.52
1:B:189:ARG:HA	1:B:240:TYR:CZ	2.45	0.51
1:A:255:GLU:HB2	1:A:262:PRO:HD3	1.93	0.51
1:A:341:CYS:O	1:A:349:ARG:HD2	2.11	0.51
1:B:118:LEU:HB3	1:B:129:PHE:CD2	2.46	0.50
1:B:310:ILE:H	1:B:310:ILE:HD12	1.76	0.50
1:A:181:ARG:HD2	1:A:242:VAL:HG11	1.95	0.49
1:A:188:HIS:HD2	1:A:190:ASP:N	2.01	0.48
1:B:57:LYS:O	1:B:60:VAL:HG22	2.12	0.48
1:A:189:ARG:HA	1:A:240:TYR:CZ	2.50	0.46
1:A:241:SER:HB3	6:A:546:HOH:O	2.16	0.46
1:A:188:HIS:CD2	1:A:190:ASP:H	2.15	0.46
1:B:308:MET:CE	1:B:316:TYR:CD2	2.99	0.46
1:B:308:MET:HE2	1:B:313:LEU:HA	1.97	0.45
1:B:69:GLU:HB2	1:B:88:LYS:HE2	1.99	0.45
1:A:247:TRP:CD1	1:A:247:TRP:C	2.95	0.44
1:B:379:ILE:HD11	1:B:381:LEU:HD12	1.99	0.43
1:A:200:SER:HB3	1:A:381:LEU:HD13	1.99	0.43
1:B:208:ASP:HA	4:B:403:T3W:C5	2.49	0.42
1:B:188:HIS:ND1	1:B:209:PHE:HB3	2.34	0.42
1:B:105:PRO:HA	1:B:108:ARG:HG2	2.01	0.42
1:B:312:GLU:O	1:B:316:TYR:HD2	2.03	0.42
1:A:169:VAL:O	1:A:173:VAL:HG23	2.20	0.41
1:B:94:MET:CE	1:B:142:CYS:HB3	2.50	0.41
1:B:215:LEU:O	1:B:219:MET:HG2	2.20	0.41
1:A:163:GLU:HG3	1:A:334:PHE:CD2	2.55	0.41
1:B:108:ARG:HD3	1:B:134:TYR:CD1	2.56	0.41
1:B:118:LEU:HB3	1:B:129:PHE:CG	2.56	0.40
1:A:208:ASP:HA	4:A:403:T3W:C5	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/326 (90%)	285 (97%)	9 (3%)	1 (0%)	36	35
1	B	289/326 (89%)	281 (97%)	8 (3%)	0	100	100
All	All	584/652 (90%)	566 (97%)	17 (3%)	1 (0%)	43	42

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	42	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	262/280 (94%)	242 (92%)	20 (8%)	12	9
1	B	255/280 (91%)	234 (92%)	21 (8%)	10	7
All	All	517/560 (92%)	476 (92%)	41 (8%)	11	8

All (41) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	A	41	GLU
1	A	62	GLU
1	A	82	VAL
1	A	108	ARG
1	A	136	ASP
1	A	139	ILE
1	A	154	VAL
1	A	157	LYS
1	A	177	LEU
1	A	186	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	197	LEU
1	A	215	LEU
1	A	234	ARG
1	A	308	MET
1	A	313	LEU
1	A	325	LEU
1	A	329	VAL
1	A	375	LEU
1	A	382	ASN
1	B	56	GLN
1	B	62	GLU
1	B	82	VAL
1	B	102	GLU
1	B	107	ILE
1	B	118	LEU
1	B	139	ILE
1	B	154	VAL
1	B	177	LEU
1	B	183	LYS
1	B	187	MET
1	B	197	LEU
1	B	211	VAL
1	B	215	LEU
1	B	242	VAL
1	B	260	ARG
1	B	325	LEU
1	B	344	LYS
1	B	349	ARG
1	B	375	LEU
1	B	377	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	36	GLN
1	A	46	GLN
1	A	188	HIS
1	A	214	GLN
1	A	345	ASN
1	A	354	GLN
1	B	119	HIS
1	B	122	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	188	HIS
1	B	354	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 2 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	T3W	A	403	-	16,19,19	2.85	2 (12%)	17,27,27	2.07	4 (23%)
2	ANP	B	401	3	33,33,33	0.25	0	45,52,52	0.72	2 (4%)
5	SO4	A	404	-	4,4,4	0.25	0	6,6,6	0.15	0
2	ANP	A	401	3	33,33,33	0.47	0	45,52,52	0.75	3 (6%)
4	T3W	B	403	-	16,19,19	2.89	3 (18%)	17,27,27	1.88	4 (23%)
5	SO4	B	404	-	4,4,4	0.28	0	6,6,6	0.08	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	T3W	B	403	-	-	0/4/14/14	0/3/3/3
2	ANP	A	401	3	-	2/18/38/38	0/3/3/3
2	ANP	B	401	3	-	3/18/38/38	0/3/3/3
4	T3W	A	403	-	-	0/4/14/14	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	403	T3W	C7-C11	-10.89	1.33	1.51
4	A	403	T3W	C7-C11	-10.69	1.34	1.51
4	A	403	T3W	C11-N14	2.65	1.37	1.32
4	B	403	T3W	C11-N14	2.38	1.36	1.32
4	B	403	T3W	C8-C10	-2.04	1.33	1.37

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	T3W	C8-N15-C11	-5.84	105.86	107.66
4	B	403	T3W	C8-N15-C11	-5.37	106.00	107.66
4	A	403	T3W	C5-C10-C8	-4.13	123.34	129.65
4	B	403	T3W	C5-C10-C8	-3.51	124.28	129.65
4	A	403	T3W	C8-C10-N14	2.72	113.35	110.75
2	A	401	ANP	O2G-PG-O1G	-2.58	106.97	113.45
2	A	401	ANP	O1G-PG-N3B	2.55	115.52	111.77
2	A	401	ANP	O3G-PG-O1G	-2.52	107.14	113.45
4	B	403	T3W	C8-C10-N14	2.40	113.04	110.75
4	A	403	T3W	C7-C11-N14	2.15	127.57	123.47
2	B	401	ANP	O1B-PB-N3B	2.10	114.87	111.77
2	B	401	ANP	O3G-PG-O1G	-2.06	108.30	113.45
4	B	403	T3W	C7-C11-N14	2.00	127.28	123.47

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	401	ANP	PB-N3B-PG-O1G
2	A	401	ANP	PG-N3B-PB-O3A
2	B	401	ANP	PB-N3B-PG-O1G
2	B	401	ANP	PA-O3A-PB-O2B
2	B	401	ANP	PA-O3A-PB-O1B


There are no ring outliers.


2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	403	T3W	1	0
4	B	403	T3W	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	301/326 (92%)	0.45	13 (4%) 40 39	31, 45, 74, 114	0
1	B	295/326 (90%)	0.63	20 (6%) 23 22	29, 50, 91, 112	0
All	All	596/652 (91%)	0.54	33 (5%) 30 29	29, 48, 83, 114	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	220	ALA	5.6
1	A	226	THR	3.7
1	A	102	GLU	3.6
1	B	240	TYR	3.5
1	A	33	LEU	3.3
1	B	264	GLY	3.2
1	A	310	ILE	3.1
1	B	41	GLU	3.0
1	B	239	HIS	3.0
1	B	53	PHE	2.9
1	A	101	LEU	2.8
1	A	263	ILE	2.8
1	B	68	PHE	2.8
1	A	103	ILE	2.8
1	B	63	LEU	2.8
1	B	49	ARG	2.7
1	A	219	MET	2.6
1	A	107	ILE	2.5
1	B	42	LEU	2.5
1	B	117	VAL	2.4
1	B	316	TYR	2.4
1	B	139	ILE	2.3
1	B	71	ILE	2.3
1	A	329	VAL	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	237	GLY	2.3
1	A	316	TYR	2.3
1	B	310	ILE	2.1
1	A	186	ILE	2.1
1	B	216	ILE	2.1
1	B	219	MET	2.1
1	B	377	SER	2.1
1	B	378	THR	2.0
1	B	56	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	B	404	5/5	0.56	0.13	143,143,144,144	0
5	SO4	A	404	5/5	0.82	0.09	83,85,86,88	0
4	T3W	B	403	17/17	0.85	0.18	91,96,111,117	0
4	T3W	A	403	17/17	0.90	0.16	61,66,84,93	0
2	ANP	A	401	31/31	0.94	0.07	32,43,58,61	0
2	ANP	B	401	31/31	0.95	0.07	34,46,59,62	0
3	MG	A	402	1/1	0.99	0.03	44,44,44,44	0
3	MG	B	402	1/1	0.99	0.03	38,38,38,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around T3W B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around T3W A 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ANP A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP B 401:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.