



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 06:02 AM UTC

PDB ID : 8B91 / pdb_00008b91
Title : Crystal structure of mutant PPARG (C313A) and NCOR2 with an inverse agonist (compound SI-1)
Authors : Friberg, A.; Orsi, D.L.; Pook, E.; Siegel, S.; Lemke, C.T.; Stellfeld, T.; Puetter, V.; Goldstein, J.
Deposited on : 2022-10-05
Resolution : 2.23 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	
1	B	279	
2	C	23	
2	D	23	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4438 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisome proliferator-activated receptor gamma.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	261	Total	C	N	O	S	0	2	0
			2109	1361	347	392	9			
1	B	241	Total	C	N	O	S	0	1	0
			1928	1244	317	359	8			


There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	199	GLY	-	expression tag	UNP P37231
A	200	SER	-	expression tag	UNP P37231
A	201	HIS	-	expression tag	UNP P37231
A	202	MET	-	expression tag	UNP P37231
A	285	ALA	CYS	engineered mutation	UNP P37231
B	199	GLY	-	expression tag	UNP P37231
B	200	SER	-	expression tag	UNP P37231
B	201	HIS	-	expression tag	UNP P37231
B	202	MET	-	expression tag	UNP P37231
B	285	ALA	CYS	engineered mutation	UNP P37231

- Molecule 2 is a protein called Nuclear receptor corepressor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	11	Total	C	N	O	S	0	0	0
			83	54	15	13	1			
2	D	12	Total	C	N	O	S	0	0	0
			87	56	16	14	1			

- Molecule 3 is {N}3-[4-[bis(fluoranyl)methoxy]-3-fluoranyl-phenyl]-4-chloranyl-6-fluoranyl-1-{N}1-[(2-methoxyphenyl)methyl]benzene-1,3-dicarboxamide (CCD ID: Q4O) (formula: C₂₃H₁₇ClF₄N₂O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	F	N	O	0	0
			34	23	1	4	2	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	122	Total	O	0	0
			122	122		
4	B	75	Total	O	0	0
			75	75		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peroxisome proliferator-activated receptor gamma




- Molecule 1: Peroxisome proliferator-activated receptor gamma



- Molecule 2: Nuclear receptor corepressor 2



- Molecule 2: Nuclear receptor corepressor 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.35Å 86.13Å 120.86Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.47 – 2.23 49.47 – 2.23	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.47-2.23) 99.4 (49.47-2.23)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.57 (at 2.22Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.202 , 0.228 0.209 , 0.231	Depositor DCC
R_{free} test set	1611 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.051	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4438	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: Q4O

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.99	0/2144	1.47	5/2886 (0.2%)
1	B	1.00	0/1958	1.46	2/2635 (0.1%)
2	C	1.05	0/82	1.82	0/107
2	D	1.18	0/86	1.76	0/112
All	All	1.00	0/4270	1.48	7/5740 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	363	PHE	CA-CB-CG	6.50	120.30	113.80
1	B	380	ASP	CA-CB-CG	5.53	118.13	112.60
1	A	441	ASP	CA-C-N	5.46	129.01	120.82
1	A	441	ASP	C-N-CA	5.46	129.01	120.82
1	B	363	PHE	CA-CB-CG	5.44	119.24	113.80
1	A	465	LEU	CA-C-N	5.16	127.99	120.51
1	A	465	LEU	C-N-CA	5.16	127.99	120.51

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2109	0	2166	29	0
1	B	1928	0	1985	20	0
2	C	83	0	97	0	0
2	D	87	0	100	2	0
3	A	34	0	0	1	0
4	A	122	0	0	2	0
4	B	75	0	0	0	0
All	All	4438	0	4348	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:327[B]:TYR:CE2	1:A:445:ILE:HG21	1.88	1.09
1:B:370:PHE:HB2	1:B:445:ILE:HD11	1.46	0.98
1:A:327[B]:TYR:CE2	1:A:445:ILE:CG2	2.46	0.98
1:A:327[B]:TYR:CZ	1:A:445:ILE:CG2	2.66	0.77
1:B:370:PHE:CB	1:B:445:ILE:HD11	2.17	0.75
1:A:327[A]:TYR:OH	1:A:449:HIS:CD2	2.41	0.73
1:A:217:HIS:HE1	1:A:302:SER:O	1.74	0.70
1:A:327[B]:TYR:CD2	1:A:445:ILE:HG21	2.27	0.69
1:B:357:ARG:O	1:B:357:ARG:HG3	2.03	0.58
1:A:327[B]:TYR:CZ	1:A:445:ILE:HG22	2.38	0.58
1:A:205:ASN:HB2	1:A:206:PRO:HD2	1.88	0.56
1:A:327[B]:TYR:CE2	1:A:367:LYS:HG3	2.41	0.55
1:A:379:LEU:HD11	1:A:435:LEU:HD21	1.90	0.54
1:B:324:GLU:HA	1:B:442:LEU:HD23	1.91	0.53
1:B:324:GLU:O	1:B:442:LEU:HD21	2.08	0.53
1:A:383:ASP:OD2	1:A:425:HIS:HE1	1.93	0.51
1:A:205:ASN:HB2	1:A:206:PRO:CD	2.40	0.51
1:B:459:THR:C	1:B:460:GLU:HG2	2.36	0.51
1:B:297:THR:HG21	2:D:2357:ALA:HB3	1.93	0.50
1:B:327:TYR:OH	1:B:449:HIS:CD2	2.65	0.49
1:B:226:PHE:CG	1:B:329:MET:HE1	2.46	0.49
1:B:383:ASP:OD2	1:B:425:HIS:HE1	1.94	0.49
1:A:204:LEU:HD23	4:A:613:HOH:O	2.13	0.48
1:A:449:HIS:NE2	1:A:453:LEU:HD11	2.29	0.48
1:A:433:ALA:O	1:A:437:GLN:HG3	2.13	0.48
1:B:289:SER:O	1:B:293:VAL:HG23	2.14	0.48
1:B:327:TYR:OH	1:B:449:HIS:CG	2.67	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:370:PHE:CB	1:B:445:ILE:CD1	2.89	0.48
1:B:393:LEU:HD22	1:B:409:ILE:HB	1.96	0.47
1:B:259:GLU:OE2	1:B:280:ARG:NH1	2.47	0.47
1:A:255:LEU:HD22	1:A:277:VAL:CG1	2.45	0.47
1:B:404:LYS:HB3	1:B:405:PRO:HD3	1.96	0.47
1:A:436:LEU:HD23	1:A:439:MET:HE3	1.97	0.47
1:A:284:GLY:HA3	3:A:501:Q4O:C26	2.45	0.47
1:A:327[A]:TYR:OH	1:A:449:HIS:NE2	2.49	0.45
1:A:393:LEU:HD13	1:A:413:LEU:HD11	1.98	0.45
1:A:358:LYS:HD3	4:A:687:HOH:O	2.17	0.44
1:B:455:VAL:O	1:B:459:THR:HG22	2.18	0.44
1:A:442:LEU:HD12	1:A:442:LEU:HA	1.88	0.43
1:A:440:THR:O	1:A:441:ASP:C	2.62	0.42
1:A:449:HIS:CD2	1:A:453:LEU:HD11	2.54	0.42
1:B:439:MET:O	1:B:443:ARG:HG3	2.20	0.42
1:A:255:LEU:HD22	1:A:277:VAL:HG13	2.02	0.42
1:A:404:LYS:N	1:A:405:PRO:CD	2.83	0.41
1:B:301:LYS:HE2	2:D:2358:LEU:HA	2.02	0.41
1:A:255:LEU:HD13	1:A:281:ILE:HD11	2.02	0.41
1:A:255:LEU:CD2	1:A:277:VAL:HG13	2.50	0.41
1:B:379:LEU:HD11	1:B:435:LEU:HD13	2.02	0.41
1:A:370:PHE:CE1	1:A:442:LEU:HD13	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	257/279 (92%)	251 (98%)	6 (2%)	0	100	100
1	B	236/279 (85%)	233 (99%)	2 (1%)	1 (0%)	30	30
2	C	9/23 (39%)	9 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	10/23 (44%)	10 (100%)	0	0	100	100
All	All	512/604 (85%)	503 (98%)	8 (2%)	1 (0%)	43	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	239	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	235/250 (94%)	222 (94%)	13 (6%)	19	18
1	B	214/250 (86%)	206 (96%)	8 (4%)	30	35
2	C	8/18 (44%)	7 (88%)	1 (12%)	4	2
2	D	8/18 (44%)	8 (100%)	0	100	100
All	All	465/536 (87%)	443 (95%)	22 (5%)	23	24

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	204	LEU
1	A	255	LEU
1	A	263	LYS
1	A	265	LYS
1	A	310	ASP
1	A	342	SER
1	A	358	LYS
1	A	434	LYS
1	A	439	MET
1	A	442	LEU
1	A	445	ILE
1	A	450	VAL
1	A	465	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	208	SER
1	B	232	LYS
1	B	259	GLU
1	B	260	ASP
1	B	363	PHE
1	B	393	LEU
1	B	442	LEU
1	B	459	THR
2	C	2356	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	217	HIS
1	A	425	HIS
1	A	437	GLN
1	B	203	GLN
1	B	253	ASN
1	B	425	HIS
1	B	444	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	Q4O	A	501	-	36,36,36	0.78	0	50,50,50	1.54	8 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	Q4O	A	501	-	-	3/23/23/23	0/3/3/3

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	Q4O	O33-C32-C27	4.63	122.85	115.96
3	A	501	Q4O	O33-C32-C31	-4.11	117.37	124.30
3	A	501	Q4O	C1-C23-N24	2.86	122.88	117.32
3	A	501	Q4O	C3-C9-N11	2.67	121.58	116.03
3	A	501	Q4O	F18-C16-C17	2.58	123.79	118.64
3	A	501	Q4O	C4-C3-C9	-2.52	118.49	122.56
3	A	501	Q4O	O10-C9-C3	-2.40	116.64	121.03
3	A	501	Q4O	O25-C23-N24	-2.07	118.56	122.59

There are no chirality outliers.

All (3) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	501	Q4O	C31-C32-O33-C34
3	A	501	Q4O	C27-C32-O33-C34
3	A	501	Q4O	N24-C26-C27-C32

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	Q4O	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	261/279 (93%)	0.10	7 (2%) 56 57	14, 36, 72, 90	2 (0%)
1	B	241/279 (86%)	0.11	7 (2%) 53 54	13, 38, 69, 95	1 (0%)
2	C	11/23 (47%)	1.12	1 (9%) 15 13	48, 59, 80, 85	0
2	D	12/23 (52%)	0.79	1 (8%) 17 15	39, 51, 70, 79	0
All	All	525/604 (86%)	0.14	16 (3%) 52 52	13, 38, 72, 95	3 (0%)

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	469	LEU	3.3
1	A	465	LEU	3.1
1	A	444	GLN	3.0
2	C	2359	MET	2.9
1	B	239	GLY	2.8
1	B	445	ILE	2.7
1	B	287	PHE	2.5
2	D	2359	MET	2.4
1	B	442	LEU	2.4
1	A	201	HIS	2.4
1	B	204	LEU	2.3
1	A	445	ILE	2.3
1	B	455	VAL	2.2
1	A	461	THR	2.1
1	A	442	LEU	2.1
1	B	363	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	Q4O	A	501	34/34	0.91	0.10	46,54,63,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.