



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 10:01 AM UTC

PDB ID : 3BAD / pdb_00003bad
Title : Crystal structure of D70A/H93G mutant of Human acidic fibroblast growth factor
Authors : Blaber, M.; Lee, J.
Deposited on : 2007-11-07
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	146	 73% 14% .. 10%
1	B	146	 74% 14% • 12%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2144 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Heparin-binding growth factor 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	131	Total	C	N	O	S	0	0	0
			1040	659	181	196	4			
1	B	129	Total	C	N	O	S	0	0	0
			1019	644	178	193	4			

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	HIS	-	expression tag	UNP P05230
A	-4	HIS	-	expression tag	UNP P05230
A	-3	HIS	-	expression tag	UNP P05230
A	-2	HIS	-	expression tag	UNP P05230
A	-1	HIS	-	expression tag	UNP P05230
A	0	HIS	-	expression tag	UNP P05230
A	70	ALA	ASP	engineered mutation	UNP P05230
A	93	GLY	HIS	engineered mutation	UNP P05230
B	-5	HIS	-	expression tag	UNP P05230
B	-4	HIS	-	expression tag	UNP P05230
B	-3	HIS	-	expression tag	UNP P05230
B	-2	HIS	-	expression tag	UNP P05230
B	-1	HIS	-	expression tag	UNP P05230
B	0	HIS	-	expression tag	UNP P05230
B	70	ALA	ASP	engineered mutation	UNP P05230
B	93	GLY	HIS	engineered mutation	UNP P05230

- Molecule 2 is SULFATE ION (CCD ID: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	34	Total	O	0	0
			34	34		
3	B	41	Total	O	0	0
			41	41		

i

- Molecule 1: Heparin-binding growth factor 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	35.21Å 111.08Å 36.84Å 90.00° 117.71° 90.00°	Depositor
Resolution (Å)	31.29 – 2.00 31.29 – 2.00	Depositor EDS
% Data completeness (in resolution range)	85.3 (31.29-2.00) 86.6 (31.29-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	12.44 (at 2.00Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.213 , 0.253 0.220 , 0.261	Depositor DCC
R_{free} test set	769 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	25.0	Xtriage
Anisotropy	1.094	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 38.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.380 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	2144	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 13.44% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	0/1063	0.89	2/1433 (0.1%)
1	B	0.56	0/1041	0.90	1/1404 (0.1%)
All	All	0.55	0/2104	0.90	3/2837 (0.1%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	109	VAL	N-CA-C	-7.26	99.34	109.21
1	A	55	TYR	N-CA-C	-5.58	101.08	109.95
1	A	109	VAL	N-CA-C	-5.27	101.93	108.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1040	0	1031	18	0
1	B	1019	0	1009	13	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
3	A	34	0	0	2	0
3	B	41	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	2144	0	2040	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:GLU:H	1:A:91:GLU:CD	1.91	0.77
1:B:53:GLU:OE2	1:B:100:LYS:HE3	1.98	0.64
1:A:22:PHE:CE2	1:A:37:ARG:HG2	2.35	0.61
1:A:107:TRP:CE2	1:A:121:PRO:HD3	2.38	0.58
1:B:112:LYS:HD2	1:B:116:SER:OG	2.06	0.56
1:A:121:PRO:HG2	3:A:149:HOH:O	2.05	0.55
1:A:8:TYR:HA	1:A:46:LEU:O	2.06	0.55
1:B:112:LYS:HE2	3:B:142:HOH:O	2.05	0.55
1:A:9:LYS:HE3	1:A:9:LYS:HA	1.89	0.53
1:A:8:TYR:O	1:A:9:LYS:HB2	2.08	0.52
1:A:64:TYR:CE1	1:A:79:PRO:HD3	2.45	0.52
1:B:127:GLN:HB3	3:B:178:HOH:O	2.09	0.52
1:B:10:LYS:HE2	1:B:11:PRO:HD2	1.94	0.49
1:B:26:LEU:HB3	1:B:27:PRO:HD2	1.95	0.49
1:B:91:GLU:N	1:B:91:GLU:CD	2.71	0.48
1:B:91:GLU:CD	1:B:91:GLU:H	2.21	0.48
1:A:27:PRO:HG2	3:A:172:HOH:O	2.14	0.47
1:B:72:LEU:HD23	1:B:119:ARG:HG2	1.95	0.47
1:A:50:SER:OG	1:A:51:VAL:N	2.47	0.47
1:A:136:PRO:HB3	1:B:92:ASN:O	2.16	0.46
1:A:22:PHE:HE2	1:A:37:ARG:HG2	1.81	0.44
1:A:107:TRP:CZ2	1:A:121:PRO:HD3	2.52	0.44
1:B:68:ASP:OD1	1:B:68:ASP:C	2.60	0.44
1:A:91:GLU:CD	1:A:91:GLU:N	2.67	0.43
1:A:16:CYS:HB2	1:A:132:PHE:CE2	2.52	0.43
1:A:136:PRO:HG3	1:B:93:GLY:HA3	2.01	0.43
1:B:110:GLY:HA2	1:B:129:ALA:O	2.20	0.41
1:A:73:LEU:HD12	1:A:117:CYS:HB2	2.02	0.41
1:A:104:GLU:H	1:A:104:GLU:HG2	1.72	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	129/146 (88%)	125 (97%)	2 (2%)	2 (2%)	7	3
1	B	127/146 (87%)	121 (95%)	6 (5%)	0	100	100
All	All	256/292 (88%)	246 (96%)	8 (3%)	2 (1%)	16	11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	120	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	113/127 (89%)	108 (96%)	5 (4%)	25	24
1	B	111/127 (87%)	107 (96%)	4 (4%)	31	31
All	All	224/254 (88%)	215 (96%)	9 (4%)	28	27

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	TYR
1	A	9	LYS
1	A	86	LEU
1	A	91	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	106	ASN
1	B	45	GLN
1	B	86	LEU
1	B	91	GLU
1	B	104	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	HIS
1	A	63	GLN
1	A	95	ASN
1	A	106	ASN
1	B	21	HIS
1	B	41	HIS
1	B	45	GLN
1	B	95	ASN
1	B	106	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	141	-	4,4,4	0.39	0	6,6,6	0.16	0
2	SO4	B	141	-	4,4,4	0.45	0	6,6,6	0.17	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	131/146 (89%)	-1.24	0 100 100	27, 38, 56, 67	0
1	B	129/146 (88%)	-1.25	0 100 100	27, 37, 55, 62	0
All	All	260/292 (89%)	-1.25	0 100 100	27, 38, 56, 67	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	SO4	B	141	5/5	0.97	0.06	79,79,79,80	0
2	SO4	A	141	5/5	0.98	0.05	72,73,73,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.