



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 02:04 PM UTC

PDB ID : 6BA4 / pdb_00006ba4
Title : Crystal structure of MYST acetyltransferase domain in complex with Acetyl-CoA cofactor
Authors : Hermans, S.J.; Chung, M.C.; Peat, T.S.; Baell, J.B.; Thomas, T.; Parker, M.W.
Deposited on : 2017-10-12
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	295	8% 76% 14% 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 2471 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone acetyltransferase KAT8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	1	0
			2211	1441	358	400	12			

There are 26 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	485	MET	-	initiating methionine	UNP Q9H7Z6
A	486	GLY	-	expression tag	UNP Q9H7Z6
A	487	SER	-	expression tag	UNP Q9H7Z6
A	488	SER	-	expression tag	UNP Q9H7Z6
A	489	HIS	-	expression tag	UNP Q9H7Z6
A	490	HIS	-	expression tag	UNP Q9H7Z6
A	491	HIS	-	expression tag	UNP Q9H7Z6
A	492	HIS	-	expression tag	UNP Q9H7Z6
A	493	HIS	-	expression tag	UNP Q9H7Z6
A	494	HIS	-	expression tag	UNP Q9H7Z6
A	495	SER	-	expression tag	UNP Q9H7Z6
A	496	SER	-	expression tag	UNP Q9H7Z6
A	497	GLY	-	expression tag	UNP Q9H7Z6
A	498	LEU	-	expression tag	UNP Q9H7Z6
A	499	VAL	-	expression tag	UNP Q9H7Z6
A	500	PRO	-	expression tag	UNP Q9H7Z6
A	501	ARG	-	expression tag	UNP Q9H7Z6
A	502	GLY	-	expression tag	UNP Q9H7Z6
A	503	SER	-	expression tag	UNP Q9H7Z6
A	579	HIS	TYR	conflict	UNP Q9H7Z6
A	645	SER	ALA	conflict	UNP Q9H7Z6
A	648	MET	LEU	conflict	UNP Q9H7Z6
A	649	ILE	THR	conflict	UNP Q9H7Z6
A	660	ARG	LYS	conflict	UNP Q9H7Z6
A	697	SER	TRP	conflict	UNP Q9H7Z6
A	702	ASN	ILE	conflict	UNP Q9H7Z6


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Na	0	0
			3	3		

- Molecule 4 is ACETYL COENZYME *A (CCD ID: ACO) (formula: C₂₃H₃₈N₇O₁₇P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	S	0	1
			51	23	7	17	3	1		

- Molecule 5 is S-{(3S,5R,9R)-1-[(2R,3R,4R,5R)-5-(6-amino-9H-purin-9-yl)-3-hydroxy-4-(phosphonooxy)tetrahydrofuran-2-yl]-3,5,9-trihydroxy-8,8-dimethyl-3,5-dioxido-10,14-dioxo-2,4,6-trioxa-11,15-diaza-3lambda 5 ,5lambda 5 -diphosphaheptadecan-17-yl} ethanethioate (CCD ID: 7L1) (formula: C₂₃H₃₈N₇O₁₇P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
5	A	1	Total	C	N	O	P	S	0	1
			51	23	7	17	3	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	154	Total	O	0	0
			154	154		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Histone acetyltransferase KAT8



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	46.36Å 57.92Å 120.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.68 – 1.95 41.68 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (41.68-1.95) 99.8 (41.68-1.95)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 1.95Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.184 , 0.213 0.188 , 0.214	Depositor DCC
R_{free} test set	1193 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	29.8	Xtriage
Anisotropy	0.687	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 47.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2471	wwPDB-VP
Average B, all atoms (Å ²)	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.22% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, 7L1, ACO, ALY, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/2267	0.83	9/3074 (0.3%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	587	LYS	N-CA-C	6.35	117.86	111.07
1	A	776	ALA	CA-C-N	6.33	126.89	120.38
1	A	776	ALA	C-N-CA	6.33	126.89	120.38
1	A	764	PRO	CA-C-N	-5.98	114.58	120.98
1	A	764	PRO	C-N-CA	-5.98	114.58	120.98
1	A	777	PRO	N-CA-C	5.18	117.02	110.70
1	A	651	PRO	CA-C-N	-5.17	114.75	119.82
1	A	651	PRO	C-N-CA	-5.17	114.75	119.82
1	A	586	HIS	N-CA-CB	-5.08	104.55	111.65

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2211	0	2166	30	0
2	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	3	0	0	0	0
4	A	51	0	34	1	0
5	A	51	0	0	4	0
6	A	154	0	0	5	0
All	All	2471	0	2200	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:566:PRO:HB2	1:A:586:HIS:CE1	2.12	0.83
1:A:620:GLU:OE2	1:A:658:TYR:OH	1.98	0.81
5:A:806[B]:7L1:O50	6:A:901:HOH:O	2.14	0.64
1:A:530:ASP:N	1:A:530:ASP:OD1	2.29	0.63
1:A:763:LYS:C	1:A:765:PRO:HD3	2.25	0.61
5:A:806[B]:7L1:O11	6:A:902:HOH:O	2.16	0.60
4:A:805[A]:ACO:O4A	6:A:902:HOH:O	2.17	0.60
1:A:573:LYS:NZ	6:A:910:HOH:O	2.35	0.59
1:A:585:ASP:C	1:A:586:HIS:HD2	2.13	0.56
1:A:591:GLN:HG2	1:A:608:PHE:HA	1.89	0.54
1:A:528:PRO:O	1:A:532:GLY:N	2.22	0.53
1:A:606:LEU:HD21	1:A:609:ASP:O	2.08	0.53
1:A:530:ASP:O	1:A:534:GLN:HG2	2.09	0.52
1:A:566:PRO:HB2	1:A:586:HIS:ND1	2.23	0.52
1:A:585:ASP:C	1:A:586:HIS:CD2	2.88	0.51
1:A:668:GLU:HG3	1:A:768:VAL:HG11	1.93	0.50
1:A:585:ASP:O	1:A:586:HIS:HD2	1.94	0.50
1:A:509:VAL:HG23	1:A:532:GLY:HA2	1.94	0.49
1:A:655:ARG:HH21	5:A:806[B]:7L1:P48	2.36	0.48
1:A:763:LYS:N	1:A:764:PRO:HD3	2.29	0.48
1:A:655:ARG:NH2	5:A:806[B]:7L1:O51	2.47	0.47
1:A:738:TYR:CZ	1:A:741:GLY:HA2	2.50	0.46
1:A:738:TYR:HB2	6:A:1024:HOH:O	2.15	0.46
1:A:775:TRP:CZ2	1:A:778:PRO:HD2	2.52	0.44
1:A:567:GLY:H	1:A:586:HIS:HE1	1.64	0.43
1:A:642:ASN:HA	1:A:678:SER:O	2.19	0.42
1:A:762:LYS:C	1:A:764:PRO:HD3	2.44	0.42
1:A:685:ASP:O	1:A:689:LEU:HG	2.20	0.41
1:A:562[B]:TRP:CZ3	1:A:566:PRO:HD3	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:611:GLU:CD	1:A:611:GLU:H	2.29	0.40
1:A:747:VAL:HG23	1:A:751:LEU:HD23	2.03	0.40
1:A:777:PRO:HA	1:A:778:PRO:HD2	1.65	0.40
1:A:566:PRO:HB2	1:A:586:HIS:HE1	1.77	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	264/295 (90%)	258 (98%)	5 (2%)	1 (0%)	30 22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	764	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	245/271 (90%)	241 (98%)	4 (2%)	55 46

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	507	LYS
1	A	704	ARG
1	A	742	GLN
1	A	750	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	586	HIS
1	A	592	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	ALY	A	604	1	10,11,12	0.85	0	7,12,14	1.15	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	ALY	A	604	1	-	1/9/10/12	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	604	ALY	CD-CG-CB	-2.39	104.62	113.62

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	604	ALY	CA-CB-CG-CD

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACO	A	805[A]	-	51,53,53	2.64	13 (25%)	73,79,79	2.23	12 (16%)
5	7L1	A	806[B]	-	51,53,53	2.75	12 (23%)	73,79,79	1.67	14 (19%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACO	A	805[A]	-	-	2/51/67/67	0/3/3/3
5	7L1	A	806[B]	-	-	6/51/67/67	0/3/3/3

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	806[B]	7L1	C46-C35	-10.75	1.26	1.53
4	A	805[A]	ACO	C2B-C3B	-9.76	1.31	1.53
5	A	806[B]	7L1	O34-C35	9.05	1.62	1.42
5	A	806[B]	7L1	O34-C03	-7.54	1.28	1.45
5	A	806[B]	7L1	C20-N22	6.40	1.48	1.33
4	A	805[A]	ACO	O4B-C4B	-5.91	1.31	1.45
4	A	805[A]	ACO	C9P-N8P	5.71	1.47	1.33
4	A	805[A]	ACO	O4B-C1B	5.69	1.55	1.42
4	A	805[A]	ACO	C1B-N9A	-5.27	1.31	1.46
4	A	805[A]	ACO	C6A-N6A	4.75	1.46	1.34
4	A	805[A]	ACO	C3B-C4B	4.17	1.63	1.52
4	A	805[A]	ACO	C5P-N4P	4.08	1.43	1.33
5	A	806[B]	7L1	C25-N27	3.96	1.42	1.33
4	A	805[A]	ACO	P1A-O3A	3.22	1.63	1.59
5	A	806[B]	7L1	C44-N45	3.15	1.42	1.34
4	A	805[A]	ACO	P2A-O3A	3.06	1.62	1.59
4	A	805[A]	ACO	C2A-N1A	3.03	1.39	1.33
5	A	806[B]	7L1	C37-N36	-2.68	1.33	1.37
5	A	806[B]	7L1	C04-C03	2.66	1.59	1.51
4	A	805[A]	ACO	C7P-C6P	2.54	1.59	1.51
5	A	806[B]	7L1	O02-C01	-2.41	1.37	1.43
5	A	806[B]	7L1	C24-C25	2.35	1.56	1.51
4	A	805[A]	ACO	C8A-N7A	2.30	1.36	1.31
5	A	806[B]	7L1	O19-C18	-2.15	1.38	1.42
5	A	806[B]	7L1	O47-C46	2.05	1.51	1.44

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	805[A]	ACO	N6A-C6A-N1A	-10.13	95.81	118.38
4	A	805[A]	ACO	C5A-C6A-N6A	8.48	144.29	123.29
4	A	805[A]	ACO	N3A-C2A-N1A	-5.38	120.43	128.58
5	A	806[B]	7L1	C39-C40-N41	-4.90	119.97	126.72
4	A	805[A]	ACO	N9A-C8A-N7A	-4.69	107.28	113.94
4	A	805[A]	ACO	C5A-C4A-N3A	-4.58	120.41	126.72
5	A	806[B]	7L1	C23-C24-C25	4.38	119.68	112.39
5	A	806[B]	7L1	N41-C40-N36	3.83	133.68	127.17
5	A	806[B]	7L1	N43-C42-N41	-3.77	122.87	128.58
4	A	805[A]	ACO	C4A-N9A-C8A	3.49	109.40	105.74
4	A	805[A]	ACO	N3A-C4A-N9A	3.32	132.81	127.17
5	A	806[B]	7L1	C42-N41-C40	3.31	119.92	111.83
5	A	806[B]	7L1	C40-C39-N38	-3.22	106.90	110.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	805[A]	ACO	C2A-N3A-C4A	3.03	119.23	111.83
4	A	805[A]	ACO	C5A-N7A-C8A	2.97	108.12	103.45
5	A	806[B]	7L1	C39-N38-C37	2.97	108.11	103.45
5	A	806[B]	7L1	P48-O47-C46	-2.89	115.70	123.43
5	A	806[B]	7L1	C46-C35-N36	-2.89	108.99	113.75
5	A	806[B]	7L1	N36-C37-N38	-2.77	110.01	113.94
5	A	806[B]	7L1	C40-N36-C37	2.76	108.63	105.74
5	A	806[B]	7L1	C16-C15-C18	2.50	113.03	108.77
5	A	806[B]	7L1	O21-C20-N22	-2.46	117.78	122.98
5	A	806[B]	7L1	O34-C35-C46	-2.32	102.59	106.59
4	A	805[A]	ACO	C2P-C3P-N4P	-2.30	107.61	112.41
4	A	805[A]	ACO	C7P-C6P-C5P	-2.12	108.86	112.39
4	A	805[A]	ACO	C6A-C5A-C4A	2.12	120.07	117.18

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	805[A]	ACO	S1P-C2P-C3P-N4P
5	A	806[B]	7L1	N22-C23-C24-C25
5	A	806[B]	7L1	C01-C46-O47-P48
5	A	806[B]	7L1	C35-C46-O47-P48
5	A	806[B]	7L1	C14-O13-P10-O09
5	A	806[B]	7L1	C14-O13-P10-O12
5	A	806[B]	7L1	N27-C28-C29-S30
4	A	805[A]	ACO	O4B-C4B-C5B-O5B


There are no ring outliers.


2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	805[A]	ACO	1	0
5	A	806[B]	7L1	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/295 (90%)	0.44	23 (8%) 16 18	18, 36, 65, 83	1 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	764	PRO	4.4
1	A	530	ASP	4.2
1	A	535	PRO	3.7
1	A	532	GLY	3.6
1	A	761	TYR	3.4
1	A	759	ALA	3.4
1	A	529	GLU	3.3
1	A	778	PRO	3.3
1	A	741	GLY	3.1
1	A	586	HIS	3.1
1	A	710	LEU	3.1
1	A	751	LEU	3.0
1	A	562[A]	TRP	2.9
1	A	765	PRO	2.9
1	A	763	LYS	2.8
1	A	738	TYR	2.6
1	A	760	GLN	2.6
1	A	528	PRO	2.5
1	A	606	LEU	2.3
1	A	776	ALA	2.3
1	A	750	LYS	2.2
1	A	742	GLN	2.1
1	A	749	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	ALY	A	604	12/13	0.94	0.09	25,34,42,42	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACO	A	805[A]	51/51	0.91	0.12	27,33,53,54	51
5	7L1	A	806[B]	51/51	0.91	0.12	27,33,53,57	51
3	NA	A	803	1/1	0.98	0.08	35,35,35,35	0
3	NA	A	804	1/1	0.99	0.04	31,31,31,31	0
3	NA	A	802	1/1	0.99	0.06	17,17,17,17	0
2	ZN	A	801	1/1	0.99	0.02	25,25,25,25	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ACO A 805 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 7L1 A 806 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.