



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:36 PM UTC

PDB ID : 8BA4 / pdb_00008ba4
Title : Crystal structure of JAK2 JH2-V617F in complex with Bemcentinib
Authors : Haikarainen, T.; Silvennoinen, O.
Deposited on : 2022-10-11
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	289	5% 85% 9% • 5%
1	B	289	7% 88% 7% 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8804 atoms, of which 4244 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein kinase JAK2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	274	Total	C	H	N	O	S	0	0	0
			4316	1390	2141	372	402	11			
1	B	274	Total	C	H	N	O	S	0	0	0
			4213	1370	2069	362	402	10			

There are 32 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	617	PHE	VAL	variant	UNP O60674
A	659	ALA	TRP	engineered mutation	UNP O60674
A	777	ALA	TRP	engineered mutation	UNP O60674
A	794	HIS	PHE	engineered mutation	UNP O60674
A	813	LEU	-	expression tag	UNP O60674
A	814	VAL	-	expression tag	UNP O60674
A	815	PRO	-	expression tag	UNP O60674
A	816	ARG	-	expression tag	UNP O60674
A	817	GLY	-	expression tag	UNP O60674
A	818	SER	-	expression tag	UNP O60674
A	819	HIS	-	expression tag	UNP O60674
A	820	HIS	-	expression tag	UNP O60674
A	821	HIS	-	expression tag	UNP O60674
A	822	HIS	-	expression tag	UNP O60674
A	823	HIS	-	expression tag	UNP O60674
A	824	HIS	-	expression tag	UNP O60674
B	617	PHE	VAL	variant	UNP O60674
B	659	ALA	TRP	engineered mutation	UNP O60674
B	777	ALA	TRP	engineered mutation	UNP O60674
B	794	HIS	PHE	engineered mutation	UNP O60674
B	813	LEU	-	expression tag	UNP O60674
B	814	VAL	-	expression tag	UNP O60674
B	815	PRO	-	expression tag	UNP O60674
B	816	ARG	-	expression tag	UNP O60674
B	817	GLY	-	expression tag	UNP O60674

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	818	SER	-	expression tag	UNP O60674
B	819	HIS	-	expression tag	UNP O60674
B	820	HIS	-	expression tag	UNP O60674
B	821	HIS	-	expression tag	UNP O60674
B	822	HIS	-	expression tag	UNP O60674
B	823	HIS	-	expression tag	UNP O60674
B	824	HIS	-	expression tag	UNP O60674

- Molecule 2 is 1-(3,4-diazatricyclo[9.4.0.0^{2,7}]pentadeca-1(11),2(7),3,5,12,14-hexaen-5-yl)-{N}3-[(7 {S})-7-pyrrolidin-1-yl-6,7,8,9-tetrahydro-5 {H}-benzo[7]annulen-3-yl]-1,2,4-triazole-3,5-diamine (CCD ID: Q8U) (formula: C₃₀H₃₄N₈) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	N	0	0
			72	30	34	8		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	106	Total	O	0	0
			106	106		
3	B	97	Total	O	0	0
			97	97		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein kinase JAK2



- Molecule 1: Tyrosine-protein kinase JAK2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	55.57Å 57.09Å 60.71Å 90.06° 116.01° 115.64°	Depositor
Resolution (Å)	50.02 – 2.10 50.02 – 2.10	Depositor EDS
% Data completeness (in resolution range)	95.6 (50.02-2.10) 97.9 (50.02-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.82 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.216 , 0.274 0.216 , 0.274	Depositor DCC
R_{free} test set	1682 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtriage
Anisotropy	0.121	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 33.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8804	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: Q8U

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.13	0/2224	0.30	0/3013
1	B	0.12	0/2193	0.28	0/2980
All	All	0.12	0/4417	0.29	0/5993

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2175	2141	2142	16	0
1	B	2144	2069	2069	11	0
2	A	38	34	0	0	0
3	A	106	0	0	0	0
3	B	97	0	0	0	0
All	All	4560	4244	4211	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:581:LYS:NZ	1:A:699:ASP:OD2	2.31	0.63
1:B:615:VAL:HG12	1:B:624:LEU:HD23	1.82	0.61
1:B:555:THR:OG1	1:B:715:ARG:NH2	2.34	0.58
1:A:670:ILE:HG21	1:A:730:LEU:HD21	1.87	0.55
1:A:555:THR:HG1	1:A:715:ARG:HH21	1.53	0.54
1:A:704:ILE:HD11	1:A:730:LEU:HD13	1.90	0.54
1:B:617:PHE:O	1:B:619:GLY:N	2.41	0.53
1:A:604:LEU:HD11	1:A:669:LEU:HD12	1.91	0.52
1:A:555:THR:OG1	1:A:715:ARG:NH2	2.37	0.52
1:A:618:CYS:HB2	1:A:623:ILE:HD12	1.95	0.49
1:A:670:ILE:CG2	1:A:730:LEU:HD21	2.42	0.48
1:A:579:LEU:HD13	1:A:580:LEU:N	2.29	0.47
1:A:661:MET:HE2	1:A:735:ASP:HB3	1.97	0.46
1:B:629:VAL:HG11	1:B:680:LEU:HB3	1.99	0.45
1:B:583:LEU:HD22	1:B:591:SER:HA	1.99	0.44
1:A:619:GLY:O	1:A:621:GLU:N	2.51	0.43
1:B:635:ASP:OD1	1:B:636:THR:N	2.53	0.42
1:B:713:GLN:NE2	1:B:763:LEU:HD11	2.35	0.42
1:A:743:LEU:HD21	1:A:780:LEU:HB3	2.02	0.42
1:A:598:ALA:O	1:A:612:ASN:ND2	2.53	0.41
1:B:615:VAL:HG12	1:B:624:LEU:CD2	2.48	0.41
1:B:595:PHE:HD1	1:B:624:LEU:HD21	1.86	0.41
1:B:704:ILE:HD11	1:B:730:LEU:HD13	2.01	0.41
1:A:739:PHE:CE2	1:A:743:LEU:HD13	2.55	0.41
1:A:670:ILE:HG21	1:A:730:LEU:CD2	2.50	0.41
1:B:661:MET:HE2	1:B:735:ASP:HB3	2.03	0.40
1:A:594:PHE:CZ	1:A:624:LEU:HD12	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/289 (94%)	265 (97%)	6 (2%)	1 (0%)	30	28
1	B	272/289 (94%)	265 (97%)	6 (2%)	1 (0%)	30	28
All	All	544/578 (94%)	530 (97%)	12 (2%)	2 (0%)	30	28

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	618	CYS
1	A	620	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	238/259 (92%)	234 (98%)	4 (2%)	53	62
1	B	231/259 (89%)	231 (100%)	0	100	100
All	All	469/518 (90%)	465 (99%)	4 (1%)	70	78

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	605	SER
1	A	699	ASP
1	A	728	LYS
1	A	743	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	574	HIS
1	A	662	HIS
1	A	729	ASN
1	A	760	GLN
1	B	538	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	548	ASN
1	B	587	HIS
1	B	713	GLN
1	B	770	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	Q8U	A	901	-	44,44,44	3.77	21 (47%)	55,63,63	2.30	10 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	Q8U	A	901	-	-	0/8/39/39	1/7/7/7

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	901	Q8U	C18-N17	10.80	1.54	1.36
2	A	901	Q8U	C03-N04	-9.91	1.23	1.47
2	A	901	Q8U	C06-N04	-8.50	1.27	1.48
2	A	901	Q8U	C21-N23	8.24	1.48	1.34
2	A	901	Q8U	C01-C05	-6.45	1.29	1.51
2	A	901	Q8U	C07-C06	5.87	1.68	1.53
2	A	901	Q8U	C29-C28	5.61	1.48	1.39
2	A	901	Q8U	C38-C27	-4.69	1.44	1.49
2	A	901	Q8U	C05-N04	4.10	1.57	1.47
2	A	901	Q8U	C24-N20	3.44	1.47	1.41
2	A	901	Q8U	C16-C09	3.41	1.45	1.39
2	A	901	Q8U	C13-C10	3.38	1.44	1.39
2	A	901	Q8U	C16-C15	2.87	1.43	1.38
2	A	901	Q8U	C27-N26	2.74	1.38	1.33
2	A	901	Q8U	C11-C10	2.69	1.57	1.51
2	A	901	Q8U	C36-C35	2.66	1.44	1.38
2	A	901	Q8U	C12-C06	2.54	1.59	1.53
2	A	901	Q8U	N20-N19	-2.51	1.33	1.39
2	A	901	Q8U	C21-N22	-2.45	1.29	1.33
2	A	901	Q8U	C30-C28	2.29	1.56	1.51
2	A	901	Q8U	C13-C14	2.04	1.42	1.39

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	901	Q8U	N19-C18-N22	-6.72	107.72	116.67
2	A	901	Q8U	C18-N19-N20	6.24	107.85	100.66
2	A	901	Q8U	C24-N20-N19	5.67	126.12	117.84
2	A	901	Q8U	C27-C38-C33	5.57	124.28	120.66
2	A	901	Q8U	N25-C24-N20	5.50	119.27	116.02
2	A	901	Q8U	C05-N04-C06	5.29	119.23	114.12
2	A	901	Q8U	C24-N25-N26	4.17	123.18	118.97
2	A	901	Q8U	C12-C11-C10	2.31	118.52	113.48
2	A	901	Q8U	C11-C10-C13	-2.15	114.69	119.35
2	A	901	Q8U	C29-C24-N25	-2.07	120.77	123.82

There are no chirality outliers.


There are no torsion outliers.

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	901	Q8U	C06-C07-C08-C09-C10-C11-C12

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/289 (94%)	0.36	14 (5%) 33 35	28, 46, 79, 114	0
1	B	274/289 (94%)	0.38	20 (7%) 21 22	27, 45, 78, 104	0
All	All	548/578 (94%)	0.37	34 (6%) 26 28	27, 46, 79, 114	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	809	PHE	5.5
1	A	809	PHE	5.4
1	B	617	PHE	4.8
1	B	536	VAL	4.4
1	A	617	PHE	4.0
1	A	730	LEU	3.9
1	B	594	PHE	3.8
1	B	595	PHE	3.6
1	B	598	ALA	3.5
1	A	619	GLY	3.4
1	A	536	VAL	3.1
1	A	604	LEU	3.1
1	A	728	LYS	2.8
1	B	622	ASN	2.8
1	B	599	SER	2.8
1	A	618	CYS	2.7
1	B	601	MET	2.6
1	B	624	LEU	2.6
1	B	616	CYS	2.5
1	A	731	ASN	2.5
1	B	604	LEU	2.5
1	B	687	ARG	2.4
1	B	626	GLN	2.3
1	B	583	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	620	ASP	2.2
1	B	537	PHE	2.2
1	A	538	HIS	2.2
1	B	688	LYS	2.2
1	A	586	ALA	2.2
1	B	586	ALA	2.1
1	A	622	ASN	2.1
1	A	537	PHE	2.1
1	B	618	CYS	2.0
1	B	615	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	Q8U	A	901	38/38	0.87	0.13	36,58,108,131	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.