



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:51 AM UTC

PDB ID : 3BBL / pdb\_00003bbl  
Title : Crystal structure of a regulatory protein of LacI family from *Chloroflexus aggregans*  
Authors : Kumaran, D.; Burley, S.K.; Swaminathan, S.; New York SGX Research Center for Structural Genomics (NYSGXRC)  
Deposited on : 2007-11-09  
Resolution : 2.35 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	287	

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2331 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Regulatory protein of LacI family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	281	2219	1413	389	408	1	8	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	60	MSE	-	expression tag	UNP A0H3S5
A	61	SER	-	expression tag	UNP A0H3S5
A	62	LEU	-	expression tag	UNP A0H3S5
A	339	GLU	-	expression tag	UNP A0H3S5
A	340	GLY	-	expression tag	UNP A0H3S5
A	341	HIS	-	expression tag	UNP A0H3S5
A	342	HIS	-	expression tag	UNP A0H3S5
A	343	HIS	-	expression tag	UNP A0H3S5
A	344	HIS	-	expression tag	UNP A0H3S5
A	345	HIS	-	expression tag	UNP A0H3S5
A	346	HIS	-	expression tag	UNP A0H3S5

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	96	Total O 96 96	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Regulatory protein of LacI family



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 63 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	122.55Å 122.55Å 123.30Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	35.38 – 2.35 35.38 – 2.35	Depositor EDS
% Data completeness (in resolution range)	93.9 (35.38-2.35) 93.9 (35.38-2.35)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.90 (at 2.16Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, $R_{free}$	0.240 , 0.258 0.230 , 0.258	Depositor DCC
$R_{free}$ test set	1063 reflections (4.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.6	Xtrriage
Anisotropy	0.349	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 39.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2331	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.45	0/2262	0.88	4/3057 (0.1%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	VAL	N-CA-C	-5.76	100.04	108.11
1	A	160	ASP	N-CA-C	5.54	118.27	109.24
1	A	121	SER	N-CA-C	-5.28	106.79	113.18
1	A	163	TRP	N-CA-C	5.13	116.63	108.79

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2219	0	2196	54	0
2	A	16	0	24	0	0
3	A	96	0	0	8	0
All	All	2331	0	2220	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:LEU:HD22	1:A:235:MSE:HE3	1.53	0.87
1:A:109:ARG:HH21	1:A:131:SER:HB3	1.38	0.86
1:A:65:MSE:HE2	1:A:101:LEU:HB2	1.59	0.82
1:A:255:ASN:C	1:A:255:ASN:HD22	1.90	0.78
1:A:255:ASN:ND2	1:A:258:MSE:H	1.83	0.76
1:A:336:ARG:HG2	1:A:341:HIS:HB3	1.70	0.73
1:A:156:ASN:HB3	3:A:536:HOH:O	1.88	0.72
1:A:255:ASN:HD21	1:A:258:MSE:H	1.39	0.69
1:A:324:GLN:OE1	1:A:324:GLN:HA	1.94	0.68
1:A:65:MSE:HE1	1:A:101:LEU:HD13	1.77	0.66
1:A:255:ASN:C	1:A:255:ASN:ND2	2.58	0.62
1:A:255:ASN:HD21	1:A:258:MSE:HG3	1.64	0.61
1:A:198:VAL:HG23	3:A:563:HOH:O	2.00	0.60
1:A:282:PHE:O	1:A:283:ASP:CG	2.45	0.59
1:A:159:TRP:HB2	3:A:536:HOH:O	2.04	0.58
1:A:190:LEU:HB3	1:A:235:MSE:CE	2.35	0.57
1:A:182:ARG:NH2	3:A:594:HOH:O	2.37	0.57
1:A:338:SER:O	1:A:339:GLU:HG3	2.05	0.57
1:A:255:ASN:ND2	1:A:258:MSE:HG3	2.19	0.57
1:A:190:LEU:HB3	1:A:235:MSE:HE1	1.87	0.57
1:A:336:ARG:HE	1:A:341:HIS:CE1	2.23	0.56
1:A:79:ASN:ND2	3:A:597:HOH:O	2.42	0.53
1:A:112:ILE:HD12	1:A:141:PHE:CD1	2.44	0.52
1:A:74:GLU:H	1:A:74:GLU:CD	2.19	0.51
1:A:173:ARG:O	1:A:177:GLU:HG3	2.11	0.51
1:A:78:VAL:HG12	1:A:285:ALA:CB	2.41	0.51
1:A:283:ASP:O	1:A:284:ASP:C	2.55	0.50
1:A:109:ARG:HH11	1:A:109:ARG:HG3	1.77	0.49
1:A:337:ALA:O	1:A:339:GLU:N	2.45	0.49
1:A:112:ILE:HD12	1:A:141:PHE:CG	2.49	0.48
1:A:73:THR:HG21	1:A:80:HIS:CE1	2.47	0.48
1:A:140:GLN:HG2	1:A:144:LYS:HE3	1.96	0.48
1:A:132:ILE:HG22	1:A:155:SER:HB2	1.96	0.47
1:A:92:GLU:HG3	1:A:307:ARG:HA	1.97	0.47
1:A:108:ASP:OD2	1:A:111:GLN:HG3	2.14	0.47
1:A:191:ALA:HA	1:A:254:LEU:HG	1.97	0.47
1:A:79:ASN:CG	3:A:597:HOH:O	2.58	0.47
1:A:182:ARG:HD2	3:A:508:HOH:O	2.14	0.47
1:A:132:ILE:CG2	1:A:155:SER:HB2	2.45	0.46
1:A:79:ASN:HB2	1:A:285:ALA:HB2	1.97	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:255:ASN:HA	1:A:282:PHE:O	2.16	0.46
1:A:65:MSE:CE	1:A:101:LEU:HD13	2.43	0.46
1:A:218:GLU:CD	1:A:246:ARG:HH22	2.24	0.45
1:A:286:PRO:O	1:A:287:MSE:CB	2.65	0.44
1:A:229:PHE:CE1	1:A:261:GLY:HA2	2.53	0.43
1:A:231:VAL:O	1:A:235:MSE:HG3	2.18	0.43
1:A:191:ALA:HB1	3:A:505:HOH:O	2.17	0.43
1:A:282:PHE:C	1:A:283:ASP:CG	2.86	0.43
1:A:230:GLU:H	1:A:230:GLU:CD	2.26	0.42
1:A:192:TRP:HB3	1:A:193:PRO:CD	2.51	0.41
1:A:310:ILE:O	1:A:314:VAL:HG23	2.21	0.41
1:A:324:GLN:NE2	1:A:326:HIS:O	2.52	0.40
1:A:337:ALA:O	1:A:338:SER:C	2.64	0.40
1:A:223:LEU:HD12	1:A:235:MSE:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	279/287 (97%)	265 (95%)	9 (3%)	5 (2%)	<b>6</b> <b>5</b>

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	338	SER
1	A	157	PRO
1	A	156	ASN
1	A	283	ASP
1	A	340	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	235/232 (101%)	226 (96%)	9 (4%)	29 39

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	GLU
1	A	78	VAL
1	A	157	PRO
1	A	246	ARG
1	A	255	ASN
1	A	283	ASP
1	A	287	MSE
1	A	324	GLN
1	A	341	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	80	HIS
1	A	84	GLN
1	A	140	GLN
1	A	255	ASN
1	A	289	GLN
1	A	341	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	EDO	A	501	-	3,3,3	0.48	0	2,2,2	0.23	0
2	EDO	A	502	-	3,3,3	0.49	0	2,2,2	0.32	0
2	EDO	A	500	-	3,3,3	0.48	0	2,2,2	0.24	0
2	EDO	A	503	-	3,3,3	0.44	0	2,2,2	0.33	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	501	-	-	1/1/1/1	-
2	EDO	A	502	-	-	0/1/1/1	-
2	EDO	A	500	-	-	1/1/1/1	-
2	EDO	A	503	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	500	EDO	O1-C1-C2-O2
2	A	501	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	273/287 (95%)	0.58	37 (13%) <b>7</b>   <b>7</b>	15, 30, 64, 79	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	342	HIS	7.2
1	A	78	VAL	6.9
1	A	340	GLY	6.9
1	A	75	PRO	6.3
1	A	157	PRO	5.8
1	A	338	SER	5.7
1	A	341	HIS	5.6
1	A	159	TRP	4.8
1	A	158	ASP	4.6
1	A	79	ASN	4.4
1	A	283	ASP	4.4
1	A	74	GLU	4.2
1	A	80	HIS	4.1
1	A	324	GLN	4.1
1	A	339	GLU	4.0
1	A	77	GLN	3.6
1	A	337	ALA	3.6
1	A	156	ASN	3.5
1	A	81	ILE	3.2
1	A	76	GLY	3.2
1	A	111	GLN	3.2
1	A	160	ASP	3.1
1	A	144	LYS	3.0
1	A	322	PRO	2.9
1	A	140	GLN	2.9
1	A	120	ARG	2.8
1	A	108	ASP	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	135	ASN	2.6
1	A	323	GLU	2.5
1	A	73	THR	2.5
1	A	123	ASN	2.4
1	A	121	SER	2.3
1	A	95	ALA	2.2
1	A	155	SER	2.2
1	A	185	ARG	2.1
1	A	325	LYS	2.1
1	A	72	GLN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	EDO	A	502	4/4	0.88	0.21	37,41,41,42	0
2	EDO	A	503	4/4	0.89	0.16	32,33,33,37	0
2	EDO	A	501	4/4	0.91	0.12	28,28,29,30	0
2	EDO	A	500	4/4	0.96	0.07	22,24,25,28	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.