



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:41 PM UTC

PDB ID : 2BCQ / pdb_00002bcq
Title : DNA polymerase lambda in complex with a DNA duplex containing an unpaired Dtmp
Authors : Garcia-Diaz, M.; Bebenek, K.; Krahn, J.M.; Pedersen, L.C.; Kunkel, T.A.
Deposited on : 2005-10-19
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.66-1.66)
Clashscore	190562	2662 (1.66-1.66)
Ramachandran outliers	187476	2621 (1.66-1.66)
Sidechain outliers	187428	2621 (1.66-1.66)
RSRZ outliers	180081	2564 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	4	50% 50%
2	P	7	100%
3	T	12	92% 8%
4	A	335	18% 79% 18% .

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 3624 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(P*GP*CP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	4	Total	C	N	O	P	0	0	0
			83	38	16	25	4			

- Molecule 2 is a DNA chain called 5'-D(*CP*AP*GP*TP*AP*CP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	P	7	Total	C	N	O	P	0	0	0
			141	68	28	39	6			

- Molecule 3 is a DNA chain called 5'-D(*CP*GP*GP*CP*CP*GP*TP*TP*AP*CP*TP*G)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	T	12	Total	C	N	O	P	0	0	0
			242	116	43	72	11			

- Molecule 4 is a protein called DNA polymerase lambda.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	A	324	Total	C	N	O	S	0	5	0
			2571	1615	470	475	11			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	241	MET	-	initiating methionine	UNP Q9UGP5


- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	3	Total	Na	0	0
			3	3		

- Molecule 6 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Mg	0	0
			1	1		

- Molecule 7 is PYROPHOSPHATE (CCD ID: PPV) (formula: $\text{H}_4\text{O}_7\text{P}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	O	P	0	0
			9	7	2		

- Molecule 8 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $\text{C}_2\text{H}_6\text{O}_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	D	22	Total	O	0	0
			22	22		
9	P	49	Total	O	0	0
			49	49		
9	T	72	Total	O	0	0
			72	72		
9	A	427	Total	O	0	0
			427	427		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5'-D(P*GP*CP*CP*G)-3'




- Molecule 2: 5'-D(*CP*AP*GP*TP*AP*CP*G)-3'




There are no outlier residues recorded for this chain.

- Molecule 3: 5'-D(*CP*GP*GP*CP*CP*GP*TP*TP*AP*CP*TP*G)-3'



- Molecule 4: DNA polymerase lambda



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.04Å 62.54Å 139.64Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.74 – 1.65 41.74 – 1.65	Depositor EDS
% Data completeness (in resolution range)	92.3 (41.74-1.65) 92.3 (41.74-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.61 (at 1.64Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.201 , 0.230 0.206 , 0.232	Depositor DCC
R_{free} test set	2873 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	19.7	Xtriage
Anisotropy	0.484	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3624	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.06% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, MG, NA, PPV

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.47	0/92	0.52	0/138
2	P	0.29	0/158	0.57	0/242
3	T	0.28	0/270	0.70	0/415
4	A	0.31	0/2624	0.79	2/3542 (0.1%)
All	All	0.32	0/3144	0.76	2/4337 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	T	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
4	A	273	LYS	N-CA-C	6.82	118.72	111.28
4	A	531	ALA	N-CA-C	5.24	116.10	108.14

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	T	11	DT	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	83	0	45	1	0
2	P	141	0	80	0	0
3	T	242	0	137	0	0
4	A	2571	0	2544	54	0
5	A	3	0	0	0	0
6	A	1	0	0	0	0
7	A	9	0	0	0	0
8	A	4	0	6	1	0
9	A	427	0	0	7	0
9	D	22	0	0	0	0
9	P	49	0	0	0	0
9	T	72	0	0	0	0
All	All	3624	0	2812	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:467:ASN:HD21	4:A:469:GLN:HB2	1.50	0.75
4:A:256:ILE:HD12	4:A:257:THR:N	2.04	0.72
4:A:519:LEU:HD23	4:A:565:LEU:HD11	1.71	0.72
4:A:312[B]:LYS:HE2	9:A:3152:HOH:O	1.95	0.67
4:A:396:GLU:HG3	4:A:414:ALA:HB2	1.77	0.65
4:A:467:ASN:ND2	4:A:469:GLN:HB2	2.13	0.63
4:A:256:ILE:HD13	4:A:286:LEU:HD13	1.80	0.63
4:A:256:ILE:CD1	4:A:286:LEU:HD13	2.29	0.61
4:A:346[B]:THR:HG23	9:A:2839:HOH:O	2.02	0.60
4:A:537:VAL:HB	4:A:546:GLY:HA3	1.85	0.58
4:A:313:ILE:O	4:A:317:LEU:HD13	2.04	0.58
4:A:256:ILE:HG22	4:A:316:ILE:CG2	2.34	0.57
4:A:346[A]:THR:O	4:A:350:GLN:HG3	2.05	0.57
4:A:431:LEU:C	4:A:431:LEU:HD23	2.30	0.56
4:A:290:HIS:CG	4:A:291:LYS:H	2.24	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:471:GLN:HB3	8:A:2801:EDO:H22	1.88	0.55
4:A:256:ILE:HG22	4:A:316:ILE:HG21	1.89	0.55
4:A:312[A]:LYS:O	4:A:316:ILE:HG13	2.08	0.53
4:A:516:MET:HE1	4:A:560:PHE:CE2	2.44	0.53
4:A:437:GLY:HA2	9:A:2841:HOH:O	2.09	0.53
4:A:308:ARG:HG3	4:A:308:ARG:HH11	1.75	0.51
4:A:289:PHE:HB3	9:A:2922:HOH:O	2.10	0.51
4:A:265:LYS:HD2	4:A:336:GLU:OE1	2.11	0.51
4:A:572:GLU:HG3	9:A:3111:HOH:O	2.11	0.51
4:A:568:ARG:NH1	4:A:575:TRP:HZ2	2.10	0.50
4:A:312[B]:LYS:O	4:A:316:ILE:HG13	2.12	0.49
4:A:346[B]:THR:O	4:A:350:GLN:HG3	2.11	0.49
4:A:290:HIS:CG	4:A:291:LYS:N	2.79	0.49
4:A:308:ARG:O	4:A:312[B]:LYS:HG2	2.13	0.49
4:A:316:ILE:O	4:A:320:GLY:HA2	2.12	0.49
4:A:351:MET:O	4:A:355:GLN:HG3	2.13	0.48
4:A:292:PRO:O	4:A:294:THR:HG23	2.13	0.48
1:D:3:DC:H2"	1:D:4:DG:C8	2.50	0.47
4:A:519:LEU:CD2	4:A:565:LEU:HD11	2.43	0.47
4:A:255:HIS:ND1	4:A:256:ILE:HG23	2.30	0.46
4:A:253:ASN:O	4:A:256:ILE:HG13	2.16	0.46
4:A:308:ARG:O	4:A:312[A]:LYS:HG3	2.16	0.46
4:A:315:GLU:HG2	4:A:321:HIS:O	2.16	0.46
4:A:256:ILE:HG21	4:A:317:LEU:HD11	1.99	0.45
4:A:519:LEU:HD21	4:A:565:LEU:HD21	1.98	0.45
4:A:350:GLN:HG2	9:A:2901:HOH:O	2.16	0.45
4:A:554:PRO:HG2	4:A:558:ASP:OD2	2.17	0.45
4:A:332:VAL:HB	4:A:333:PRO:HD3	1.98	0.44
4:A:256:ILE:HD12	4:A:257:THR:HG23	2.00	0.43
4:A:396:GLU:CG	4:A:414:ALA:HB2	2.45	0.43
4:A:310:ALA:O	4:A:314:ILE:HG12	2.18	0.43
4:A:287:LYS:C	4:A:289:PHE:H	2.26	0.43
4:A:505:TYR:CG	4:A:529:GLU:HB3	2.55	0.42
4:A:265:LYS:HE2	9:A:2960:HOH:O	2.18	0.42
4:A:434:HIS:HA	4:A:435:PRO:HD3	1.92	0.42
4:A:431:LEU:HD23	4:A:432:ILE:N	2.35	0.42
4:A:467:ASN:CG	4:A:469:GLN:H	2.28	0.41
4:A:259:LYS:NZ	4:A:321:HIS:HA	2.36	0.41
4:A:261[A]:GLU:CG	4:A:283:ILE:HD13	2.52	0.40
4:A:309:MET:O	4:A:313:ILE:HG13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	A	327/335 (98%)	315 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	A	270/281 (96%)	269 (100%)	1 (0%)	84	77

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	A	506	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
4	A	270	GLN
4	A	355	GLN
4	A	366	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	PPV	A	738	6	6,8,8	0.94	1 (16%)	12,13,13	1.32	1 (8%)
8	EDO	A	2801	-	3,3,3	0.52	0	2,2,2	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	PPV	A	738	6	-	0/6/6/6	-
8	EDO	A	2801	-	-	1/1/1/1	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	A	738	PPV	P2-O12	-2.12	1.46	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	738	PPV	O32-P2-OPP	-2.31	96.90	104.64

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	2801	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	2801	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	D	4/4 (100%)	0.29	0	100 100	28, 28, 28, 28	0
2	P	7/7 (100%)	-0.96	0	100 100	13, 14, 17, 18	0
3	T	12/12 (100%)	-0.37	0	100 100	15, 18, 29, 39	0
4	A	324/335 (96%)	0.92	61 (18%)	3 4	9, 26, 65, 81	5 (1%)
All	All	347/358 (96%)	0.83	61 (17%)	4 4	9, 25, 64, 81	5 (1%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	A	293	VAL	8.0
4	A	256	ILE	6.4
4	A	302	ILE	5.6
4	A	286	LEU	5.2
4	A	252	HIS	5.0
4	A	545	VAL	5.0
4	A	299	ALA	5.0
4	A	292	PRO	4.9
4	A	317	LEU	4.8
4	A	294	THR	4.8
4	A	313	ILE	4.7
4	A	289	PHE	4.6
4	A	255	HIS	4.5
4	A	254	LEU	4.5
4	A	300	CYS	4.2
4	A	316	ILE	4.2
4	A	546	GLY	4.1
4	A	540	THR	4.0
4	A	314	ILE	3.8
4	A	291	LYS	3.7
4	A	319	SER	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	A	305	ILE	3.6
4	A	285	ALA	3.5
4	A	296	TYR	3.5
4	A	295	SER	3.5
4	A	322	LEU	3.5
4	A	405	ALA	3.4
4	A	575	TRP	3.4
4	A	284	ASN	3.3
4	A	288	SER	3.3
4	A	290	HIS	3.3
4	A	303	PRO	3.1
4	A	543	CYS	3.1
4	A	537	VAL	3.0
4	A	257	THR	3.0
4	A	310	ALA	3.0
4	A	406	PHE	2.9
4	A	467	ASN	2.9
4	A	253	ASN	2.9
4	A	298	GLU	2.8
4	A	320	GLY	2.7
4	A	442	GLY	2.7
4	A	536	VAL	2.6
4	A	283	ILE	2.6
4	A	411	LEU	2.6
4	A	311	GLU	2.6
4	A	301	SER	2.6
4	A	511[A]	HIS	2.5
4	A	453	GLN	2.4
4	A	541	HIS	2.4
4	A	304	GLY	2.3
4	A	325	LEU	2.3
4	A	443	ILE	2.3
4	A	297	GLN	2.2
4	A	308	ARG	2.2
4	A	539	ASN	2.2
4	A	306	GLY	2.2
4	A	282	ALA	2.2
4	A	437	GLY	2.1
4	A	318	GLU	2.1
4	A	538	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	EDO	A	2801	4/4	0.81	0.21	40,40,40,42	0
5	NA	A	712	1/1	0.89	0.11	59,59,59,59	0
7	PPV	A	738	9/9	0.92	0.11	25,29,32,32	0
5	NA	A	4	1/1	0.97	0.05	27,27,27,27	0
5	NA	A	1	1/1	0.98	0.06	15,15,15,15	0
6	MG	A	713	1/1	0.98	0.04	21,21,21,21	0

6.5 Other polymers [i](#)

There are no such residues in this entry.