



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:13 PM UTC

PDB ID : 4BDT / pdb_00004bdt
Title : Human acetylcholinesterase in complex with huprine W and fasciculin 2
Authors : Nachon, F.; Carletti, E.; Ronco, C.; Trovaslet, M.; Nicolet, Y.; Jean, L.; Renard, P.-Y.
Deposited on : 2012-10-06
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	583	% 69% 24% . .
2	B	61	49% 43% 8%
3	C	3	33% 67%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 5057 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ACETYLCHOLINESTERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	563	Total	C	N	O	S	17	0	0
			4399	2822	770	793	14			

- Molecule 2 is a protein called FASCICULIN-2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	61	Total	C	N	O	S	0	0	0
			464	276	88	90	10			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	3	Total	C	N	O	0	0	0
			38	22	2	14			

- Molecule 4 is HUPRINE W (CCD ID: HUW) (formula: C₁₈H₁₉ClN₂O).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	Cl	N	O	0	0
			22	18	1	2	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	10	Total	Cl	0	0
			10	10		

- Molecule 6 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	113	Total	O	0	0
			113	113		
7	B	6	Total	O	0	0
			6	6		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ACETYLCHOLINESTERASE



• Molecule 2: FASCICULIN-2



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[beta-L-fucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	151.60Å 151.60Å 246.40Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	57.93 – 3.10 57.93 – 3.10	Depositor EDS
% Data completeness (in resolution range)	98.3 (57.93-3.10) 98.4 (57.93-3.10)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.79 (at 3.13Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.159 , 0.219 0.165 , 0.222	Depositor DCC
R_{free} test set	984 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	66.4	Xtriage
Anisotropy	0.723	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 65.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5057	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, HUW, FUL, NAG, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.52	0/4538	0.95	12/6204 (0.2%)
2	B	0.57	1/473 (0.2%)	0.99	2/636 (0.3%)
All	All	0.53	1/5011 (0.0%)	0.95	14/6840 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	30	PRO	CA-C	5.17	1.54	1.51

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	264	GLY	N-CA-C	-6.36	102.88	112.60
1	A	450	GLU	N-CA-C	-6.34	105.68	113.41
1	A	87	ASN	CA-C-N	6.15	127.53	119.84
1	A	87	ASN	C-N-CA	6.15	127.53	119.84
1	A	234	GLY	CA-C-N	6.14	125.83	119.56
1	A	234	GLY	C-N-CA	6.14	125.83	119.56
1	A	329	GLY	N-CA-C	6.09	120.54	110.55
1	A	112	THR	CA-C-N	6.07	126.09	119.90
1	A	112	THR	C-N-CA	6.07	126.09	119.90
1	A	439	TRP	CA-C-N	5.69	126.95	119.84
1	A	439	TRP	C-N-CA	5.69	126.95	119.84
2	B	41	CYS	CA-C-N	5.69	126.24	120.38
2	B	41	CYS	C-N-CA	5.69	126.24	120.38
1	A	336	SER	N-CA-C	5.39	117.60	111.02

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4399	0	4252	87	0
2	B	464	0	442	16	0
3	C	38	0	34	1	0
4	A	22	0	19	1	0
5	A	10	0	0	1	0
6	A	5	0	0	0	0
7	A	113	0	0	7	0
7	B	6	0	0	0	0
All	All	5057	0	4747	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:56:PRO:HB2	2:B:59:CYS:HB3	1.54	0.88
1:A:113:PRO:HG2	1:A:485:ARG:HG2	1.73	0.71
1:A:75:THR:O	2:B:37:ARG:NH2	2.27	0.67
1:A:255:VAL:HG22	1:A:280:VAL:HG12	1.78	0.65
1:A:417:ARG:NH1	1:A:421:GLN:OE1	2.31	0.64
1:A:265:ASN:CG	1:A:266:ASP:HA	2.25	0.62
2:B:25:LYS:HD3	2:B:48:LEU:HD13	1.82	0.61
2:B:37:ARG:NH1	2:B:61:TYR:O	2.34	0.60
1:A:491:GLU:O	7:A:2106:HOH:O	2.17	0.59
1:A:329:GLY:HA3	1:A:428:TYR:CE1	2.38	0.58
1:A:511:VAL:HB	1:A:518:LEU:HD22	1.86	0.58
1:A:374:ALA:HA	1:A:539:LEU:HD21	1.87	0.57
2:B:8:THR:HG23	2:B:9:THR:HG23	1.86	0.56
1:A:121:GLY:HA2	4:A:701:H UW:H92C	1.88	0.56
1:A:266:ASP:O	1:A:270:VAL:HG23	2.06	0.56
1:A:460:ASP:HB3	1:A:463:ARG:HG3	1.88	0.55
1:A:417:ARG:NH2	5:A:803:CL:CL	2.76	0.55
2:B:27:ARG:NH1	2:B:30:PRO:O	2.40	0.55
1:A:294:VAL:HG13	1:A:338:PHE:HB3	1.91	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:VAL:HG22	1:A:327:LEU:HB3	1.90	0.53
1:A:207:ALA:O	1:A:211:MET:HG3	2.09	0.52
1:A:507:ALA:HA	1:A:522:ARG:NH1	2.24	0.52
1:A:77:TYR:CZ	1:A:348:LYS:HG2	2.45	0.51
1:A:512:SER:O	7:A:2103:HOH:O	2.19	0.51
1:A:437:LEU:HD23	1:A:438:SER:H	1.76	0.51
1:A:120:GLY:HA3	1:A:203:SER:HB3	1.92	0.50
1:A:68:VAL:HG13	1:A:127:ALA:HB2	1.92	0.50
1:A:7:GLU:HA	7:A:2002:HOH:O	2.10	0.49
2:B:4:TYR:CE2	2:B:58:LYS:HG2	2.47	0.49
2:B:7:THR:H	2:B:10:SER:HB3	1.77	0.49
1:A:536:LEU:HD12	1:A:539:LEU:HD23	1.94	0.49
3:C:1:NAG:H62	3:C:2:NAG:H82	1.94	0.49
1:A:304:ASP:OD1	1:A:304:ASP:N	2.44	0.49
1:A:236:TRP:CD1	1:A:236:TRP:H	2.30	0.48
1:A:254:LEU:HB3	1:A:284:HIS:HD2	1.78	0.48
1:A:265:ASN:OD1	1:A:266:ASP:HA	2.13	0.48
1:A:111:PRO:HB2	1:A:193:ASP:HB2	1.94	0.48
1:A:177:ARG:CZ	1:A:217:PRO:HB2	2.43	0.47
1:A:75:THR:HA	7:A:2022:HOH:O	2.14	0.47
1:A:193:ASP:OD1	1:A:195:THR:OG1	2.30	0.47
1:A:245:ARG:NH2	1:A:266:ASP:OD2	2.47	0.47
1:A:177:ARG:NH1	7:A:2049:HOH:O	2.45	0.47
1:A:434:ALA:O	1:A:437:LEU:HB2	2.15	0.46
1:A:24:THR:OG1	1:A:140:GLN:HG3	2.15	0.46
1:A:541:SER:O	1:A:545:THR:HG23	2.15	0.46
1:A:241:MET:HE3	1:A:241:MET:HB2	1.83	0.46
1:A:254:LEU:HB3	1:A:284:HIS:CD2	2.51	0.46
1:A:356:ARG:HA	1:A:394:LEU:HD13	1.97	0.46
2:B:30:PRO:HA	2:B:31:PRO:C	2.40	0.46
1:A:39:GLU:HA	1:A:40:PRO:HD2	1.79	0.45
1:A:69:CYS:HB2	7:A:2018:HOH:O	2.16	0.45
1:A:208:SER:O	1:A:211:MET:N	2.49	0.45
1:A:72:TYR:HE2	2:B:33:MET:HE2	1.82	0.45
1:A:468:GLU:H	1:A:468:GLU:HG3	1.40	0.45
1:A:329:GLY:HA3	1:A:428:TYR:CZ	2.53	0.44
1:A:73:VAL:HA	7:A:2019:HOH:O	2.17	0.44
1:A:499:GLN:O	1:A:501:PRO:HD3	2.18	0.44
1:A:162:PRO:HB3	1:A:241:MET:HG2	1.99	0.44
1:A:45:ARG:HA	1:A:48:LEU:HD12	1.99	0.43
1:A:24:THR:HB	1:A:25:PRO:HD2	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:364:ARG:NH1	1:A:370:VAL:O	2.42	0.43
2:B:1:THR:HG22	2:B:57:ASP:HA	2.00	0.43
2:B:6:HIS:HB3	2:B:12:ALA:HA	2.00	0.43
2:B:27:ARG:HD3	2:B:31:PRO:O	2.19	0.43
1:A:81:GLU:H	1:A:81:GLU:HG2	1.43	0.43
2:B:28:ARG:N	2:B:47:ASN:O	2.52	0.43
1:A:506:GLY:O	1:A:508:GLN:HG3	2.18	0.43
1:A:308:LEU:HD23	1:A:308:LEU:HA	1.82	0.43
1:A:275:THR:OG1	1:A:276:ARG:NH1	2.52	0.42
2:B:4:TYR:HB2	2:B:37:ARG:HD2	2.00	0.42
1:A:159:LEU:C	1:A:159:LEU:HD23	2.44	0.42
1:A:179:ALA:O	1:A:183:VAL:HG23	2.20	0.42
1:A:390:ASP:HA	1:A:391:PRO:HD3	1.87	0.42
1:A:452:GLU:OE2	1:A:465:TYR:OH	2.25	0.42
1:A:105:TYR:HA	1:A:106:PRO:HA	1.78	0.42
1:A:7:GLU:HG2	1:A:7:GLU:O	2.20	0.42
1:A:68:VAL:HG23	1:A:90:ARG:HB2	2.02	0.42
1:A:265:ASN:CB	1:A:266:ASP:HA	2.49	0.42
1:A:491:GLU:HB2	1:A:493:ARG:O	2.19	0.42
1:A:77:TYR:CE1	1:A:348:LYS:HG2	2.55	0.41
1:A:491:GLU:HG3	1:A:495:PRO:HG2	2.03	0.41
1:A:43:GLY:HA3	1:A:44:PRO:HD3	1.83	0.41
1:A:142:GLU:OE2	1:A:478:ARG:NH1	2.53	0.41
1:A:328:VAL:HG21	1:A:415:ALA:HB2	2.02	0.41
1:A:328:VAL:O	1:A:427:ALA:HA	2.20	0.41
1:A:482:ASN:O	1:A:486:THR:HG23	2.20	0.41
1:A:561:SER:O	1:A:565:VAL:HG13	2.21	0.41
2:B:6:HIS:CE1	2:B:36:GLY:HA2	2.56	0.41
1:A:218:SER:HA	1:A:221:LEU:HD12	2.02	0.41
1:A:463:ARG:HD2	1:A:465:TYR:HE1	1.86	0.41
1:A:497:ALA:HA	1:A:498:PRO:HD3	1.96	0.41
1:A:261:GLY:HA2	1:A:264:GLY:O	2.21	0.41
1:A:540:LEU:HD23	1:A:540:LEU:HA	1.92	0.41
1:A:40:PRO:HA	1:A:41:PRO:HD3	1.78	0.40
1:A:257:CYS:HA	1:A:258:PRO:HD3	1.79	0.40
1:A:107:ARG:HA	1:A:108:PRO:HD3	1.99	0.40
1:A:327:LEU:HD11	1:A:500:TRP:CH2	2.56	0.40
1:A:440:PRO:HB2	1:A:442:TRP:CD1	2.57	0.40
1:A:97:LEU:HD23	1:A:97:LEU:HA	1.78	0.40
1:A:474:GLN:HA	1:A:477:MET:HE2	2.04	0.40
1:A:102:TRP:CE3	1:A:146:LEU:HD12	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:ALA:O	1:A:326:VAL:HA	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	561/583 (96%)	518 (92%)	36 (6%)	7 (1%)	10	37
2	B	59/61 (97%)	52 (88%)	5 (8%)	2 (3%)	3	16
All	All	620/644 (96%)	570 (92%)	41 (7%)	9 (2%)	8	32

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	495	PRO
1	A	259	PRO
1	A	447	HIS
2	B	20	ASN
1	A	290	PRO
1	A	28	PRO
1	A	306	ASP
2	B	57	ASP
1	A	491	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	457/476 (96%)	417 (91%)	40 (9%)	9	33
2	B	55/55 (100%)	46 (84%)	9 (16%)	2	11
All	All	512/531 (96%)	463 (90%)	49 (10%)	8	30

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLU
1	A	33	LEU
1	A	59	VAL
1	A	60	VAL
1	A	81	GLU
1	A	83	THR
1	A	100	ASN
1	A	110	SER
1	A	112	THR
1	A	149	MET
1	A	161	LEU
1	A	165	ARG
1	A	187	VAL
1	A	200	PHE
1	A	243	GLU
1	A	250	GLN
1	A	255	VAL
1	A	266	ASP
1	A	267	THR
1	A	291	GLN
1	A	294	VAL
1	A	324	LEU
1	A	326	VAL
1	A	337	TYR
1	A	348	LYS
1	A	363	VAL
1	A	437	LEU
1	A	441	LEU
1	A	466	THR
1	A	468	GLU
1	A	471	ILE
1	A	486	THR
1	A	490	ASN
1	A	493	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	496	LYS
1	A	545	THR
1	A	554	LYS
1	A	559	ARG
1	A	564	MET
1	A	565	VAL
2	B	1	THR
2	B	7	THR
2	B	8	THR
2	B	11	ARG
2	B	14	LEU
2	B	19	GLU
2	B	21	SER
2	B	35	LEU
2	B	54	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	66	GLN
1	A	279	GLN
1	A	284	HIS
1	A	317	ASN
1	A	325	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	1,3	14,14,15	0.61	0	17,19,21	2.32	4 (23%)
3	NAG	C	2	3	14,14,15	0.71	0	17,19,21	1.09	2 (11%)
3	FUL	C	3	3	10,10,11	0.78	0	14,14,16	1.32	3 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	1,3	-	4/6/23/26	0/1/1/1
3	NAG	C	2	3	-	0/6/23/26	0/1/1/1
3	FUL	C	3	3	-	-	0/1/1/1

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1	NAG	C1-O5-C5	8.08	123.01	112.19
3	C	1	NAG	C4-C3-C2	3.06	115.50	111.02
3	C	2	NAG	C3-C4-C5	-2.64	105.45	110.23
3	C	3	FUL	C1-O5-C5	2.62	119.14	112.97
3	C	1	NAG	O6-C6-C5	2.27	119.08	111.33
3	C	3	FUL	C2-C3-C4	-2.26	106.88	110.86
3	C	1	NAG	O5-C5-C6	2.17	111.88	107.66
3	C	2	NAG	O4-C4-C3	2.08	115.27	110.38
3	C	3	FUL	O2-C2-C1	2.05	113.92	109.22

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
3	C	1	NAG	O5-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C4-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	2	NAG	1	0
3	C	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry

Of 12 ligands modelled in this entry, 10 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HUW	A	701	-	25,25,25	1.96	2 (8%)	27,37,37	1.71	6 (22%)
6	SO4	A	812	-	4,4,4	0.25	0	6,6,6	0.17	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HUW	A	701	-	-	0/3/23/23	0/5/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	701	HUW	C13-C4	-6.09	1.35	1.42
4	A	701	HUW	C4-N1	5.98	1.39	1.32

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	701	HUW	C13-C4-N1	-3.84	121.97	123.65
4	A	701	HUW	C9-C11-C13	-3.74	107.41	111.33
4	A	701	HUW	C14-C15-C3	-3.31	116.38	118.23
4	A	701	HUW	C4-N1-C3	3.19	121.45	117.67
4	A	701	HUW	C15-C3-N1	-2.43	120.23	122.80
4	A	701	HUW	C2-C3-C15	2.10	121.93	119.58

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	701	HUW	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	563/583 (96%)	-0.82	5 (0%) 81 63	20, 40, 90, 176	17 (3%)
2	B	61/61 (100%)	-0.66	0 100 100	29, 53, 84, 134	0
All	All	624/644 (96%)	-0.81	5 (0%) 82 65	20, 41, 91, 176	17 (2%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	553	TRP	2.2
1	A	564	MET	2.2
1	A	560	TRP	2.2
1	A	567	TRP	2.2
1	A	566	HIS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	FUL	C	3	10/11	0.74	0.13	91,106,111,116	0
3	NAG	C	2	14/15	0.80	0.11	62,83,93,101	0
3	NAG	C	1	14/15	0.85	0.11	88,96,99,99	0

The following is a graphical depiction of the model fit to experimental electron density for oligosac-

charide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CL	A	805	1/1	0.83	0.06	58,58,58,58	0
5	CL	A	808	1/1	0.86	0.12	69,69,69,69	0
5	CL	A	809	1/1	0.90	0.06	79,79,79,79	0
5	CL	A	804	1/1	0.92	0.11	71,71,71,71	0
6	SO4	A	812	5/5	0.92	0.10	72,77,82,96	0
5	CL	A	810	1/1	0.93	0.05	44,44,44,44	0
5	CL	A	806	1/1	0.93	0.06	87,87,87,87	0
5	CL	A	811	1/1	0.94	0.17	67,67,67,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	CL	A	807	1/1	0.95	0.03	53,53,53,53	0
5	CL	A	803	1/1	0.96	0.04	52,52,52,52	0
4	HUW	A	701	22/22	0.98	0.05	20,40,55,58	0
5	CL	A	802	1/1	0.98	0.03	36,36,36,36	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.