



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 18, 2026 – 04:04 PM UTC

PDB ID : 4BDU / pdb\_00004bdu  
Title : Bax BH3-in-Groove dimer (GFP)  
Authors : Czabotar, P.E.; Colman, P.M.  
Deposited on : 2012-10-08  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	 68% 26% . .
1	B	309	 66% 28% . .
1	C	309	 65% 29% . .
1	D	309	 68% 26% . .

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 9548 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GREEN FLUORESCENT PROTEIN, APOPTOSIS REGULATOR BAX.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	296	2366	1506	400	451	9	0	0	0
1	B	296	2366	1506	400	451	9	0	0	0
1	C	296	2366	1506	400	451	9	0	0	0
1	D	296	2366	1506	400	451	9	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP P42212
A	-1	SER	-	expression tag	UNP P42212
A	0	HIS	-	expression tag	UNP P42212
A	65	CR2	SER	engineered mutation	UNP P42212
A	65	CR2	SER	chromophore	UNP P42212
A	65	CR2	TYR	chromophore	UNP P42212
A	65	CR2	GLY	chromophore	UNP P42212
A	206	ASN	ALA	engineered mutation	UNP P42212
A	231	GLY	-	linker	UNP P42212
A	232	SER	-	linker	UNP P42212
A	1126	SER	CYS	engineered mutation	UNP Q07812
B	-2	GLY	-	expression tag	UNP P42212
B	-1	SER	-	expression tag	UNP P42212
B	0	HIS	-	expression tag	UNP P42212
B	65	CR2	SER	engineered mutation	UNP P42212
B	65	CR2	SER	chromophore	UNP P42212
B	65	CR2	TYR	chromophore	UNP P42212
B	65	CR2	GLY	chromophore	UNP P42212
B	206	ASN	ALA	engineered mutation	UNP P42212
B	231	GLY	-	linker	UNP P42212

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	232	SER	-	linker	UNP P42212
B	1126	SER	CYS	engineered mutation	UNP Q07812
C	-2	GLY	-	expression tag	UNP P42212
C	-1	SER	-	expression tag	UNP P42212
C	0	HIS	-	expression tag	UNP P42212
C	65	CR2	SER	engineered mutation	UNP P42212
C	65	CR2	SER	chromophore	UNP P42212
C	65	CR2	TYR	chromophore	UNP P42212
C	65	CR2	GLY	chromophore	UNP P42212
C	206	ASN	ALA	engineered mutation	UNP P42212
C	231	GLY	-	linker	UNP P42212
C	232	SER	-	linker	UNP P42212
C	1126	SER	CYS	engineered mutation	UNP Q07812
D	-2	GLY	-	expression tag	UNP P42212
D	-1	SER	-	expression tag	UNP P42212
D	0	HIS	-	expression tag	UNP P42212
D	65	CR2	SER	engineered mutation	UNP P42212
D	65	CR2	SER	chromophore	UNP P42212
D	65	CR2	TYR	chromophore	UNP P42212
D	65	CR2	GLY	chromophore	UNP P42212
D	206	ASN	ALA	engineered mutation	UNP P42212
D	231	GLY	-	linker	UNP P42212
D	232	SER	-	linker	UNP P42212
D	1126	SER	CYS	engineered mutation	UNP Q07812

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	18	Total O 18 18	0	0
2	A	2	Total O 2 2	0	0
2	B	18	Total O 18 18	0	0
2	B	4	Total O 4 4	0	0
2	C	19	Total O 19 19	0	0
2	C	4	Total O 4 4	0	0
2	D	18	Total O 18 18	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	D	1	Total	O	0	0
			1	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GREEN FLUORESCENT PROTEIN, APOPTOSIS REGULATOR BAX



- Molecule 1: GREEN FLUORESCENT PROTEIN, APOPTOSIS REGULATOR BAX



- Molecule 1: GREEN FLUORESCENT PROTEIN, APOPTOSIS REGULATOR BAX





• Molecule 1: GREEN FLUORESCENT PROTEIN, APOPTOSIS REGULATOR BAX



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 64	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	112.31Å 112.31Å 293.27Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.96 – 3.00 19.96 – 3.00	Depositor EDS
% Data completeness (in resolution range)	99.4 (19.96-3.00) 99.5 (19.96-3.00)	Depositor EDS
$R_{merge}$	0.17	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.37 (at 2.98Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.215 , 0.246 0.209 , 0.238	Depositor DCC
$R_{free}$ test set	2074 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	82.1	Xtrriage
Anisotropy	0.110	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.23 , 53.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.480 for h,-h-k,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	9548	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	120.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 46.24 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.1773e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CR2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.52	2/2397 (0.1%)	0.92	5/3234 (0.2%)
1	B	0.51	0/2397	0.94	6/3234 (0.2%)
1	C	0.51	0/2397	0.94	6/3234 (0.2%)
1	D	0.50	0/2397	0.93	5/3234 (0.2%)
All	All	0.51	2/9588 (0.0%)	0.93	22/12936 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	139	HIS	CE1-NE2	-5.41	1.27	1.32
1	A	64	PHE	C-O	5.34	1.34	1.23

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	195	LEU	CA-C-N	6.52	126.48	120.03
1	A	195	LEU	C-N-CA	6.52	126.48	120.03
1	D	1115	TYR	N-CA-C	-6.35	104.58	112.90
1	C	195	LEU	CA-C-N	6.28	126.19	119.85
1	C	195	LEU	C-N-CA	6.28	126.19	119.85

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2366	0	2299	89	0
1	B	2366	0	2299	102	0
1	C	2366	0	2299	102	0
1	D	2366	0	2299	86	0
2	A	20	0	0	19	0
2	B	22	0	0	29	0
2	C	23	0	0	23	0
2	D	19	0	0	18	0
All	All	9548	0	9196	334	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 334 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:218:MET:HA	2:C:2014:HOH:O	1.33	1.27
1:C:103:ASP:OD1	1:C:104:GLY:N	1.94	0.99
1:B:103:ASP:OD1	1:B:104:GLY:N	1.94	0.99
1:C:134:GLY:HA3	2:C:2010:HOH:O	1.64	0.98
1:C:56:PRO:HG2	1:C:141:LEU:HD12	1.43	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	289/309 (94%)	263 (91%)	21 (7%)	5 (2%)	<b>7</b> 32
1	B	289/309 (94%)	264 (91%)	23 (8%)	2 (1%)	18 <b>53</b>
1	C	289/309 (94%)	264 (91%)	22 (8%)	3 (1%)	12 <b>45</b>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	289/309 (94%)	263 (91%)	21 (7%)	5 (2%)	7	32
All	All	1156/1236 (94%)	1054 (91%)	87 (8%)	15 (1%)	9	38

5 of 15 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	79	LYS
1	B	69	GLN
1	B	79	LYS
1	C	69	GLN
1	C	79	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/268 (96%)	254 (99%)	3 (1%)	63	82
1	B	257/268 (96%)	254 (99%)	3 (1%)	63	82
1	C	257/268 (96%)	255 (99%)	2 (1%)	73	86
1	D	257/268 (96%)	253 (98%)	4 (2%)	55	79
All	All	1028/1072 (96%)	1016 (99%)	12 (1%)	63	82

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	206	ASN
1	D	43	THR
1	D	206	ASN
1	D	123	ILE
1	B	101	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	149	ASN
1	D	25	HIS
1	C	25	HIS
1	B	25	HIS
1	C	1104	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
1	CR2	B	65	1	20,20,21	3.86	7 (35%)	25,27,29	3.04	11 (44%)
1	CR2	A	65	1	20,20,21	4.11	7 (35%)	25,27,29	3.16	11 (44%)
1	CR2	D	65	1	20,20,21	4.08	8 (40%)	25,27,29	3.05	11 (44%)
1	CR2	C	65	1	20,20,21	3.87	8 (40%)	25,27,29	2.88	11 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CR2	B	65	1	-	0/6/25/26	0/2/2/2
1	CR2	A	65	1	-	0/6/25/26	0/2/2/2
1	CR2	D	65	1	-	0/6/25/26	0/2/2/2
1	CR2	C	65	1	-	0/6/25/26	0/2/2/2

The worst 5 of 30 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	65	CR2	CB2-CA2	14.82	1.49	1.35
1	D	65	CR2	CB2-CA2	14.74	1.49	1.35
1	B	65	CR2	CB2-CA2	14.16	1.48	1.35
1	C	65	CR2	CB2-CA2	14.04	1.48	1.35
1	A	65	CR2	CA1-C1	6.42	1.56	1.49

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	65	CR2	O2-C2-CA2	-8.06	125.88	131.02
1	A	65	CR2	O2-C2-CA2	-7.79	126.05	131.02
1	C	65	CR2	O2-C2-CA2	-7.66	126.13	131.02
1	D	65	CR2	CB2-CA2-C2	6.91	130.73	122.36
1	A	65	CR2	CB2-CA2-C2	6.89	130.71	122.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	65	CR2	4	0
1	A	65	CR2	3	0
1	D	65	CR2	1	0
1	C	65	CR2	1	0

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	295/309 (95%)	-0.35	1 (0%) 90 79	67, 107, 196, 264	0
1	B	295/309 (95%)	-0.40	1 (0%) 90 79	67, 107, 196, 264	0
1	C	295/309 (95%)	-0.46	0 100 100	64, 104, 197, 264	0
1	D	295/309 (95%)	-0.39	3 (1%) 79 59	69, 107, 197, 264	0
All	All	1180/1236 (95%)	-0.40	5 (0%) 88 76	64, 107, 197, 264	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	1101	SER	4.3
1	A	1101	SER	2.8
1	B	1098	ASP	2.6
1	D	1056	THR	2.4
1	D	1095	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	CR2	A	65	19/20	0.97	0.07	62,85,99,99	0
1	CR2	B	65	19/20	0.98	0.06	47,86,110,113	0
1	CR2	C	65	19/20	0.98	0.06	47,89,97,100	0
1	CR2	D	65	19/20	0.98	0.07	62,82,100,103	0

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.