



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 03:19 AM UTC

PDB ID : 1BEN / pdb_00001ben
Title : INSULIN COMPLEXED WITH 4-HYDROXYBENZAMIDE
Authors : Smith, G.D.; Ciszak, E.; Pangborn, W.
Deposited on : 1996-02-15
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	21	62% 33% 5%
1	C	21	76% 19% 5%
2	B	30	80% 13% . .
2	D	30	67% 23% . 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 1713 atoms, of which 706 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HUMAN INSULIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	21	Total	C	H	N	O	S	0	1	0
			303	97	140	26	35	5			
1	C	21	Total	C	H	N	O	S	0	2	0
			315	103	144	26	38	4			


- Molecule 2 is a protein called HUMAN INSULIN.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	30	Total	C	H	N	O	S	0	3	0
			473	165	221	43	42	2			
2	D	28	Total	C	H	N	O	S	0	1	0
			431	152	201	37	39	2			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Zn	0	0
			1	1		
3	D	1	Total	Zn	0	0
			1	1		

- Molecule 4 is 4-HYDROXYBENZAMIDE (CCD ID: HBD) (formula: C₇H₇NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	C	1	Total	C	N	O	0	0
			10	7	1	2		
4	D	1	Total	C	N	O	0	0
			10	7	1	2		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total	Cl	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	39	Total	O	0	0
			39	39		
6	B	50	Total	O	0	0
			50	50		
6	C	42	Total	O	0	0
			42	42		
6	D	37	Total	O	0	0
			37	37		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: HUMAN INSULIN

Chain A: 



- Molecule 1: HUMAN INSULIN

Chain C: 



- Molecule 2: HUMAN INSULIN

Chain B: 



- Molecule 2: HUMAN INSULIN

Chain D: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	80.72Å 80.72Å 37.63Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	8.00 – 1.40	Depositor
% Data completeness (in resolution range)	76.5 (8.00-1.40)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROFFT	Depositor
R, R_{free}	(Not available) , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1713	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, HBD, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.79	0/150	1.82	1/199 (0.5%)
1	C	0.94	0/160	1.75	4/213 (1.9%)
2	B	2.77	2/254 (0.8%)	1.85	7/343 (2.0%)
2	D	0.92	0/227	1.73	4/307 (1.3%)
All	All	1.74	2/791 (0.3%)	1.79	16/1062 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	30	THR	CB-OG1	-38.09	0.82	1.43
2	B	30	THR	CB-CG2	-15.32	1.02	1.52

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	27	THR	CA-C-N	7.26	126.89	119.56
2	B	27	THR	C-N-CA	7.26	126.89	119.56
2	B	30	THR	OG1-CB-CG2	7.10	123.51	109.30
2	B	30	THR	CA-CB-CG2	-6.94	98.70	110.50
1	A	2	ILE	N-CA-C	6.52	117.27	110.62
2	B	30	THR	CA-CB-OG1	-6.03	100.55	109.60
1	C	6	CYS	N-CA-C	6.00	120.62	112.88

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	7	CYS	CA-C-N	5.96	126.60	119.98
2	D	7	CYS	C-N-CA	5.96	126.60	119.98
1	C	20	CYS	O-C-N	5.93	129.61	122.85
1	C	20	CYS	CA-C-O	-5.80	114.97	121.81
1	C	5	GLN	CB-CG-CD	5.25	121.53	112.60
2	D	5	HIS	CA-CB-CG	-5.14	108.66	113.80
2	D	17	LEU	CA-C-O	-5.08	115.47	120.70
2	B	12[1]	VAL	O-C-N	-5.03	117.00	121.87
2	B	12[2]	VAL	O-C-N	-5.03	117.00	121.87

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	11[2]	CYS	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	163	140	147	8	0
1	C	171	144	159	2	0
2	B	252	221	251	9	0
2	D	230	201	210	9	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	C	10	0	7	0	0
4	D	10	0	7	1	0
5	D	1	0	0	0	0
6	A	39	0	0	3	0
6	B	50	0	0	2	0
6	C	42	0	0	0	0
6	D	37	0	0	3	0
All	All	1007	706	781	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:13:GLU:OE2	2:D:13:GLU:HG3	1.77	0.85
2:D:21:GLU:CD	2:D:21:GLU:H	2.00	0.67
2:B:13:GLU:OE2	2:D:13:GLU:CG	2.48	0.61
2:B:27:THR:O	2:B:30:THR:HG22	2.01	0.60
1:A:4:GLU:OE1	6:A:25:HOH:O	2.17	0.59
6:A:24:HOH:O	2:B:30:THR:HG21	2.04	0.57
2:B:27:THR:O	2:B:30:THR:CG2	2.54	0.55
1:A:5:GLN:OE1	6:A:32:HOH:O	2.18	0.54
6:B:79:HOH:O	2:D:21:GLU:CB	2.55	0.54
2:D:21:GLU:CD	2:D:21:GLU:N	2.65	0.54
2:D:22:ARG:NH2	6:D:112:HOH:O	2.41	0.53
1:A:11[2]:CYS:HB2	1:A:15:GLN:HB2	1.91	0.53
2:B:30:THR:HG1	2:B:30:THR:C	2.05	0.52
1:C:5:GLN:OE1	1:C:19:TYR:OH	2.12	0.52
6:B:79:HOH:O	2:D:21:GLU:HG2	2.13	0.48
1:A:13:LEU:HD11	2:B:18[2]:VAL:CG2	2.43	0.47
1:A:13:LEU:HD11	2:B:18[2]:VAL:HG23	1.99	0.45
1:A:4:GLU:H	1:A:4:GLU:CD	2.26	0.44
1:C:13:LEU:HD23	1:C:13:LEU:C	2.42	0.44
4:D:102:HBD:C1'	6:D:111:HOH:O	2.68	0.41
1:A:1:GLY:HA3	2:B:30:THR:HG23	2.04	0.40
2:D:28:PRO:HB3	6:D:122:HOH:O	2.20	0.40
1:A:21:ASN:ND2	2:D:27:THR:HG23	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	20/21 (95%)	19 (95%)	1 (5%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	21/21 (100%)	21 (100%)	0	0	100	100
2	B	31/30 (103%)	30 (97%)	1 (3%)	0	100	100
2	D	27/30 (90%)	27 (100%)	0	0	100	100
All	All	99/102 (97%)	97 (98%)	2 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	18/20 (90%)	18 (100%)	0	100	100
1	C	18/20 (90%)	18 (100%)	0	100	100
2	B	23/26 (88%)	21 (91%)	2 (9%)	9	0
2	D	22/26 (85%)	21 (96%)	1 (4%)	24	4
All	All	81/92 (88%)	78 (96%)	3 (4%)	30	5

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	29	LYS
2	B	30	THR
2	D	27	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
2	B	5	HIS
1	C	21	ASN
2	D	4	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HBD	D	102	-	10,10,10	1.20	1 (10%)	13,13,13	1.73	2 (15%)
4	HBD	C	101	-	10,10,10	1.00	0	13,13,13	2.39	4 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HBD	D	102	-	-	0/4/4/4	0/1/1/1
4	HBD	C	101	-	-	0/4/4/4	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	102	HBD	C1'-N1'	2.40	1.37	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	C	101	HBD	O1'-C1'-N1'	-4.89	115.55	122.62
4	D	102	HBD	O1'-C1'-C1	4.80	125.47	119.60
4	C	101	HBD	C1-C1'-N1'	4.68	123.51	117.74
4	C	101	HBD	C5-C6-C1	-3.52	117.04	120.80
4	D	102	HBD	C1-C1'-N1'	-2.69	114.42	117.74
4	C	101	HBD	C6-C1-C2	2.52	121.77	118.57

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	102	HBD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.