



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 04:44 AM UTC

PDB ID : 2BEO / pdb_00002beo
Title : PrfA, Transcriptional Regulator In Listeria Monocytogenes
Authors : Eiting, M.; Hagelueken, G.; Schubert, W.-D.; Heinz, D.W.
Deposited on : 2004-11-26
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3538 (2.70-2.70)
Clashscore	190562	3843 (2.70-2.70)
Ramachandran outliers	187476	3778 (2.70-2.70)
Sidechain outliers	187428	3778 (2.70-2.70)
RSRZ outliers	180081	3538 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	236	 74% 17% . . .
1	B	236	 77% 16% . . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACM	B	1239	-	-	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 4031 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LISTERIOLYSIN REGULATORY PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	16	0
			1985	1294	314	369	8			
1	B	231	Total	C	N	O	S	0	10	0
			1980	1291	311	371	7			

- Molecule 2 is GLUTAMINE (CCD ID: GLN) (formula: C₅H₁₀N₂O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			10	5	2	3		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Cl	0	0
			1	1		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cl	0	0
			1	1		

- Molecule 4 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			11	7	4		

- Molecule 5 is ACETAMIDE (CCD ID: ACM) (formula: C_2H_5NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			4	2	1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	26	Total	O	0	0
			26	26		
6	B	13	Total	O	0	0
			13	13		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: LISTERIOLYSIN REGULATORY PROTEIN



• Molecule 1: LISTERIOLYSIN REGULATORY PROTEIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	55.95Å 80.71Å 60.97Å 90.00° 111.91° 90.00°	Depositor
Resolution (Å)	56.80 – 2.70 56.56 – 2.70	Depositor EDS
% Data completeness (in resolution range)	91.9 (56.80-2.70) 89.0 (56.56-2.70)	Depositor EDS
R_{merge}	0.01	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.36 (at 2.72Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.244 , 0.289 0.244 , 0.296	Depositor DCC
R_{free} test set	602 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å ²)	46.6	Xtriage
Anisotropy	0.753	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 57.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	4031	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ACM, CL, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.90	1/2028 (0.0%)	1.24	22/2736 (0.8%)
1	B	0.88	3/2025 (0.1%)	1.40	19/2731 (0.7%)
All	All	0.89	4/4053 (0.1%)	1.32	41/5467 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	5
All	All	0	7

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	203	ASN	N-CA	-6.14	1.34	1.46
1	B	149	ILE	CA-CB	6.06	1.62	1.54
1	B	158	THR	CA-CB	5.32	1.61	1.53
1	B	158	THR	CA-C	5.04	1.57	1.52

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	202	LYS	O-C-N	-28.25	94.21	122.18
1	B	202	LYS	CA-C-O	-24.28	99.25	121.11
1	A	203	ASN	CA-C-N	16.22	150.89	121.70
1	A	203	ASN	C-N-CA	16.22	150.89	121.70
1	A	202	LYS	O-C-N	-13.65	105.50	123.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	158	THR	CB-CA-C	11.51	127.82	109.97
1	B	3	ALA	N-CA-C	11.19	127.57	112.68
1	A	202	LYS	N-CA-C	-11.00	93.09	110.14
1	A	65	GLY	CA-C-N	10.43	140.47	121.70
1	A	65	GLY	C-N-CA	10.43	140.47	121.70
1	B	203	ASN	CA-C-N	9.87	137.19	122.68
1	B	203	ASN	C-N-CA	9.87	137.19	122.68
1	A	203	ASN	O-C-N	-9.52	107.77	123.00
1	B	202	LYS	CB-CA-C	-7.83	98.40	111.24
1	A	187	SER	N-CA-C	-7.59	104.01	113.20
1	B	175	GLY	N-CA-C	7.28	130.43	113.18
1	B	158	THR	CA-C-N	7.17	128.80	119.84
1	B	158	THR	C-N-CA	7.17	128.80	119.84
1	B	176	TYR	N-CA-C	-7.11	102.73	111.40
1	A	202	LYS	CA-C-O	-6.68	113.47	121.56
1	B	202	LYS	CA-C-N	-6.49	109.14	121.54
1	B	202	LYS	C-N-CA	-6.49	109.14	121.54
1	B	198	VAL	N-CA-C	-6.42	106.28	111.81
1	A	65	GLY	O-C-N	-6.33	114.47	122.70
1	A	186	VAL	N-CA-CB	6.23	121.50	111.23
1	B	113	PHE	N-CA-C	-6.20	104.44	111.07
1	A	113	PHE	N-CA-C	-6.14	104.47	111.10
1	B	186	VAL	N-CA-C	-5.84	105.04	110.53
1	A	158	THR	CA-C-N	5.73	127.01	119.84
1	A	158	THR	C-N-CA	5.73	127.01	119.84
1	B	31	GLN	CB-CA-C	-5.72	99.03	110.42
1	A	74[A]	ILE	N-CA-C	5.65	116.38	110.62
1	A	74[B]	ILE	N-CA-C	5.65	116.38	110.62
1	B	158	THR	N-CA-CB	5.40	118.86	111.25
1	B	110	LEU	N-CA-C	5.40	118.08	111.82
1	A	110	LEU	N-CA-C	5.34	118.01	111.82
1	A	204	SER	N-CA-CB	-5.33	101.43	110.50
1	A	187	SER	CA-C-N	5.32	127.35	120.44
1	A	187	SER	C-N-CA	5.32	127.35	120.44
1	A	202	LYS	CA-C-N	5.05	130.78	121.70
1	A	202	LYS	C-N-CA	5.05	130.78	121.70

There are no chirality outliers.

All (7) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	202	LYS	Mainchain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	A	203	ASN	Mainchain
1	B	202	LYS	Peptide,Mainchain
1	B	203	ASN	Peptide
1	B	65	GLY	Peptide,Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1985	0	1984	27	0
1	B	1980	0	1959	26	0
2	A	10	0	7	0	0
3	A	1	0	0	0	0
3	B	1	0	0	1	0
4	A	11	0	13	0	0
5	B	4	0	5	3	0
6	A	26	0	0	0	0
6	B	13	0	0	1	0
All	All	4031	0	3968	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118[A]:GLN:OE1	1:B:75[A]:ASP:OD2	1.78	0.99
1:B:74:ILE:O	1:B:103:LYS:HE2	1.87	0.73
1:B:202:LYS:HE2	1:B:203:ASN:HD21	1.66	0.59
1:A:67:PHE:CE1	1:A:119[B]:THR:HG23	2.38	0.58
1:A:70:MET:HE2	1:A:102:LEU:CD2	2.33	0.58
1:A:31:GLN:NE2	1:A:84:ASN:HD21	2.02	0.57
1:B:199:ILE:HD12	1:B:206[A]:PHE:CZ	2.40	0.56
1:A:205:CYS:SG	1:A:206:PHE:N	2.78	0.56
1:A:118[A]:GLN:HG2	1:A:119[A]:THR:N	2.21	0.56
1:B:103:LYS:NZ	6:B:2007:HOH:O	2.39	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:169:LEU:O	1:A:172:GLN:HG3	2.07	0.55
1:B:126:TYR:CD2	5:B:1239:ACM:H22	2.42	0.54
1:B:158:THR:HB	1:B:159:PRO:HD3	1.89	0.54
1:A:202:LYS:CE	1:A:203:ASN:HA	2.40	0.52
1:A:97:ILE:HD11	1:A:102:LEU:HB2	1.92	0.51
1:A:31:GLN:HE22	1:A:51:ILE:HB	1.76	0.51
1:A:110:LEU:HD11	1:B:110:LEU:HD11	1.94	0.50
1:A:41:LEU:O	1:A:66:ALA:HA	2.11	0.50
1:A:68:VAL:HG13	1:A:119[B]:THR:HG21	1.94	0.49
1:A:118[A]:GLN:CG	1:A:119[A]:THR:N	2.76	0.48
1:A:185:ALA:O	1:A:186:VAL:HG22	2.14	0.47
1:B:61:GLN:NE2	3:B:1238:CL:CL	2.85	0.47
1:A:31:GLN:NE2	1:A:51:ILE:HB	2.30	0.46
1:A:149[B]:ILE:HD11	1:A:221:LEU:HD22	1.98	0.46
1:B:11:TYR:O	1:B:15:ASN:ND2	2.47	0.45
1:B:194:LYS:HD3	1:B:201[B]:TYR:CD1	2.51	0.45
1:B:164:ILE:HG12	1:B:206[B]:PHE:CZ	2.51	0.45
1:A:10:LYS:O	1:A:14:THR:HG23	2.17	0.45
1:B:158:THR:HB	1:B:159:PRO:CD	2.47	0.45
1:A:163:LYS:HG2	1:A:205:CYS:SG	2.58	0.44
1:B:145:GLY:O	1:B:149:ILE:HG23	2.18	0.43
1:A:202:LYS:HE3	1:A:203:ASN:HA	2.00	0.43
1:B:61:GLN:NE2	5:B:1239:ACM:H21	2.34	0.43
1:B:194:LYS:HD3	1:B:201[B]:TYR:HD1	1.83	0.43
1:B:174:LEU:O	1:B:174:LEU:HD23	2.18	0.42
1:A:198:VAL:HA	1:A:210:ASN:HB3	2.01	0.42
1:B:48:LEU:HD22	1:B:83:TYR:HB3	2.02	0.42
1:B:199:ILE:HB	1:B:206[A]:PHE:CE1	2.55	0.42
1:A:186:VAL:HG23	1:A:189:ILE:HD12	2.02	0.41
1:B:18:LYS:HG3	1:B:19:PRO:HD2	2.01	0.41
1:B:158:THR:CB	1:B:159:PRO:HD3	2.50	0.41
1:A:97:ILE:HD11	1:A:102:LEU:CB	2.50	0.41
1:B:123:GLN:HG2	5:B:1239:ACM:H23	2.02	0.41
1:A:128:LEU:HD21	1:B:61:GLN:HB2	2.03	0.41
1:A:113:PHE:HE1	1:B:117:PHE:CD1	2.39	0.40
1:A:117:PHE:CD2	1:B:74:ILE:CG1	3.05	0.40
1:A:74[B]:ILE:CG1	1:B:117:PHE:CD2	3.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	238/236 (101%)	228 (96%)	7 (3%)	3 (1%)	9	25
1	B	237/236 (100%)	224 (94%)	10 (4%)	3 (1%)	9	25
All	All	475/472 (101%)	452 (95%)	17 (4%)	6 (1%)	9	25

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	168	ASN
1	A	186	VAL
1	B	175	GLY
1	B	66	ALA
1	B	65	GLY
1	A	159	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	220/211 (104%)	196 (89%)	24 (11%)	6	16
1	B	218/211 (103%)	191 (88%)	27 (12%)	4	11
All	All	438/422 (104%)	387 (88%)	51 (12%)	6	13

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LYS
1	A	18	LYS
1	A	48	LEU
1	A	53[A]	GLU
1	A	53[B]	GLU
1	A	54[A]	ASN
1	A	54[B]	ASN
1	A	68	VAL
1	A	97	ILE
1	A	110	LEU
1	A	120	LEU
1	A	122	LYS
1	A	123	GLN
1	A	127	SER
1	A	150	LEU
1	A	156	LYS
1	A	158	THR
1	A	172	GLN
1	A	188	ARG
1	A	200	VAL
1	A	202	LYS
1	A	203	ASN
1	A	206	PHE
1	A	221	LEU
1	B	9	LYS
1	B	18	LYS
1	B	30	ASN
1	B	48	LEU
1	B	74	ILE
1	B	88	ILE
1	B	99	ILE
1	B	110	LEU
1	B	120	LEU
1	B	122[A]	LYS
1	B	122[B]	LYS
1	B	123	GLN
1	B	127	SER
1	B	139[A]	LYS
1	B	139[B]	LYS
1	B	140	LEU
1	B	150	LEU
1	B	158	THR
1	B	166	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	168[A]	ASN
1	B	168[B]	ASN
1	B	172	GLN
1	B	176	TYR
1	B	204	SER
1	B	206[A]	PHE
1	B	206[B]	PHE
1	B	221	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	ASN
1	A	31	GLN
1	A	35	GLN
1	A	123	GLN
1	A	137	ASN
1	A	172	GLN
1	A	209	GLN
1	B	30	ASN
1	B	61	GLN
1	B	91	GLN
1	B	203	ASN
1	B	209	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 5 ligands modelled in this entry, 2 are monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	ACM	B	1239	-	3,3,3	0.43	0	3,3,3	0.24	0
2	GLN	A	1240	-	8,9,9	0.84	1 (12%)	8,11,11	0.95	1 (12%)
4	PG4	A	1239	-	10,10,12	0.52	0	9,9,11	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLN	A	1240	-	-	1/9/9/9	-
4	PG4	A	1239	-	-	1/8/8/10	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1240	GLN	OXT-C	-2.03	1.24	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1240	GLN	OXT-C-O	-2.54	118.31	124.08

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1239	PG4	O2-C3-C4-O3
2	A	1240	GLN	O-C-CA-N

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	1239	ACM	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	226/236 (95%)	-0.04	2 (0%) 81 80	6, 21, 40, 56	16 (7%)
1	B	231/236 (97%)	-0.05	1 (0%) 88 87	4, 21, 42, 78	10 (4%)
All	All	457/472 (96%)	-0.04	3 (0%) 84 83	4, 21, 41, 78	26 (5%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	189	ILE	2.4
1	A	160	ASP	2.1
1	B	206[A]	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CL	B	1238	1/1	0.79	0.10	49,49,49,49	1
4	PG4	A	1239	11/13	0.79	0.19	68,75,86,92	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GLN	A	1240	10/10	0.81	0.13	48,55,62,66	0
3	CL	A	1238	1/1	0.87	0.06	61,61,61,61	0
5	ACM	B	1239	4/4	0.92	0.07	43,46,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.