



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:42 PM UTC

PDB ID : 2BEQ / pdb_00002beq
Title : Structure of a Proteolytically Resistant Core from the Severe Acute Respiratory Syndrome Coronavirus S2 Fusion Protein
Authors : Supekar, V.M.; Bruckmann, C.; Ingallinella, P.; Bianchi, E.; Pessi, A.; Carfi, A.
Deposited on : 2004-11-29
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4673 (1.60-1.60)
Clashscore	190562	4931 (1.60-1.60)
Ramachandran outliers	187476	4831 (1.60-1.60)
Sidechain outliers	187428	4830 (1.60-1.60)
RSRZ outliers	180081	4672 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	38	 87% 8% . .
1	B	38	 89% 11%
1	C	38	 95% . .
2	D	48	 4% 98% .
2	E	48	 4% 83% 15% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	48	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '8%', a large green segment in the middle labeled '88%', and a yellow segment on the right labeled '12%'.</p>

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2110 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	37	270	163	48	59	0	1	1
1	B	38	272	165	48	59	0	0	1
1	C	38	273	165	48	60	0	1	1

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	913	ACE	-	expression tag	UNP P59594
A	950	NH2	-	expression tag	UNP P59594
B	913	ACE	-	expression tag	UNP P59594
B	950	NH2	-	expression tag	UNP P59594
C	913	ACE	-	expression tag	UNP P59594
C	950	NH2	-	expression tag	UNP P59594

- Molecule 2 is a protein called Spike glycoprotein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	48	370	230	62	78	0	1	1
2	E	47	359	223	61	75	0	0	2
2	F	48	364	227	61	76	0	0	1

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	1147	ACE	-	expression tag	UNP P59594
D	1194	NH2	-	expression tag	UNP P59594

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	1147	ACE	-	expression tag	UNP P59594
E	1194	NH2	-	expression tag	UNP P59594
F	1147	ACE	-	expression tag	UNP P59594
F	1194	NH2	-	expression tag	UNP P59594


- Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	24	Total O 24 24	0	0
3	B	28	Total O 28 28	0	0
3	C	23	Total O 23 23	0	0
3	D	46	Total O 46 46	0	0
3	E	47	Total O 47 47	0	0
3	F	34	Total O 34 34	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike glycoprotein

Chain A:  87% 8% . .




- Molecule 1: Spike glycoprotein

Chain B:  89% 11%




- Molecule 1: Spike glycoprotein

Chain C:  95% . .




- Molecule 2: Spike glycoprotein

Chain D:  4% 98% .





- Molecule 2: Spike glycoprotein

Chain E:  4% 83% 15% .



- Molecule 2: Spike glycoprotein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	99.91Å 42.90Å 59.83Å 90.00° 116.69° 90.00°	Depositor
Resolution (Å)	30.00 – 1.60 30.00 – 1.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-1.60) 99.6 (30.00-1.60)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.26 (at 1.60Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.199 , 0.241 0.207 , 0.247	Depositor DCC
R_{free} test set	1521 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtrriage
Anisotropy	0.583	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	2110	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.86% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.01	0/273	1.02	0/370
1	B	1.04	0/268	1.02	0/364
1	C	0.99	0/274	1.05	0/372
2	D	0.88	0/374	1.01	0/503
2	E	1.05	0/359	1.05	0/481
2	F	1.00	0/362	1.13	1/488 (0.2%)
All	All	0.99	0/1910	1.05	1/2578 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1188	TYR	N-CA-C	5.76	118.02	111.11

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	270	0	279	1	0
1	B	272	0	282	3	0
1	C	273	0	283	2	0
2	D	370	0	371	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	E	359	0	361	9	0
2	F	364	0	364	0	0
3	A	24	0	0	0	0
3	B	28	0	0	0	0
3	C	23	0	0	0	0
3	D	46	0	0	0	0
3	E	47	0	0	0	0
3	F	34	0	0	0	0
All	All	2110	0	1940	10	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 10 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:916:ILE:HD11	2:E:1179:LEU:HD11	1.45	0.99
2:E:1149:GLY:C	2:E:1151:ILE:H	1.92	0.77
2:E:1149:GLY:C	2:E:1151:ILE:N	2.61	0.57
1:B:916:ILE:CD1	2:E:1179:LEU:HD11	2.26	0.57
2:E:1149:GLY:HA2	2:E:1151:ILE:HG12	1.87	0.56

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	36/38 (95%)	36 (100%)	0	0	100 100
1	B	36/38 (95%)	36 (100%)	0	0	100 100
1	C	37/38 (97%)	37 (100%)	0	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	D	47/48 (98%)	45 (96%)	2 (4%)	0	100	100
2	E	45/48 (94%)	43 (96%)	2 (4%)	0	100	100
2	F	46/48 (96%)	46 (100%)	0	0	100	100
All	All	247/258 (96%)	243 (98%)	4 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	33/32 (103%)	29 (88%)	4 (12%)	5	0
1	B	32/32 (100%)	29 (91%)	3 (9%)	8	1
1	C	33/32 (103%)	30 (91%)	3 (9%)	9	1
2	D	42/41 (102%)	41 (98%)	1 (2%)	43	19
2	E	40/41 (98%)	38 (95%)	2 (5%)	22	5
2	F	40/41 (98%)	35 (88%)	5 (12%)	4	0
All	All	220/219 (100%)	202 (92%)	18 (8%)	11	2

5 of 18 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	1175	LEU
2	F	1192	ILE
2	F	1190	GLN
1	C	924[B]	SER
2	F	1163	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	1176	ASN
2	E	1190	GLN
2	F	1169	ASN
1	C	936	GLN
1	C	939	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	36/38 (94%)	0.17	0	100 100	23, 29, 43, 44	1 (2%)
1	B	36/38 (94%)	-0.05	0	100 100	21, 27, 37, 42	0
1	C	36/38 (94%)	-0.16	0	100 100	20, 26, 37, 39	1 (2%)
2	D	46/48 (95%)	0.50	2 (4%)	40 41	26, 36, 46, 48	1 (2%)
2	E	46/48 (95%)	0.23	2 (4%)	40 41	23, 31, 42, 50	0
2	F	46/48 (95%)	0.67	4 (8%)	16 16	26, 39, 50, 53	0
All	All	246/258 (95%)	0.25	8 (3%)	49 52	20, 32, 46, 53	3 (1%)

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	1148	LEU	3.6
2	F	1191	TYR	2.9
2	E	1149	GLY	2.5
2	D	1182	LEU	2.3
2	F	1177	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.