



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:27 AM UTC

PDB ID : 3BEX / pdb_00003bex
Title : Type III pantothenate kinase from *Thermotoga maritima* complexed with pantothenate
Authors : Yang, K.; Huerta, C.; Strauss, E.; Zhang, H.
Deposited on : 2007-11-20
Resolution : 1.51 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5890 (1.54-1.50)
Clashscore	190562	6116 (1.54-1.50)
Ramachandran outliers	187476	6002 (1.54-1.50)
Sidechain outliers	187428	5999 (1.54-1.50)
RSRZ outliers	180081	5891 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	249	
1	B	249	
1	C	249	
1	D	249	
1	E	249	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	249	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a small red segment on the left labeled '4%', a large green segment in the middle labeled '88%', and a small yellow segment on the right labeled '10%'. A small black dot is located at the end of the bar.</p>

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 13501 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type III pantothenate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	248	1929	1244	323	353	9	0	0	0
1	B	248	1929	1244	323	353	9	0	0	0
1	C	248	1929	1244	323	353	9	0	0	0
1	D	248	1929	1244	323	353	9	0	0	0
1	E	248	1929	1244	323	353	9	0	0	0
1	F	248	1929	1244	323	353	9	0	0	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	MET	-	expression tag	UNP Q9WZY5
A	-1	ASP	-	expression tag	UNP Q9WZY5
A	0	PRO	-	expression tag	UNP Q9WZY5
B	-2	MET	-	expression tag	UNP Q9WZY5
B	-1	ASP	-	expression tag	UNP Q9WZY5
B	0	PRO	-	expression tag	UNP Q9WZY5
C	-2	MET	-	expression tag	UNP Q9WZY5
C	-1	ASP	-	expression tag	UNP Q9WZY5
C	0	PRO	-	expression tag	UNP Q9WZY5
D	-2	MET	-	expression tag	UNP Q9WZY5
D	-1	ASP	-	expression tag	UNP Q9WZY5
D	0	PRO	-	expression tag	UNP Q9WZY5
E	-2	MET	-	expression tag	UNP Q9WZY5
E	-1	ASP	-	expression tag	UNP Q9WZY5
E	0	PRO	-	expression tag	UNP Q9WZY5
F	-2	MET	-	expression tag	UNP Q9WZY5
F	-1	ASP	-	expression tag	UNP Q9WZY5

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
F	0	PRO	-	expression tag	UNP Q9WZY5

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is PANTOTHENOIC ACID (CCD ID: PAU) (formula: C₉H₁₇NO₅).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	15	9	1	5	0	0
3	B	1	15	9	1	5	0	0
3	C	1	15	9	1	5	0	0
3	D	1	15	9	1	5	0	0
3	E	1	15	9	1	5	0	0
3	F	1	15	9	1	5	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	292	292	292	0	0
4	B	293	293	293	0	0
4	C	331	331	331	0	0
4	D	305	305	305	0	0
4	E	296	296	296	0	0
4	F	290	290	290	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Type III pantothenate kinase




- Molecule 1: Type III pantothenate kinase




- Molecule 1: Type III pantothenate kinase



- Molecule 1: Type III pantothenate kinase






- Molecule 1: Type III pantothenate kinase





- Molecule 1: Type III pantothenate kinase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	74.92Å 136.99Å 74.97Å 90.00° 109.62° 90.00°	Depositor
Resolution (Å)	50.00 – 1.51 50.00 – 1.51	Depositor EDS
% Data completeness (in resolution range)	99.4 (50.00-1.51) 99.4 (50.00-1.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 1.51Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.184 , 0.213 (Not available) , 0.217	Depositor DCC
R_{free} test set	11057 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	19.0	Xtrriage
Anisotropy	0.065	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.146 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13501	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.76% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PAU, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.52	0/1971	0.76	1/2668 (0.0%)
1	B	0.53	0/1971	0.73	0/2668
1	C	0.55	0/1971	0.77	0/2668
1	D	0.55	0/1971	0.74	0/2668
1	E	0.54	0/1971	0.76	0/2668
1	F	0.58	0/1971	0.77	0/2668
All	All	0.55	0/11826	0.75	1/16008 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	LEU	CA-CB-CG	5.16	134.36	116.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1929	0	1946	26	0
1	B	1929	0	1946	37	0
1	C	1929	0	1946	27	0
1	D	1929	0	1946	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1929	0	1946	42	0
1	F	1929	0	1946	35	0
2	A	5	0	0	1	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
3	A	15	0	15	0	0
3	B	15	0	15	0	0
3	C	15	0	15	0	0
3	D	15	0	15	0	0
3	E	15	0	16	0	0
3	F	15	0	16	0	0
4	A	292	0	0	8	0
4	B	293	0	0	10	0
4	C	331	0	0	8	0
4	D	305	0	0	7	0
4	E	296	0	0	9	0
4	F	290	0	0	5	0
All	All	13501	0	11768	161	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 161 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:124:ILE:HD11	1:B:199:ILE:HD11	1.33	1.05
1:E:124:ILE:HD11	1:E:199:ILE:HD11	1.33	1.02
1:B:98:ASN:HB2	4:B:527:HOH:O	1.64	0.96
1:B:171:ALA:H	1:B:189:ASN:HD21	1.14	0.93
1:C:69:VAL:HG23	4:C:578:HOH:O	1.65	0.93

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/249 (99%)	241 (98%)	4 (2%)	1 (0%)	30	12
1	B	246/249 (99%)	243 (99%)	2 (1%)	1 (0%)	30	12
1	C	246/249 (99%)	240 (98%)	6 (2%)	0	100	100
1	D	246/249 (99%)	243 (99%)	3 (1%)	0	100	100
1	E	246/249 (99%)	241 (98%)	4 (2%)	1 (0%)	30	12
1	F	246/249 (99%)	244 (99%)	2 (1%)	0	100	100
All	All	1476/1494 (99%)	1452 (98%)	21 (1%)	3 (0%)	43	23

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	31	GLY
1	B	31	GLY
1	E	90	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	211/212 (100%)	208 (99%)	3 (1%)	59	31
1	B	211/212 (100%)	208 (99%)	3 (1%)	59	31
1	C	211/212 (100%)	208 (99%)	3 (1%)	59	31
1	D	211/212 (100%)	205 (97%)	6 (3%)	38	10

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	211/212 (100%)	208 (99%)	3 (1%)	59	31
1	F	211/212 (100%)	207 (98%)	4 (2%)	50	21
All	All	1266/1272 (100%)	1244 (98%)	22 (2%)	53	24

5 of 22 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	225	ILE
1	E	236	ILE
1	E	225	ILE
1	F	-1	ASP
1	C	-2	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	182	ASN
1	F	182	ASN
1	E	66	GLN
1	F	66	GLN
1	E	11	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	PAU	C	248	-	12,14,14	2.18	4 (33%)	17,19,19	1.47	3 (17%)
3	PAU	D	248	-	12,14,14	1.68	2 (16%)	17,19,19	1.26	2 (11%)
2	PO4	A	247	-	4,4,4	1.27	1 (25%)	6,6,6	1.51	1 (16%)
3	PAU	E	248	-	12,14,14	1.65	3 (25%)	17,19,19	1.21	2 (11%)
2	PO4	D	247	-	4,4,4	1.27	1 (25%)	6,6,6	1.62	2 (33%)
2	PO4	C	247	-	4,4,4	1.50	1 (25%)	6,6,6	1.47	1 (16%)
3	PAU	F	248	-	12,14,14	1.73	4 (33%)	17,19,19	1.15	2 (11%)
2	PO4	B	247	-	4,4,4	1.24	0	6,6,6	1.14	1 (16%)
3	PAU	B	248	-	12,14,14	1.88	2 (16%)	17,19,19	1.24	1 (5%)
2	PO4	E	247	-	4,4,4	1.27	0	6,6,6	1.47	1 (16%)
2	PO4	F	247	-	4,4,4	1.55	1 (25%)	6,6,6	0.91	0
3	PAU	A	248	-	12,14,14	1.99	4 (33%)	17,19,19	1.26	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PAU	D	248	-	-	2/19/19/19	-
3	PAU	E	248	-	-	2/19/19/19	-
3	PAU	F	248	-	-	2/19/19/19	-
3	PAU	B	248	-	-	2/19/19/19	-
3	PAU	C	248	-	-	2/19/19/19	-
3	PAU	A	248	-	-	2/19/19/19	-

The worst 5 of 23 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	248	PAU	C6-N	-4.17	1.24	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	248	PAU	C2-C1	-4.01	1.51	1.53
3	D	248	PAU	C6-N	-3.99	1.24	1.33
3	C	248	PAU	O6'-C6	3.96	1.31	1.23
3	C	248	PAU	C6-N	-3.75	1.24	1.33

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	248	PAU	C8-N-C6	3.25	128.38	122.55
3	D	248	PAU	C8-N-C6	3.05	128.03	122.55
3	C	248	PAU	O-C-C9	-3.03	113.48	123.09
2	D	247	PO4	O2-P-O1	-2.95	100.51	110.95
2	C	247	PO4	O3-P-O1	-2.87	100.79	110.95

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	248	PAU	O-C-C9-C8
3	B	248	PAU	OXT-C-C9-C8
3	A	248	PAU	OXT-C-C9-C8
3	F	248	PAU	OXT-C-C9-C8
3	C	248	PAU	OXT-C-C9-C8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	247	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	248/249 (99%)	0.22	7 (2%) 55 61	11, 18, 29, 35	0
1	B	248/249 (99%)	0.28	11 (4%) 39 45	12, 19, 29, 34	0
1	C	248/249 (99%)	0.08	9 (3%) 46 53	11, 17, 27, 34	0
1	D	248/249 (99%)	0.11	8 (3%) 50 56	11, 17, 27, 34	0
1	E	248/249 (99%)	0.22	12 (4%) 35 41	11, 18, 28, 44	0
1	F	248/249 (99%)	0.18	10 (4%) 42 49	11, 17, 29, 50	0
All	All	1488/1494 (99%)	0.18	57 (3%) 44 51	11, 18, 29, 50	0

The worst 5 of 57 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	32	VAL	8.2
1	E	91	CYS	6.8
1	D	33	PHE	6.0
1	C	31	GLY	6.0
1	F	31	GLY	5.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	C	247	5/5	0.81	0.20	20,23,26,26	0
2	PO4	A	247	5/5	0.84	0.20	21,25,27,28	0
2	PO4	B	247	5/5	0.86	0.16	19,21,24,26	0
2	PO4	D	247	5/5	0.89	0.16	19,22,23,25	0
2	PO4	F	247	5/5	0.89	0.15	20,20,23,25	0
2	PO4	E	247	5/5	0.90	0.16	20,22,24,26	0
3	PAU	A	248	15/15	0.96	0.05	13,13,15,16	0
3	PAU	B	248	15/15	0.96	0.05	12,13,15,16	0
3	PAU	C	248	15/15	0.96	0.06	12,13,15,15	0
3	PAU	E	248	15/15	0.96	0.05	11,13,15,16	0
3	PAU	F	248	15/15	0.96	0.06	12,13,13,15	0
3	PAU	D	248	15/15	0.97	0.04	10,13,15,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.